

		Depression	Hypomania	Mania	Psychosis
Depression	<i>Case</i>		14.4% [12 17.2]	6.4% [4.8 8.6]	12.3% [10.0 15.0]
	<i>Control</i>		3.5% [2.8 4.4]	0.5% [0.2 0.9]	5.3% [4.3 6.3]
	<i>p-value</i>		1.40E-24	9.90E-22	4.10E-10
Hypomania	<i>Case</i>	58.9% [51.2 66.2]		% [0 NA]*	18.9% [13.5 25.6]
	<i>Control</i>	23.6% [22 25.3]		2.2% [1.6 2.8]	6.3% [5.4 7.3]
	<i>p-value</i>	1.40E-24			8.80E-10
Mania	<i>Case</i>	82.1% [69.2 90.7]	% [0 NA]*		21.4% [12 34.8]
	<i>Control</i>	24.6% [23 26.3]	6.4% [5.6 7.5]		6.8% [5.9 7.8]
	<i>p-value</i>	9.90E-22			7.20E-05
Psychosis	<i>Case</i>	44.9% [37.9 52.1]	16.8% [12 23]	6.1% [3.3 10.7]	
	<i>Control</i>	24.4% [22.7 26.1]	5.5% [4.7 6.5]	1.7% [1.3 2.3]	
	<i>p-value</i>	4.10E-10	8.80E-10	7.20E-05	

Supplementary 3: Comorbidity pattern between the clinical syndromes

* Mania and Hypomania are mutually exclusive.

NB. Read rows first followed by columns, e.g. of participants with MDD 14.4% also had Hypomania