Table S1. Estimates of Fixed Effects From Mixed Models Testing Treatment Effects Over Course of Follow-Up for Acute Treatment Responders^a

	β	SE	p
CYBOCS	·		-
Group	.05	1.29	.97
Baseline to follow-up	-15.79	1.06	<.001
Posttreatment to follow-up	-0.88	0.99	.38
COIS-RP			
Group	1.49	3.93	.71
Baseline to follow-up	-9.25	2.28	<.001
Posttreatment to follow-up	-0.57	1.72	.74
<u>FAS</u>			
Group	1.06	2.20	.63
Baseline to follow-up	-18.85	2.30	<.001
Posttreatment to follow-up	-0.46	1.29	.72
FES-Conflict			
Group	0.35	0.66	.60
Baseline to follow-up	-0.89	0.28	.003
Posttreatment to follow-up	0.09	0.16	.60
FES-Cohesion			
Group	.07	0.57	.90
Baseline to follow-up	0.47	0.27	.09
Posttreatment to follow-up	0.11	0.26	.66
<u>PABS</u>			
Group	.02	0.93	.99
Baseline to follow-up	-3.54	0.84	<.001
Posttreatment to follow-up	-0.30	0.47	.53

Note. For treatment condition, standard treatment serves as the reference group. For time, follow-up serves as the reference group. COIS-RP = Child Obsessive-Compulsive Disorder (OCD) Impairment Scale – Parent-Report Revised; CYBOCS = Children's Yale-Brown Obsessive Compulsive Scale; FAS = Family Accommodation Scale; FES = Family Environment Scale; PABS = Parental Attitudes and Behaviors Scale.

^a Regardless of condition, acute treatment responders maintained improvements at follow-up, with improved outcomes at follow-up compared to baseline.