

Additional file 8. Multivariate analysis of variance (MANOVA) with post hoc pairwise comparisons of cranium and mandible shape between major groups of squamates. *F*-values and *p*-values are shown in upper and lower off-diagonal cells, respectively. Significant values are highlighted in bold.

CRANIUM	Anguimorpha	Gekkota	Iguania	Lacertoidea	Scincoidea	Serpentes
Anguimorpha	-	2.71	3.5	1.41	5.44	5.58
Gekkota	0.1307	-	1.17	2.49	3.23	11.95
Iguania	0.0696	0.2882	-	16.98	18.16	21.03
Lacertoidea	0.2496	0.1344	0.0002	-	0.69	5.25
Scincoidea	0.0351	0.0996	0.0002	0.417	-	7.07
Serpentes	0.0216	0.0011	<0.0001	0.0253	0.0101	-

MANDIBLE	Anguimorpha	Gekkota	Iguania	Lacertoidea	Scincoidea	Serpentes
Anguimorpha	-	0.21	16.25	1.20	2.34	7.70
Gekkota	0.6533	-	8.73	1.01	1.78	4.32
Iguania	0.0004	0.0069	-	2.55	6.65	7.14
Lacertoidea	0.2868	0.3309	0.1197	-	0.12	7.93
Scincoidea	0.1486	0.2092	0.0155	0.7296	-	5.33
Serpentes	0.0074	0.0425	0.0089	0.0065	0.0246	-