

The TIDieR (Template for Intervention Description and Replication) Checklist*:

Information to include when describing an intervention and the location of the information

Item	Item	Where located **	
number		Primary paper	Other † (details)
		(page or appendix	
		number)	
	BRIEF NAME		
1.	Provide the name or a phrase that describes the intervention.	Title, Abstract (1),	
		Introduction (3)	
	WHY		
2.	Describe any rationale, theory, or goal of the elements essential to the intervention.	Introduction (2-3),	
		Methods: The	
		Bandebereho	
		intervention (9-12)	
	WHAT		
3.	Materials: Describe any physical or informational materials used in the intervention, including	Methods: The	
	those provided to participants or used in intervention delivery or in training of intervention	Bandebereho	
	providers. Provide information on where the materials can be accessed (e.g. online appendix,	intervention (9-10)	
	URL).		
	Dreadures Describe each of the presedures estivities and/or presedured in the	Mathadar Tha	
4.	Procedures: Describe each of the procedures, activities, and/or processes used in the	Methods: The	
	intervention, including any enabling or support activities.	Bandebereho	
	WHO PROVIDED	intervention (9-12)	
-	WHO PROVIDED	Mathada, The	
5. 	For each category of intervention provider (e.g. psychologist, nursing assistant), describe their	Methods: The	

	expertise, background and any specific training given.	Bandebereho	
		intervention (12)	
	HOW		
6.	Describe the modes of delivery (e.g. face-to-face or by some other mechanism, such as internet	Methods: The	
	or telephone) of the intervention and whether it was provided individually or in a group.	Bandebereho	
		intervention (12)	
	WHERE		
7.	Describe the type(s) of location(s) where the intervention occurred, including any necessary	Methods: The	
	infrastructure or relevant features.	Bandebereho	
		intervention (12)	
	WHEN and HOW MUCH		
8.	Describe the number of times the intervention was delivered and over what period of time	Methods: Procedure	
	including the number of sessions, their schedule, and their duration, intensity or dose.	(7), Methods: The	
		Bandebereho	
		intervention (9-12)	
	TAILORING		
9.	If the intervention was planned to be personalised, titrated or adapted, then describe what, why,	N/A	
	when, and how.		
	MODIFICATIONS		
10. [‡]	If the intervention was modified during the course of the study, describe the changes (what, why,	N/A	
	when, and how).		
	HOW WELL		
11.	Planned: If intervention adherence or fidelity was assessed, describe how and by whom, and if	Methods: The	
	any strategies were used to maintain or improve fidelity, describe them.	Bandebereho	
		intervention (12)	
12. [‡]	Actual: If intervention adherence or fidelity was assessed, describe the extent to which the	Methods: The	

intervention was delivered as planned.	Bandebereho
	intervention (9-12)

- ** **Authors** use N/A if an item is not applicable for the intervention being described. **Reviewers** use '?' if information about the element is not reported/not sufficiently reported.
- † If the information is not provided in the primary paper, give details of where this information is available. This may include locations such as a published protocol or other published papers (provide citation details) or a website (provide the URL).
- ‡ If completing the TIDieR checklist for a protocol, these items are not relevant to the protocol and cannot be described until the study is complete.
- * We strongly recommend using this checklist in conjunction with the TIDieR guide (see BMJ 2014;348:g1687) which contains an explanation and elaboration for each item.
- * The focus of TIDieR is on reporting details of the intervention elements (and where relevant, comparison elements) of a study. Other elements and methodological features of studies are covered by other reporting statements and checklists and have not been duplicated as part of the TIDieR checklist. When a randomised trial is being reported, the TIDieR checklist should be used in conjunction with the CONSORT statement (see www.consort-statement.org) as an extension of ttem 5 of the CONSORT 2010 Statement. When a clinical trial protocol is being reported, the TIDieR checklist should be used in conjunction with the SPIRIT statement as an extension of ttem 11 of the SPIRIT 2013
 Statement (see www.spirit-statement.org). For alternate study designs, TIDieR can be used in conjunction with the appropriate checklist for that study design (see www.equator-network.org).