

Title;

The oscillation of intracellular Ca²⁺ influx associated with the circadian expression of *Piezo1* and *TRPV4* in the bladder urothelium.

Tatsuya Ihara¹, Takahiko Mitsui¹, Yuki Nakamura², Mie Kanda¹, Sachiko Tsuchiya¹,
Satoru Kira¹, Hiroshi Nakagomi¹, Norifumi Sawada¹, Manabu Kamiyama¹, Yuri Hirayama³,
Eiji Shigetomi³, Youichi Shinozaki³, Mitsuharu Yoshiyama¹, Atsuhito Nakao², Masayuki
Takeda^{1*} and Schuichi Koizumi^{3*}

¹Department of Urology, Interdisciplinary Graduate School of Medicine, University of
Yamanashi, Chuo, Yamanashi, Japan

²Department of Immunology, Interdisciplinary Graduate School of Medicine, University of
Yamanashi, Chuo, Yamanashi, Japan

³Department of Neuropharmacology, Interdisciplinary Graduate School of Medicine,
University of Yamanashi, Chuo, Yamanashi, Japan

*Correspondence author:

Schuichi Koizumi

1110 Shimokato, Chuo, Yamanashi, 409-3898 Japan

Tel: +81-55-273-9503

Fax: +81-55-273-6739

E-mail: skoizumi@yamanashi.ac.jp

Masayuki Takeda


1110 Shimokato, Chuo, Yamanashi, 409-3898 Japan

Tel: +81-55-273-9643

Fax: +81-55-273-9659

E-mail: matakeda@yamanashi.ac.jp

Supplementary Figure 1



Supplementary Figure 1. *In vitro* cell stretch system.

A, Elastic silicone chambers. B and C, The silicone chamber was set in the 2 arms of the extension device on the microscope stage. The arrow in C indicates the direction of extension.