

S8 Table. Lobe volume determination on manual and automatic segmentation maps (Expiration B60f scans).

		Baseline	3 months	12 months	24 months	Overall
RUL	manually corrected (mean±sd)	230±86.22	245.4±94.3	271.73±104.96	310.56±114.78	264.31±102.89
	automatic (mean±sd)	228.89±112	249.51±138.37	313.5±162.71	346.75±159.26	284.2±148.66
	Δ (mean±sd)	1.11±60.57	-4.1±83.06	-41.77±85.52	-36.19±88.6	-19.9±80.44
	LoA (2.5% limit, 97.5% limit)	(-117.61, 119.83)	(-166.91, 158.7)	(-209.38, 125.85)	(-209.85, 137.47)	(-177.55, 137.76)
	Regr. (Intercept / Slope)	-79.73 / 1.41	-29.28 / 1.2	-68.56 / 1.39	-19.03 / 1.11	-46.57 / 1.25
	Regr (Pearson's r)	0.84	0.81	0.88	0.84	0.86
RML	manually corrected (mean±sd)	134.68±39.51	134.6±59.44	144.07±49.14	153.96±43.4	141.87±48.05
	automatic (mean±sd)	154.91±67.34	123.64±90.66	131.46±44.5	168.03±56.86	144.77±68.42
	Δ (mean±sd)	-20.24±59.69	10.96±68.71	12.61±34.88	-14.07±34.35	-2.9±52.78
	LoA (2.5% limit, 97.5% limit)	(-137.22, 96.75)	(-123.71, 145.63)	(-55.76, 80.98)	(-81.39, 53.26)	(-106.35, 100.55)
	Regr. (Intercept / Slope)	-210.72 / 2.68	-160.25 / 2.18	1.3 / 0.99	-58.98 / 1.41	-88.39 / 1.63
	Regr (Pearson's r)	0.48	0.65	0.73	0.8	0.64
RLL	manually corrected (mean±sd)	338.15±92.55	349.91±146.68	368.35±119.43	426.88±148.03	370.86±130.39
	automatic (mean±sd)	309.35±124.27	349.18±174.76	328.55±117.29	366.43±123.65	338.53±135.66
	Δ (mean±sd)	28.8±86.71	0.73±87.77	39.8±100.63	60.46±109.79	32.33±96.77
	LoA (2.5% limit, 97.5% limit)	(-141.15, 198.74)	(-171.3, 172.75)	(-157.44, 237.03)	(-154.73, 275.64)	(-157.34, 221.99)
	Regr. (Intercept / Slope)	-116.94 / 1.24	-93.61 / 1.19	-55.61 / 1.08	-42.27 / 1	-60.05 / 1.08
	Regr (Pearson's r)	0.72	0.87	0.64	0.69	0.74
LUL	manually corrected (mean±sd)	172.68±68.44	171.37±68.27	205.89±86.89	230.07±89.51	194.83±80.76
	automatic (mean±sd)	232.02±157.69	185.97±95.79	257.42±120.77	229.93±79.9	225.84±117.22
	Δ (mean±sd)	-59.34±136.7	-14.6±90	-51.53±82.52	0.14±74.64	-31.01±99.97
	LoA (2.5% limit, 97.5% limit)	(-327.27, 208.58)	(-191, 161.8)	(-213.27, 110.21)	(-146.15, 146.44)	(-226.96, 164.94)
	Regr. (Intercept / Slope)	-330.22 / 3.22	-46.38 / 1.39	-19.03 / 1.17	41.71 / 0.84	-40.92 / 1.33
	Regr (Pearson's r)	0.5	0.44	0.73	0.62	0.54
LLi	manually corrected (mean±sd)	95.99±33.01	108.06±49.29	117.96±35.3	123.31±36.67	111±39.73
	automatic (mean±sd)	80.64±72.05	106.46±104.34	106.85±63.94	148.98±121.23	110.93±95.66
	Δ (mean±sd)	15.35±75.75	1.6±94.11	11.11±50.38	-25.67±102.31	0.08±83.9
	LoA (2.5% limit, 97.5% limit)	(-133.12, 163.81)	(-182.85, 186.06)	(-87.64, 109.85)	(-226.2, 174.85)	(-164.36, 164.52)
	Regr. (Intercept / Slope)	-383.66 / 4.24	-163.23 / 2.6	-107.6 / 1.83	-99.37 / 1.87	-149.42 / 2.26
	Regr (Pearson's r)	0.11	0.43	0.62	0.63	0.49
LLL	manually corrected (mean±sd)	296.68±93.5	301.04±126.38	328.45±124.97	370.27±128.35	324.04±119.91
	automatic (mean±sd)	232.45±122.29	281.1±136.01	272.03±137.58	338.1±134.75	281.06±135.02
	Δ (mean±sd)	64.23±96.55	19.95±39.41	56.42±48.63	32.17±49.97	42.98±64.01
	LoA (2.5% limit, 97.5% limit)	(-125.01, 253.48)	(-57.3, 97.19)	(-38.89, 151.74)	(-65.78, 130.11)	(-82.47, 168.43)
	Regr. (Intercept / Slope)	-111.34 / 1.25	-63.87 / 1.11	-94.44 / 1.15	-57.57 / 1.09	-82.19 / 1.14
	Regr (Pearson's r)	0.63	0.96	0.94	0.93	0.88
LUL+LLi	manually corrected (mean±sd)	268.67±94.49	279.43±114.16	318.03±119.55	353.38±122.37	304.67±115.4
	automatic (mean±sd)	312.67±132.63	293.07±120.05	350.03±127.98	378.91±142.08	333.41±132.13
	Δ (mean±sd)	-44±95.35	-13.64±33.05	-31.99±39.85	-25.53±40.93	-28.74±57.9
	LoA (2.5% limit, 97.5% limit)	(-230.89, 142.9)	(-78.42, 51.14)	(-110.09, 46.1)	(-105.75, 54.69)	(-142.21, 84.74)
	Regr. (Intercept / Slope)	-89.92 / 1.5	-12.2 / 1.09	32.2 / 0.96	-27.99 / 1.12	-9.1 / 1.1
	Regr (Pearson's r)	0.7	0.96	0.95	0.96	0.9

Lobar volumes [cm³]calculated for fully automatic and manually corrected segmentation on expiration B60f scans at baseline, 3, 12 and 24 months.

The last column summarizes all time points. All values are separately calculated for the right upper (RUL), middle (RML) and lower lobe (RLL), the

left upper lobe (LUL), the lingula (LLi), the left lower lobe (LLL), and also combining left upper lobe and lingula into one lobe (LUL+LLi). Both methods are compared in accordance with the approach of Bland-Altman giving mean differences (Δ), limits of agreement (LoA) and two regression coefficients (Intercept / Slope and Pearson's correlation coefficient).