

Online Supplemental Appendix

Supplemental Table 1: Logistic regression analysis of Model 3.

	OR (95% CI)	p-Value	AUC
Age <60	0.54 (0.44, 0.67)	<0.001	0.63
Medical History	0.67 (0.50, 0.91)	0.009	
Tobacco Use	0.91 (0.81, 1.01)	0.07	
Weight (per 10 lbs)	1.00 (1.00, 1.00)	0.002	
Prior VTE	1.21 (0.97, 1.50)	0.09	
Alcohol use	0.49 (0.31, 0.77)	0.002	

VTE, venous thromboembolism; OR, Odds Ratio; CI, Confidence Interval; AUC, Area Under the Receiver Operator Curve.

Supplemental Table 2: Relationship between SAMe-TT₂R₂ scores and adverse clinical outcomes.

SAMe-TT₂R₂	N	Recurrent VTE (%)	ISTH Major Bleeding (%)	Overall Adverse Events (%)	Total follow-up years
0	155	1 (0.6)	7 (4.5)	8 (5.2)	197.4
1	510	6 (1.2)	16 (3.1)	22 (4.3)	592.4
2	432	12 (2.8)	15 (3.5)	27 (6.3)	463.3
3	431	17 (3.9)	22 (5.1)	39 (9.0)	462.4
4	289	16 (5.5)	6 (2.1)	22 (7.6)	275.0
>5	126	4 (3.2)	4 (3.2)	8 (6.3)	136.8

VTE, venous thromboembolism; ISTH, International Society of Thrombosis and Haemostasis