


Supplementary Table 1: Descriptive characteristics of the community controls in the study according to angina frequency score categories

Variables	No angina in past 4 weeks (N= 99)	Angina in past 4 weeks (N= 11)	Total population (N= 110)
Demographic factors			
Age, Mean (SD)	50 (9)	48 (11)	50 (9)
Women, N (%)	50 (50%)	7 (63%)	57 (52%)
African-American, N (%)	38 (38%)	8 (72%)	46 (42%)
Below poverty line, N (%)	7 (7%)	2 (18%)	9 (9%)
Current smokers, N (%)	3 (3%)	2 (18%)	5 (5%)
Psychological factors			
BDI-II score, Mean (SD) ^a	5 (5)	17 (14)	6 (7)
STAI Anxiety-Trait score, Mean (SD) ^a	31 (8)	40 (15)	32 (10)
STAXI Anger-Trait score, Mean (SD)	14 (3)	15 (4)	14 (3)
PCL score, Mean (SD) ^a	23 (8)	39 (20)	24 (11)
Perceived-stress score, Mean (SD) ^a	10 (6)	17 (9)	10 (6)

SD: standard deviation; BDI: Beck Depression Inventory; STAI: State- Trait Anxiety Inventory; STAXI: State-Trait Anger Expression Inventory; PCL: PTSD Symptom Checklist (Civilian)

^a P value < 0.05



Angina and MSIMI_AmJMed_figure1_103117.jpg