

SUPPLEMENTARY TABLE S3. PAIRWISE COMPARISON OF DOSE LEVELS FOR SURVIVAL  
(ODDS OF DOSE 1 OVER ODDS OF DOSE 2)

<i>Dose 1 (mM)</i>	<i>Dose 2 (mM)</i>	<i>t</i>	<i>Pr &gt;  t </i>	<i>Odds ratio</i>	<i>Lower confidence limit for odds ratio</i>	<i>Upper confidence limit for odds ratio</i>
0	1.3	6.35	<0.0001	3.520	2.350	5.274
0	1.4	2.90	0.0068	1.779	1.187	2.665
0	1.5	8.43	<0.0001	3.634	2.659	4.967
0	1.6	10.55	<0.0001	9.267	6.025	14.253
0	1.7	10.77	<0.0001	61.297	28.123	133.604
1.3	1.4	-2.76	0.0097	0.505	0.305	0.837
1.3	1.5	0.15	0.8823	1.032	0.668	1.596
1.3	1.6	3.75	0.0007	2.632	1.555	4.455
1.3	1.7	6.97	<0.0001	17.413	7.551	40.156
1.4	1.5	3.36	0.0021	2.043	1.324	3.155
1.4	1.6	6.43	<0.0001	5.210	3.087	8.794
1.4	1.7	8.66	<0.0001	34.464	14.974	79.322
1.5	1.6	4.16	0.0002	2.550	1.612	4.032
1.5	1.7	7.25	<0.0001	16.866	7.620	37.333
1.6	1.7	4.55	<0.0001	6.615	2.838	15.418

Survival percent was negatively correlated with nicotine dose (Spearman's correlation = -0.62,  $p < 0.0001$ ). Data is presented as a bar graph in Figure 1A.