

SUPPLEMENTARY TABLE S4. PAIRWISE COMPARISON OF DOSE LEVELS FOR HEART DEFECTS
(ODDS OF DOSE 1 OVER ODDS OF DOSE 2)

<i>Doses (mM)</i>	<i>t</i>	<i>Pr > t </i>	<i>Odds ratio</i>	<i>Odds ratio (lower)</i>	<i>Odds ratio (upper)</i>
1.4 vs. 1.3	16.14	<0.0001	3.4769	2.9885	4.0450
1.5 vs. 1.3	-0.87	0.3858	0.9409	0.8200	1.0797
1.6 vs. 1.3	29.58	<0.0001	25.1883	20.3403	31.1919
1.7 vs. 1.3	16.42	<0.0001	273.76	140.12	534.87
1.5 vs. 1.4	-21.51	<0.0001	0.2706	0.2402	0.3049
1.6 vs. 1.4	19.12	<0.0001	7.2445	5.9132	8.8756
1.7 vs. 1.4	12.87	<0.0001	78.7366	40.4880	153.12
1.6 vs. 1.5	33.41	<0.0001	26.7697	22.0740	32.4644
1.7 vs. 1.5	16.76	<0.0001	290.94	149.88	564.76
1.7 vs. 1.6	6.84	<0.0001	10.8684	5.4877	21.5247

There is a strong positive correlation between nicotine dose and level of heart defect (Spearman's correlation = 0.80, $p < 0.0001$). Data are presented as a bar graph in Figure 1G.