

# BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## Impact of dexmedetomidine infusion during general anesthesia on incidence of postoperative delirium in elderly patients after major non-cardiac surgery: study protocol of a randomized, double-blinded and placebo-controlled trial

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-019549
Article Type:	Protocol
Date Submitted by the Author:	12-Sep-2017
Complete List of Authors:	Wang, Bojie; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Li, Chunjing; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Hu, Jian; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Li, Huaijin; Peking University First Hospital, Department of Anesthesiology and Critical Care Guo, Chao; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Wang, Zhenhan; Dongping People's Hospital, Department of anesthesiology Zhang, Qiaochu; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Mu, Dongliang; Department of Anesthesiology and Critical Care Medicine Wang, Dong-Xin; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine
<b>Primary Subject Heading</b>:	Anaesthesia
Secondary Subject Heading:	Anaesthesia
Keywords:	Adult anaesthesia < ANAESTHETICS, Delirium & cognitive disorders < PSYCHIATRY, Pain management < RADIOTHERAPY

SCHOLARONE™  
Manuscripts

## Note from the Editors: Instructions for reviewers of study protocols

---

Since launching in 2011, BMJ Open has published study protocols for planned or ongoing research studies. If data collection is complete, we will not consider the manuscript.

Publishing study protocols enables researchers and funding bodies to stay up to date in their fields by providing exposure to research activity that may not otherwise be widely publicised. This can help prevent unnecessary duplication of work and will hopefully enable collaboration. Publishing protocols in full also makes available more information than is currently required by trial registries and increases transparency, making it easier for others (editors, reviewers and readers) to see and understand any deviations from the protocol that occur during the conduct of the study.

The scientific integrity and the credibility of the study data depend substantially on the study design and methodology, which is why the study protocol requires a thorough peer-review.

*BMJ Open* will consider for publication protocols for any study design, including observational studies and systematic reviews.

Some things to keep in mind when reviewing the study protocol:

- Protocol papers should report planned or ongoing studies. The dates of the study should be included in the manuscript.
- Unfortunately we are unable to customize the reviewer report form for study protocols. As such, some of the items (i.e., those pertaining to results) on the form should be scores as Not Applicable (N/A).
- While some baseline data can be presented, there should be no results or conclusions present in the study protocol.
- For studies that are ongoing, it is generally the case that very few changes can be made to the methodology. As such, requests for revisions are generally clarifications for the rationale or details relating to the methods. If there is a major flaw in the study that would prevent a sound interpretation of the data, we would expect the study protocol to be rejected.

1  
2  
3 **Impact of dexmedetomidine infusion during general anesthesia on incidence of**  
4 **postoperative delirium in elderly patients after major non-cardiac surgery: study**  
5 **protocol of a randomized, double-blinded and placebo-controlled trial**  
6  
7  
8  
9

10  
11 Bo-Jie Wang<sup>1</sup>, Chun-Jing Li<sup>1</sup>, Jian Hu<sup>1</sup>, Huai-Jin Li<sup>1</sup>, Chao Guo<sup>1</sup>, Zhen-Han Wang<sup>2</sup>,  
12 Qiao-Chu Zhang<sup>1</sup>, Dong-Liang Mu<sup>1,\*</sup>, Dong-Xin Wang<sup>1</sup>  
13  
14  
15

16  
17  
18 <sup>1</sup>Department of Anesthesiology and Critical Care Medicine, Peking University First  
19 Hospital, Beijing, China  
20  
21

22 <sup>2</sup>Department of Anesthesiology, Dongping People's Hospital, Shandong, China  
23  
24  
25

26 The first two authors (BJ Wang and CJ Li) contributed equally to this manuscript  
27  
28  
29

30  
31 Corresponding author: Dong-Liang Mu, M.D., Department of Anesthesiology and  
32 Critical Care Medicine, Peking University First Hospital, Beijing, China. Tel: +86  
33 010 83572784 (O), +86 13810702725 (M); E-mail: mudongliang@icloud.com  
34  
35  
36  
37

38  
39 Bo-Jie Wang: ivyxbd@hotmail.com  
40

41 Chun-Jing Li: gogolion@163.com  
42

43 Jian Hu: hj19871204@163.com  
44

45 Huai-Jin Li: sophie.lee.coffee@gmail.com  
46  
47

48 Chao Guo: guochao1988@yeah.com  
49

50 Zhen-Han Wang: 1468432914@qq.com  
51

52 Qiao-Chu Zhang: 372193745@qq.com  
53

54 Dong-Liang Mu: mudongliang@icloud.com  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Dong-Xin Wang: wangdongxin@hotmail.com

Word Count: 2547

For peer review only

## Abstract for protocol

**Introduction:** Delirium is a common complication in the elderly after surgery and is associated with worse outcomes. Multiple risk factors are related with postoperative delirium, such as exposure to general anesthetics, pain and inflammatory response.

Preclinical and clinical studies reported that dexmedetomidine could attenuate neurotoxicity induced by general anesthetics, improve pain management and inhibit inflammatory response. Several studies observed the relationship between intraoperative use of dexmedetomidine and postoperative delirium, but the results were inconsistent. This study is designed to investigate the impact of dexmedetomidine administered during general anesthesia in preventing delirium in elderly patients after major non-cardiac surgery.

**Method and analysis:** This is a randomized, double-blinded, and placebo-controlled trial. 620 elderly patients (age  $\geq 60$  years) who are scheduled to undertake elective major non-cardiac surgery (with an expected duration  $\geq 2$  hours) are randomly divided into two groups. For patients in dexmedetomidine group, a loading dose dexmedetomidine (0.6  $\mu\text{g}/\text{kg}$ ) will be administered in 10 minutes before anesthesia induction, followed by a continuous infusion at a rate of 0.5  $\mu\text{g}/\text{kg}/\text{h}$  till 1 hour before the end of surgery. For patients in control group, normal saline will be administered in the same rate and volume as in the dexmedetomidine group. The primary endpoint is the incidence of delirium during the first five postoperative days. Secondary endpoints include pain intensity, cumulative opioid consumption and subjective sleep quality during the first 3 postoperative days, and the incidence of non-delirium complications and all-cause mortality within 30 days after surgery.

1  
2  
3 **Ethics and dissemination:** The study protocol was approved by Clinical Research  
4  
5 Ethics Committee of Peking University First Hospital (2015-987) and registered at  
6  
7 Chinese Clinical Trial Registry (<http://www.chictr.org.cn>) with identifier ChiCTR-  
8  
9 IPR-15007654. Results of the study will be presented at academic conferences and  
10  
11 submitted to peer-reviewed journals.  
12  
13

14  
15 **Key words:** intraoperative dexmedetomidine; postoperative delirium; old patient;  
16  
17 major non-cardiac surgery  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Strengths and limitations

1. This study is designed as a randomized, double-blinded and placebo-controlled trial with a large sample size.
2. Results of our study will provide evidence on the impact of dexmedetomidine infusion during general anesthesia on the incidence of delirium in elderly patients after non-cardiac surgery.
3. Bispectral index is monitored to guide anesthesia maintenance which will overcome an important limitation in a recent trial.
4. Safety outcome data will be recorded in detail.
5. As a single site trial, the generalizability of our results will be limited.



## Introduction

Delirium is considered as an organic brain syndrome which is characterized by altered consciousness, inattention, and changes in cognition or perception; it develops acutely with clinical manifestations fluctuate during the course of the day.<sup>1</sup> Prevalence of delirium varies from 14.8% to 23.0% in patients after major non-cardiac surgery.<sup>2-3</sup>

The occurrence of postoperative delirium (POD) is associated with worsen outcomes including prolonged mechanical ventilation and intensive care unit stay, increased postoperative complications, high mortality rate and long-term cognitive decline.<sup>2-5</sup>

The etiology of POD is multifactorial and includes several intraoperative factors.<sup>6</sup> For example, exposure to general anesthetics (e.g., propofol or sevoflurane) might produce neurotoxicity,<sup>7-8</sup> whereas reduced anesthetic exposure by avoiding deep anesthesia reduced the occurrence of delirium in elderly patients undergoing major non-cardiac surgery.<sup>9</sup> Poor pain management is another risk factor of POD.<sup>6, 10</sup> It was reported that the risk of postoperative delirium was 1.2 times higher for every unit increment in visual analogue pain score (an 11-point pain scale where 0 indicates no pain and 10 the most severe pain).<sup>11</sup> Inflammation is also proposed to play an important role in the pathogenesis of POD.<sup>2</sup> Inflammatory responses induced by surgery and anesthesia are manifested by elevated levels of interleukins (IL), C-reactive protein (CRP) and tumor necrosis factor (TNF).<sup>2, 12</sup> Studies by our group and other found that higher levels of inflammatory mediators are associated with increased risk of POD.<sup>2, 13-16.</sup>

Dexmedetomidine is a highly selective  $\alpha_2$ -receptor agonist with sedative, analgesic and anxiolytic effects.<sup>17-21</sup> When used as a supplement during intraoperative anesthesia, it reduces the consumption of general anesthetics.<sup>17</sup> Preclinical evidences

1  
2  
3 suggest that use of dexmedetomidine might attenuate neurotoxicity induced by  
4 general anesthetics.<sup>18-19</sup> In a meta-analysis, intraoperative administration of  
5 dexmedetomidine lowers postoperative pain intensity and reduces opioid  
6 consumption.<sup>20</sup> Clinical evidence also showed that intraoperative dexmedetomidine  
7 significantly inhibits hyper-secretion of inflammatory cytokines during and after  
8 surgery.<sup>21-22</sup>

15  
16 Use of dexmedetomidine during general anesthesia may reduce POD. In  
17 pediatric patients undergoing tonsillectomy and cardiac surgery, intraoperative  
18 infusion of dexmedetomidine lowered the incidence of emergence delirium.<sup>23,24</sup> In  
19 adult patients undergoing cardiac surgery and microvascular free flap surgery,  
20 intraoperative dexmedetomidine (comparison with normal saline) slightly decreased  
21 the incidence of delirium, although the differences were not statistically significant  
22 between two groups possibly due to underpowered sample size.<sup>25,26</sup> In a recent study  
23 of Deiner et al.,<sup>27</sup> use of dexmedetomidine during general anesthesia did not reduce  
24 delirium after major non-cardiac surgery in the elderly. However, in that study,  
25 anesthesia depth was not monitored and the consumption of anesthetics (such as  
26 propofol and fentanyl) was similar between the two groups. It was possible that  
27 patients in the dexmedetomidine group received deeper anesthesia which might have  
28 increased the risk of delirium.<sup>27</sup> Therefore, the effects of dexmedetomidine  
29 administered during general anesthesia on the occurrence of POD need to be  
30 evaluated further.

31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48 This study is designed to investigate whether dexmedetomidine use during  
49 general anesthesia can decrease the incidence of POD in elderly patients after major  
50 non-cardiac surgery.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Method and analysis

### Study design

This randomized, double-blinded, and placebo-controlled trial with two parallel arms is designed to test the superiority of dexmedetomidine administered during general anesthesia on the incidence of delirium after surgery. Patients will be randomized into either the dexmedetomidine group or the control group (Figure 1). The study is conducted in the Department of Anesthesiology and Critical Care Medicine of Peking University First Hospital.

### Ethics approval

The study protocol (version 1.1, issue date Nov, 2015) was approved by Clinical Research Ethics Committee of Peking University First Hospital (2015-987). Any protocol modifications will be submitted for review and approval by the Ethics Committee. The trial was registered at Chinese Clinical Trial Registry (<http://www.chictr.org.cn>) on December 1, 2015, with identifier ChiCTR-IPR-15007654. Written informed consents are obtained from patients or their surrogate in law.

### Participants

Elderly (age  $\geq 60$  years) patients who are scheduled to undergo elective non-cardiac surgery with expected duration  $\geq 2$  hours under general anesthesia are screened for inclusion. Those who meet any of the following criteria will be excluded: (1) do not provide written informed consents; (2) previous history of schizophrenia, epilepsy or Parkinson disease; (3) visual, hearing, language or other barrier which impede communication and preoperative delirium assessment; (4) neurosurgery or traumatic brain injury; (5) severe bradycardia (heart rate less than 40 beats per minute), sick

1  
2  
3 sinus syndrome or atrioventricular block of degree 2 or above; (6) severe hepatic  
4 dysfunction (Child-Pugh grade C); or (7) renal failure (requirement of renal  
5 replacement therapy).  
6  
7

### 8 9 **Patient recruitment and baseline data collection**

10  
11 Potential participants are screened by qualified investigators the day before surgery  
12 (or on Friday for those who will undergo surgery next Monday). The study protocol  
13 including potential risks and benefits is explained in detail. Those who meet the  
14 inclusion and exclusion are invited to participate in the study.  
15  
16  
17  
18

19  
20 After obtaining written informed consents, the following baseline data are  
21 collected: demographic data, preoperative diagnosis, comorbidity, current medical  
22 therapy, previous surgery, and main results of physical and laboratory examinations.  
23  
24 Barthel index is used for evaluation of activities of daily living.<sup>28</sup> Cognitive function  
25 is assessed with Mini-Mental State Examination (MMSE).<sup>29</sup> Preoperative delirium is  
26 assessed with confusion assessment method (CAM).<sup>30</sup>  
27  
28  
29  
30  
31

### 32 33 **Randomization, grouping and blinding**

34  
35 Random numbers were created by an independent statistician using SAS statistical  
36 package version 9.3 (SAS Institute, Cary, NC, USA) in a 1:1 ratio and were sealed in  
37 sequentially numbered envelopes. A study coordinator, who does not participate in  
38 anesthesia and postoperative follow-up of enrolled patients, will open envelop for  
39 random numbers and prepare study drugs before induction of anesthesia. In this way,  
40 patients are randomly divided into either dexmedetomidine or control group.  
41  
42  
43  
44  
45  
46  
47

48 Information regarding randomization, study drug preparation and group  
49 allocation will be masked from investigators who perform data collection and  
50 postoperative follow-up, anesthesiologists, patients and other healthcare team  
51 members. Blinding will be maintained throughout the entire study period.  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 To ensure patients' safety, study group allocation can be unmasked in the  
4 following conditions, i.e., occurrence of severe adverse events or any unexpected  
5 deterioration in the patient's clinical status. These situations will be documented in  
6 the Case Report Forms (CRFs). The unmasked patients will be included in the  
7 intention-to-treat population but excluded from per-protocol analysis.  
8  
9

### 10 11 12 13 **Interventions, anesthesia and analgesia**

14  
15 Intraoperative monitoring includes electrocardiogram, noninvasive blood pressure,  
16 pulse oxygen saturation, end-tidal carbon dioxide, nasopharyngeal temperature, urine  
17 output, and bispectral index (BIS). Intraarterial pressure (including derivative  
18 dynamic parameters such as stroke volume variation by FlowTrac system) and central  
19 venous pressure are monitored according to patients' conditions.  
20  
21

22  
23 Study drugs, either 200 µg (2 ml) dexmedetomidine (Jiangsu Hengrui Medicine  
24 Co, Ltd, Jiangsu, China) or 2 ml normal saline, are diluted in 50 ml normal saline. All  
25 study drugs are colorless solution provided in syringes of the same size and brand.  
26  
27 The regimen of study drug administration includes a loading dose of 0.15 ml/kg (i.e.,  
28 0.6 µg/kg dexmedetomidine for patients in the dexmedetomidine group) administered  
29 during a 10-minute period before anesthesia induction. This is followed by a  
30 continuous infusion at a rate of 0.125 ml/kg/h (i.e., a rate of 0.5 µg/kg/h  
31 dexmedetomidine for patients in the dexmedetomidine group) till 1 hour before the  
32 end of surgery. Study drug infusion is performed by an infusion pump specially  
33 designed for dexmedetomidine administration (Slgo<sup>®</sup> CP1000, Beijing Slgo medical  
34 technology Co., Ltd.).  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50  
51 Attending anesthesiologists can decrease or stop study drug infusion in the  
52 following conditions: (1) severe bradycardia or hypotension which does not improve  
53 after routine treatment; (2) new onset atrioventricular block which does not improve  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 after routine treatment; or (3) other conditions that anesthesiologists consider it  
4 necessary. In these conditions the reasons that lead to any protocol deviations will be  
5 recorded in the Case Report Forms. These patients will be included in the intention-  
6 to-treat analysis but excluded from the per-protocol analysis.  
7  
8  
9  
10

11 Total intravenous anesthesia is performed for all patients. Anesthesia is induced  
12 with intravenous sufentanil (target controlled infusion with effect-site concentration  
13 from 0.2 to 0.5 ng/ml) and propofol (2-3 mg/kg) and maintained with sufentanil  
14 (effect-site concentration from 0.2 to 0.5 ng/ml) and propofol (4-12 mg/kg/h).  
15 Rocuronium and/or cisatracurium are administered for muscle relaxation. The mean  
16 arterial pressure is maintained above 60 mmHg or within 20% from baseline. BIS is  
17 maintained between 40 and 60. Mechanical ventilated is performed with a 1:1 nitrous  
18 oxide-oxygen mixture.  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 All patients are transferred to post-anesthesia care unit (PACU) or intensive care  
29 unit (ICU) before they are sent back to general wards. Patient-controlled intravenous  
30 analgesia (PCIA) is provided for all patients, which is established with 0.5 mg/ml  
31 morphine in 100 ml normal saline and programmed to deliver a 1 mg bolus with a  
32 lock-out interval of 8 minutes and a background infusion at 0.5 mg/h. Supplemental  
33 morphine at dose of 2 to 4 mg will be administered at 10-minute intervals if the  
34 numeric rating scale (NRS) pain score (a 11-point scale where 0 indicates no pain and  
35 10 indicates the worst pain) remains above 4 after 3 consecutive PCIA boluses.<sup>31</sup>  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

#### 46 **Outcome assessment**

47 Patients are followed up twice daily during the first 5 postoperative days and then  
48 weekly until 30 days after surgery.  
49  
50

#### 51 **Primary endpoint**

1  
2  
3 The primary endpoint is the incidence of delirium during the first 5 days after surgery.  
4  
5 Delirium is assessed twice daily (at 08:00-09:00 and 19:00-20:00, respectively) with  
6  
7 CAM for non-intubated patients<sup>30</sup> or CAM for the Intensive Care Unit (CAM-ICU)  
8  
9 for intubated patients.<sup>32</sup> These delirium assessment methods had been used in our  
10  
11 previous studies.<sup>2-3,31</sup> For patients who are discharged or died within 5 days after  
12  
13 surgery, the results of last delirium assessment will be considered the results of the  
14  
15 missing data. These patients will be excluded when calculating daily prevalence of  
16  
17 delirium in a post-hoc analysis.  
18  
19

20  
21 Investigators who are responsible for delirium assessment and postoperative  
22  
23 follow-up are not involved in anesthesia and perioperative care. Before the beginning  
24  
25 of the study, they are trained by a psychiatrist to perform delirium assessment and the  
26  
27 training process is repeated at 4-month intervals during the study period.<sup>2-3,31</sup>  
28

### 29 ***Secondary endpoints***

30  
31 Postoperative pain intensities at rest and with movement are assessed with NRS pain  
32  
33 score at 24, 48 and 72 hours after surgery, respectively.<sup>31</sup> Cumulative morphine  
34  
35 consumptions at these time points are recorded. Subjective sleep quality is assessed  
36  
37 with NRS (an 11-point scale where 0 indicates the worst possible sleep and 10 the  
38  
39 best possible sleep) at 08:00 on the first, second and third morning after surgery.<sup>3,31</sup>  
40  
41 Other secondary endpoints include non-delirium complications within 30 days after  
42  
43 surgery, length of stay in hospital after surgery and all-cause 30-day mortality. Non-  
44  
45 delirium complications are generally defined as new-onset non-delirium conditions  
46  
47 after surgery that are harmful to patients' recovery and require therapeutic  
48  
49 intervention.  
50  
51

### 52 ***Safety outcomes***

1  
2  
3 In the present study, adverse events are monitored from the start of study drug  
4 administration until PACU discharge or 2 hours after ICU admission. Hypotension is  
5 defined as systolic blood pressure of less than 90 mmHg or a decrement of more than  
6  
7 30% from baseline. Hypertension is defined as systolic blood pressure of more than  
8  
9 180 mmHg or an increment of more than 30% from baseline. Bradycardia is defined  
10  
11 as heart rate of less than 40 beats per minute. Tachycardia is defined as heart rate of  
12  
13 more than 100 beats per minute. Desaturation is defined as SpO<sub>2</sub> of less than 90%.  
14  
15 Emergence agitation is defined as a Richmond Agitation-Sedation Scale (RASS)  
16  
17 score of more than +2 within 30 minutes after extubation. Delayed extubation is  
18  
19 defined when time to extubation is more than 2 hours (from the end of surgery) in  
20  
21 PACU patients or more than 4 hours in ICU patients.<sup>2, 22-25</sup>  
22  
23  
24  
25

26 Severe adverse events that impede patient's safety or prolong in-hospital stay will  
27  
28 be reported to Clinical Research Ethics Committee of Peking University First  
29  
30 Hospital within 24 hours. For patients who suffered harm from present trial, medical  
31  
32 treatment will be initiated as soon as possible and compensation will be completed  
33  
34 according to local laws and regulations.  
35  
36

### 37 **Data monitoring and management**

38 Original data will be recorded in the CRFs accordingly. ALL data will be kept  
39  
40 confidentially. The completed CRFs will be checked by a study coordinator who is  
41  
42 qualified by the principal investigator. Supplementations and corrections will be made  
43  
44 when necessary. Data entry will be performed in a double-input and double-check  
45  
46 way with the Data Management System (Fantastic Eight Tech. Co., Ltd, Beijing,  
47  
48 China) of the Peking University First Hospital.  
49  
50  
51

52 The conduct of the study and the quality of data will be monitored by the  
53  
54 Clinical Research Ethics Committee of Peking University First Hospital. Data  
55  
56  
57



1  
2  
3 management and statistical analysis will be performed by the Department of  
4 Biostatistics of Peking University First Hospital. Considering that dexmedetomidine  
5 has been widely used during general anesthesia and its safety has been confirmed, no  
6 interim analysis will be performed and the trial will continue until the target sample  
7 size is achieved.  
8  
9  
10  
11  
12

### 13 **Statistical analysis**

#### 14 ***Sample size calculation***

15  
16  
17 In our previous study, the incidence of delirium was 14.8% in elderly patients after  
18 non-cardiac surgery.<sup>2</sup> Previous studies reported that intraoperative dexmedetomidine  
19 decreased the incidence of POD by 60-77% in comparison with placebo.<sup>22, 25</sup> We  
20 assumed that the incidence of POD will be reduced from 14.8% to 7.4% (i.e., a 50%  
21 reduction) in the present study. With the power set at 80% and significant level at  
22 0.05, 564 patients are required to detect the difference. Considering a loss to follow-  
23 up rate of about 9%, we plan to enroll 620 patients in this study.  
24  
25  
26  
27  
28  
29  
30  
31  
32

#### 33 ***Outcome analysis***

34  
35 Continuous data with normal distribution will be compared using independent sample  
36 T-test. Continuous data with asymmetric distribution will be compared using  
37 independent sample Mann-Whitney U test. Categorical data will be compared using  
38 Chi-squared test or continuity correction Chi-squared test. The difference (and 95%  
39 confidence interval of the difference) between two means or medians will be  
40 estimated using the methodology of Levene's test or Hodges-Lehmann estimator.  
41  
42 Time-to-event data will be analyzed by survival analysis with differences between  
43 groups compared with log-rank test.  
44  
45  
46  
47  
48  
49  
50  
51

52  
53 Statistical analyses will be performed with the SPSS 14.0 (SPSS, Inc., Chicago,  
54 IL) and SAS 9.3 (SAS Institute, Cary, NC, USA). All tests are two tailed, and P  
55  
56  
57  
58  
59  
60

1  
2  
3 values of less than 0.05 are considered to be statistically significant. The Bonferroni  
4  
5 adjustment is made to control type I error for multiple testing.  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

## Discussion

This randomized, double-blinded, placebo-controlled single center trial is designed to investigate if dexmedetomidine administration during general anesthesia can decrease the incidence of POD in elderly patients after major non-cardiac surgery.

In the present study, the dosing regimen of dexmedetomidine is similar to our previous study because it does not increase drug-related adverse events (such as severe bradycardia and hypotension).<sup>24</sup> Furthermore, the CAM and CAM-ICU are used to assess delirium in patients with or without intubation, respectively.<sup>30, 32</sup> Both CAM and CAM-ICU have been translated and validated in Chinese population.<sup>33-34</sup> Feasibility of these two assessment tools have been confirmed in our previous studies.<sup>2-3, 31</sup> To maintain the quality of delirium assessment, investigators in charge of postoperative follow-up are trained by a psychiatrist before the study and will be retrained at 4-month intervals.

The strengths of the present study include the following when compared with previous studies.<sup>22-24</sup> Firstly, a randomized, double-blind, and placebo-controlled study design with a relative large sample size (620 patients) is adopted. Results of the study will provide high quality evidences. Secondly, BIS level is monitored in all enrolled patients, which will help us to avoid unnecessary and potentially harmful deep anesthesia. Thirdly, safety data will be recorded in detail.

### **Trail status:**

This study is currently at stage of patient enrollment and data collection. The current version of study protocol is V1.1 and is approved on November 27, 2015. Patient

1  
2  
3 recruitment started from December 2, 2015 and is expected to be finished by March  
4  
5 31, 2018.  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

**List of abbreviations**

1  
2  
3  
4  
5  
6  
7 POD=postoperative delirium; IL=interleukins; CRP=C-reactive protein; TNF=tumor  
8 necrosis factor; IRB=Institutional Review Board; MMSE=Mini-Mental State  
9 Examination; CAM=confusion assessment method (CAM); BIS=Bispectral index;  
10  
11 NRS=numeric rating scale; CAM-ICU=confusion assessment method-intensive care  
12  
13  
14  
15  
16 unit

For peer review only

## Declarations

**Ethics approval and consent to participate:** The study protocol was approved by Clinical Research Ethics Committee of Peking University First Hospital (2015-987). The trial was registered at Chinese Clinical Trial Registry (www.chictr.org.cn) with identifier ChiCTR-IPR-15007654 on December 1, 2015). Written informed consent will be obtained from all patients or their surrogate in law.

**Dissemination:** Results of the study will be presented at academic conferences and submitted to peer-reviewed journals.

**Competing interests:** DXW reports lecture fees and travel expenses for lectures given at academic meetings from Jiangsu Hengrui Medicine Co Ltd, China, and Yichang Humanwell Pharmaceutical Co Ltd, China. DLM is primary investigator of present study which was supported by Beijing Excellent Talent Support Program. Other authors reported no conflict of interests.

**Funding:** This trial was supported by Beijing Excellent Talent Support Program (No. 2014000020124G025). The program committee had no role in study design, data collection, management, analysis, interpretation of data, writing of the report and any other ultimate authority over any of these activities.

**Authors' contributions:** DXW and DLM designed this study. DLM draft the manuscript of protocol. DXW critically revised the manuscript. DLM, BJW, CJL, JH,

1  
2  
3 HJL, CG, ZHW and QCZ participate in the conduct of the study. All authors read and  
4  
5 approved the final manuscript.  
6  
7  
8

9 **Acknowledgements:** We thank professor Xin-Yu Sun (Department of Psychiatrics,  
10  
11 Peking University Sixth Hospital, Beijing, China) for her help in psychiatric  
12  
13 consultation and personnel training. We also thank Dr. Xue-Ying Li (Department of  
14  
15 Medical Statistic, Peking University First Hospital, Beijing, China) for her help in  
16  
17 statistical analysis of this research.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Data sharing:** No plan

## References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) Washington, DC: American Psychiatric Association; 1994.
2. Liu P, Li YW, Wang XS, et al. High serum interleukin-6 level is associated with increased risk of delirium in elderly patients after noncardiac surgery: a prospective cohort study. *Chin Med J (Engl)*. 2013; 126(19): 3621-7.
3. Su X, Meng ZT, Wu XH, et al. Dexmedetomidine for prevention of delirium in elderly patients after non-cardiac surgery: a randomised, double-blind, placebo-controlled trial. *Lancet*. 2016 Oct 15; 388(10054): 1893-1902.
4. Ely EW, Shintani A, Truman B, et al. Delirium as a predictor of mortality in mechanically ventilated patients in the intensive care unit. *JAMA*. 2004 Apr 14; 291(14): 1753-62.
5. Saczynski JS, Marcantonio ER, Quach L, et al. Cognitive trajectories after postoperative delirium. *N Engl J Med*. 2012 Jul 5; 367(1): 30-9.
6. Aldecoa C, Bettelli G, Bilotta F, et al. European Society of Anaesthesiology evidence-based and consensus-based guidelines on postoperative delirium. *Eur J Anaesthesiol*. 2017 Apr; 34(4):192-214.
7. Creeley C, Dikranian K, Dissen G, et al. Propofol-induced apoptosis of neurones and oligodendrocytes in fetal and neonatal rhesus macaque brain. *Br J Anaesth*. 2013 Jun; 110 Suppl 1: i29-38.
8. Xiong WX, Zhou GX, Wang B, et al. Impaired spatial learning and memory after sevoflurane-nitrous oxide anesthesia in aged rats is associated with down-regulated cAMP/CREB signaling. *PLoS One*. 2013 Nov 15;8(11): e79408.



- 1  
2  
3 9. Chan MT, Cheng BC, Lee TM, et al. BIS-guided anesthesia decreases  
4  
5 postoperative delirium and cognitive decline. *J Neurosurg Anesthesiol.* 2013 Jan;  
6  
7 25(1): 33-42.  
8
- 9  
10 10. Vaurio LE, Sands LP, Wang Y, et al. Postoperative delirium: the importance of  
11  
12 pain and pain management. *Anesth Analg.* 2006 Apr; 102(4): 1267-73.  
13
- 14 11. Lynch EP, Lazor MA, Gellis JE, et al. The impact of postoperative pain on the  
15  
16 development of postoperative delirium. *Anesth Analg* 1998; 86:781-5  
17
- 18 12. Hirsch J, Vacas S, Terrando N, et al. Perioperative cerebrospinal fluid and  
19  
20 plasma inflammatory markers after orthopedic surgery. *J Neuroinflammation.*  
21  
22 2016 Aug 30; 13(1): 211.  
23
- 24 13. Cape E, Hall RJ, van Munster BC, et al. Cerebrospinal fluid markers of  
25  
26 neuroinflammation in delirium: a role for interleukin-1 $\beta$  in delirium after hip  
27  
28 fracture. *J Psychosom Res.* 2014 Sep; 77(3): 219-25.  
29
- 30  
31 14. van den Boogaard M, Kox M, Quinn KL, et al. Biomarkers associated with  
32  
33 delirium in critically ill patients and their relation with long-term subjective  
34  
35 cognitive dysfunction; indications for different pathways governing delirium in  
36  
37 inflamed and noninflamed patients. *Crit Care.* 2011; 15(6): R297.  
38
- 39  
40 15. McGrane S, Girard TD, Thompson JL, et al. Procalcitonin and C-reactive protein  
41  
42 levels at admission as predictors of duration of acute brain dysfunction in  
43  
44 critically ill patients. *Crit Care.* 2011; 15(2): R78.  
45
- 46 16. Kazmierski J, Banys A, Latek J, et al. Raised IL-2 and TNF- $\alpha$  concentrations are  
47  
48 associated with postoperative delirium in patients undergoing coronary-artery  
49  
50 bypass graft surgery. *Int Psychogeriatr.* 2014 May; 26(5): 845-55.  
51
- 52  
53 17. Shin HW, Yoo HN, Kim DH, et al. Preanesthetic dexmedetomidine 1 $\mu$ g/kg  
54  
55 single infusion is a simple, easy, and economic adjuvant for general anesthesia.  
56  
57

- 1  
2  
3 Korean J Anesthesiol. 2013 Aug; 65(2): 114-20.  
4  
5 18. Li J, Xiong M, Nadavaluru PR, et al. Dexmedetomidine attenuates neurotoxicity  
6 induced by prenatal propofol exposure. J Neurosurg Anesthesiol. 2016 Jan;  
7 28(1): 51-64.  
8  
9  
10  
11 19. Sanders RD, Xu J, Shu Y, et al. Dexmedetomidine attenuates isoflurane-induced  
12 neurocognitive impairment in neonatal rats. Anesthesiology. 2009 May; 110(5):  
13 1077-85.  
14  
15  
16  
17 20. Le Bot A, Michelet D, Hilly J, et al. Efficacy of intraoperative dexmedetomidine  
18 compared with placebo for surgery in adults: a meta-analysis of published  
19 studies. Minerva Anesthesiol. 2015 Oct; 81(10): 1105-17.  
20  
21  
22  
23 21. Kang SH, Kim YS, Hong TH, et al. Effects of dexmedetomidine on  
24 inflammatory responses in patients undergoing laparoscopic cholecystectomy.  
25 Acta Anaesthesiol Scand. 2013 Apr; 57(4): 480-7.  
26  
27  
28  
29 22. Li Y, Wang B, Zhang LL, et al. Dexmedetomidine Combined with General  
30 Anesthesia Provides Similar Intraoperative Stress Response Reduction When  
31 Compared with a Combined General and Epidural Anesthetic Technique. Anesth  
32 Analg. 2016 Apr; 122(4): 1202-10  
33  
34  
35  
36 23. Cao JL, Pei YP, Wei JQ, et al. Effects of intraoperative dexmedetomidine with  
37 intravenous anesthesia on postoperative emergence agitation/delirium in  
38 pediatric patients undergoing tonsillectomy with or without adenoidectomy: A  
39 CONSORT-prospective, randomized, controlled clinical trial. Medicine  
40 (Baltimore). 2016 Dec; 95(49): e5566.  
41  
42  
43  
44 24. Sun Y, Liu J, Yuan X, et al. Effects of dexmedetomidine on emergence delirium  
45 in pediatric cardiac surgery. Minerva Pediatr. 2017 Jun; 69(3): 165-173.  
46  
47  
48  
49 25. Li X, Yang J, Nie XL, et al. Impact of dexmedetomidine on the incidence of  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2 delirium in elderly patients after cardiac surgery: A randomized controlled trial.  
3 PLoS One. 2017 Feb 9; 12(2): e0170757.  
4  
5  
6  
7 26. Yang X, Li Z, Gao C, et al. Effect of dexmedetomidine on preventing agitation  
8 and delirium after microvascular free flap surgery: a randomized, double-blind,  
9 control study. *J Oral Maxillofac Surg*. 2015 Jun; 73(6): 1065-72.  
10  
11  
12  
13 27. Deiner S, Luo X, Lin HM, et al. Intraoperative Infusion of Dexmedetomidine for  
14 Prevention of Postoperative Delirium and Cognitive Dysfunction in Elderly  
15 Patients Undergoing Major Elective Noncardiac Surgery: A Randomized Clinical  
16 Trial. *JAMA Surg*. 2017 Aug 16; 152(8): e171505.  
17  
18  
19  
20  
21  
22 28. Collin C, Wade DT, Davies S, et al. The Barthel ADL Index: a reliability study.  
23 *Int Disabil Stud*. 1988; 10: 61-3.  
24  
25  
26 29. Katzman R, Zhang MY, Ouang-Ya-Qu, et al. A Chinese version of the Mini-  
27 Mental State Examination; impact of illiteracy in a Shanghai dementia survey. *J*  
28 *Clin Epidemiol*. 1988; 41: 971-8.  
29  
30  
31  
32  
33 30. Inouye SK, van Dyck CH, Alessi CA, et al. Clarifying confusion: The Confusion  
34 Assessment Method. A new method for detecting delirium. *Ann Intern Med*  
35 1990; 113:941-8  
36  
37  
38  
39 31. Mu DL, Zhang DZ, Wang DX, et al. Parecoxib Supplementation to Morphine  
40 Analgesia Decreases Incidence of Delirium in Elderly Patients After Hip or Knee  
41 Replacement Surgery: A Randomized Controlled Trial. *Anesth Analg*. 2017 Jun;  
42 124(6):1992-2000.  
43  
44  
45  
46  
47  
48 32. Ely EW, Margolin R, Francis J, et al. Evaluation of delirium in critically ill  
49 patients: validation of the Confusion Assessment Method for the Intensive Care  
50 Unit (CAM-ICU). *Crit Care Med*. 2001 Jul; 29(7): 1370-9.  
51  
52  
53  
54  
55 33. Ji L, Ve L, Leung CM, et al. Clinical utility and validation of two instruments  
56  
57  
58  
59  
60

1  
2  
3 (the Confusion Assessment Method Algorithm and the Chinese version of  
4 Nursing Delirium Screening Scale) to detect delirium in geriatric inpatients. Gen  
5 Hosp Psychiatry. 2008; 30: 171-6.  
6  
7

- 8  
9 34. Wang C, Wu Y, Yue P, et al. Delirium assessment using Confusion Assessment  
10 Method for the Intensive Care Unit in Chinese critically ill patients. J Crit Care.  
11 2013; 28: 223-9.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


For peer review only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Figure legend**

Figure 1. Flowchart of this study.

For peer review only

**Figure 1 Flow chart of this study**



SPIRIT 2013 Checklist: Recommended items to address in a clinical trial protocol and related documents\*

Section/item	Item No	Description	Addressed on page number
<b>Administrative information</b>			
Title	1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	1
Trial registration	2a	Trial identifier and registry name. If not yet registered, name of intended registry	4 , 8, 19
	2b	All items from the World Health Organization Trial Registration Data Set	8
Protocol version	3	Date and version identifier	8
Funding	4	Sources and types of financial, material, and other support	19
Roles and responsibilities	5a	Names, affiliations, and roles of protocol contributors	19
	5b	Name and contact information for the trial sponsor	19
	5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities	19
	5d	Composition, roles, and responsibilities of the coordinating centre, steering committee, endpoint adjudication committee, data management team, and other individuals or groups overseeing the trial, if applicable (see Item 21a for data monitoring committee)	N/A

**Introduction**

1				
2				
3	Background and	6a	Description of research question and justification for undertaking the trial, including summary of relevant	6-7
4	rationale		studies (published and unpublished) examining benefits and harms for each intervention	
5				
6		6b	Explanation for choice of comparators	6-7, 10-11
7	Objectives	7	Specific objectives or hypotheses	7
8				
9	Trial design	8	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group),	
10			allocation ratio, and framework (eg, superiority, equivalence, noninferiority, exploratory)	8
11				
12				
13	<b>Methods: Participants, interventions, and outcomes</b>			
14				
15	Study setting	9	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will	8
16			be collected. Reference to where list of study sites can be obtained	
17				
18	Eligibility criteria	10	Inclusion and exclusion criteria for participants. If applicable, eligibility criteria for study centres and	8-9
19			individuals who will perform the interventions (eg, surgeons, psychotherapists)	
20				
21	Interventions	11a	Interventions for each group with sufficient detail to allow replication, including how and when they will be	10-11
22			administered	
23				
24		11b	Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose	10-11
25			change in response to harms, participant request, or improving/worsening disease)	
26				
27		11c	Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence	10-11
28			(eg, drug tablet return, laboratory tests)	
29				
30		11d	Relevant concomitant care and interventions that are permitted or prohibited during the trial	N/A
31				
32	Outcomes	12	Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood	
33			pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg,	11-13
34			median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen	
35			efficacy and harm outcomes is strongly recommended	
36				
37	Participant timeline	13	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for	Figure 1
38			participants. A schematic diagram is highly recommended (see Figure)	
39				
40				
41				
42				
43				
44				
45				
46				
47				



1				
2				
3	Sample size	14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	14
4				
5	Recruitment	15	Strategies for achieving adequate participant enrolment to reach target sample size	9
6				
7				
8	<b>Methods: Assignment of interventions (for controlled trials)</b>			
9				
10	Allocation:			
11				
12	Sequence generation	16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	9
13				
14				
15				
16				
17	Allocation concealment mechanism	16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	9-10
18				
19				
20				
21	Implementation	16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	9-10
22				
23				
24	Blinding (masking)	17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	9-10
25				
26				
27		17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	9-10
28				
29				
30				
31	<b>Methods: Data collection, management, and analysis</b>			
32				
33	Data collection methods	18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known. Reference to where data collection forms can be found, if not in the protocol	9-13
34				
35				
36				
37				
38		18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols	11-13
39				
40				
41				
42				
43				
44				
45				
46				
47				

1				
2				
3	Data management	19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	13-14
4				
5				
6				
7	Statistical methods	20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	14-15
8				
9				
10		20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	14-15
11				
12		20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any statistical methods to handle missing data (eg, multiple imputation)	14-15
13				
14				

### 15 **Methods: Monitoring**

16				
17	Data monitoring	21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent from the sponsor and competing interests; and reference to where further details about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not needed	13
18				
19				
20				
21				
22		21b	Description of any interim analyses and stopping guidelines, including who will have access to these interim results and make the final decision to terminate the trial	13-14
23				
24				
25	Harms	22	Plans for collecting, assessing, reporting, and managing solicited and spontaneously reported adverse events and other unintended effects of trial interventions or trial conduct	13
26				
27				
28	Auditing	23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent from investigators and the sponsor	13-14
29				
30				
31				

### 32 **Ethics and dissemination**

33				
34	Research ethics approval	24	Plans for seeking research ethics committee/institutional review board (REC/IRB) approval	4, 8, 19
35				
36				
37	Protocol amendments	25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes, analyses) to relevant parties (eg, investigators, REC/IRBs, trial participants, trial registries, journals, regulators)	8
38				
39				
40				
41				
42				
43				
44				
45				
46				
47				

1				
2				
3	Consent or assent	26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	8-9
4				
5				
6		26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	N/A
7				
8	Confidentiality	27	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	13
9				
10				
11	Declaration of interests	28	Financial and other competing interests for principal investigators for the overall trial and each study site	19
12				
13				
14	Access to data	29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	13-14
15				
16				
17	Ancillary and post-trial care	30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	13
18				
19				
20	Dissemination policy	31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	19
21				
22				
23				
24				
25		31b	Authorship eligibility guidelines and any intended use of professional writers	19-20
26				
27		31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	N/A
28				
29	<b>Appendices</b>			
30				
31	Informed consent materials	32	Model consent form and other related documentation given to participants and authorised surrogates	N/A
32				
33				
34	Biological specimens	33	Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies, if applicable	N/A
35				
36				

37 \*It is strongly recommended that this checklist be read in conjunction with the SPIRIT 2013 Explanation & Elaboration for important clarification on the items.  
 38 Amendments to the protocol should be tracked and dated. The SPIRIT checklist is copyrighted by the SPIRIT Group under the Creative Commons  
 39 "[Attribution-NonCommercial-NoDerivs 3.0 Unported](https://creativecommons.org/licenses/by-nc-nd/3.0/)" license.  
 40

# BMJ Open

## Impact of dexmedetomidine infusion during general anesthesia on incidence of postoperative delirium in elderly patients after major non-cardiac surgery: study protocol of a randomized, double-blinded and placebo-controlled trial

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-019549.R1
Article Type:	Protocol
Date Submitted by the Author:	13-Dec-2017
Complete List of Authors:	Wang, Bojie; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Li, Chunjing; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Hu, Jian; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Li, Huaijin; Peking University First Hospital, Department of Anesthesiology and Critical Care Guo, Chao; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Wang, Zhenhan; Dongping People's Hospital, Department of anesthesiology Zhang, Qiaochu; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Mu, Dongliang; Department of Anesthesiology and Critical Care Medicine Wang, Dong-Xin; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine
<b>Primary Subject Heading</b>:	Anaesthesia
Secondary Subject Heading:	Anaesthesia
Keywords:	Adult anaesthesia < ANAESTHETICS, Delirium & cognitive disorders < PSYCHIATRY, Pain management < RADIOTHERAPY

SCHOLARONE™  
Manuscripts

1  
2  
3 **1 Impact of dexmedetomidine infusion during general anesthesia on incidence of**  
4  
5 **2 postoperative delirium in elderly patients after major non-cardiac surgery: study**  
6  
7 **3 protocol of a randomized, double-blinded and placebo-controlled trial**  
8  
9  
10

11 5 Bo-Jie Wang<sup>1</sup>, Chun-Jing Li<sup>1</sup>, Jian Hu<sup>1</sup>, Huai-Jin Li<sup>1</sup>, Chao Guo<sup>1</sup>, Zhen-Han Wang<sup>2</sup>,  
12  
13 6 Qiao-Chu Zhang<sup>1</sup>, Dong-Liang Mu<sup>1,\*</sup>, Dong-Xin Wang<sup>1</sup>  
14  
15

16 7  
17  
18 8 <sup>1</sup>Department of Anesthesiology and Critical Care Medicine, Peking University First  
19  
20 9 Hospital, Beijing, China

21  
22 10 <sup>2</sup>Department of Anesthesiology, Dongping People's Hospital, Shandong, China  
23  
24

25  
26  
27 12 The first two authors (BJ Wang and CJ Li) contributed equally to this manuscript  
28  
29

30  
31 14 Corresponding author: Dong-Liang Mu, M.D., Department of Anesthesiology and  
32  
33 15 Critical Care Medicine, Peking University First Hospital, Beijing, China. Tel: +86  
34  
35 16 010 83572784 (O), +86 13810702725 (M); E-mail: mudongliang@icloud.com  
36  
37

38  
39 18 Bo-Jie Wang: ivyxbd@hotmail.com  
40

41  
42 19 Chun-Jing Li: gogolion@163.com  
43

44  
45 20 Jian Hu: hj19871204@163.com  
46

47  
48 21 Huai-Jin Li: sophie.lee.coffee@gmail.com  
49

50  
51 22 Chao Guo: guochao1988@yeah.com  
52

53  
54 23 Zhen-Han Wang: 1468432914@qq.com  
55

56  
57 24 Qiao-Chu Zhang: 372193745@qq.com  
58

59  
60 25 Dong-Liang Mu: mudongliang@icloud.com

1  
2  
3 1 Dong-Xin Wang: wangdongxin@hotmail.com  
4

5 2  
6

7 3 Word Count: 2843  
8

9 4  
10

11 5  
12

13 6  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

## Abstract for protocol

**Introduction:** Delirium is a common complication in the elderly after surgery and is associated with worse outcomes. Multiple risk factors are related with postoperative delirium, such as exposure to general anesthetics, pain and inflammatory response. Preclinical and clinical studies reported that dexmedetomidine could attenuate neurotoxicity induced by general anesthetics, improve pain management and inhibit inflammatory response. Several studies observed the relationship between intraoperative use of dexmedetomidine and postoperative delirium, but the results were inconsistent. This study is designed to investigate the impact of dexmedetomidine administered during general anesthesia in preventing delirium in elderly patients after major non-cardiac surgery.

**Method and analysis:** This is a randomized, double-blinded, and placebo-controlled trial. 620 elderly patients (age  $\geq 60$  years) who are scheduled to undertake elective major non-cardiac surgery (with an expected duration  $\geq 2$  hours) are randomly divided into two groups. For patients in dexmedetomidine group, a loading dose dexmedetomidine (0.6  $\mu\text{g}/\text{kg}$ ) will be administered in 10 minutes before anesthesia induction, followed by a continuous infusion at a rate of 0.5  $\mu\text{g}/\text{kg}/\text{h}$  till 1 hour before the end of surgery. For patients in control group, normal saline will be administered in the same rate and volume as in the dexmedetomidine group. The primary endpoint is the incidence of delirium during the first five postoperative days. Secondary endpoints include pain intensity, cumulative opioid consumption and subjective sleep quality during the first 3 postoperative days, and the incidence of non-delirium complications and all-cause mortality within 30 days after surgery.

1  
2  
3 1 **Ethics and dissemination:** The study protocol was approved by Clinical Research  
4  
5 2 Ethics Committee of Peking University First Hospital (2015-987) and registered at  
6  
7 3 Chinese Clinical Trial Registry (<http://www.chictr.org.cn>) with identifier ChiCTR-  
8  
9 4 IPR-15007654. Results of the study will be presented at academic conferences and  
10  
11 5 submitted to peer-reviewed journals.  
12

13  
14 6

15  
16 7 **Key words:** intraoperative dexmedetomidine; postoperative delirium; old patient;  
17  
18 8 major non-cardiac surgery  
19

20  
21 9  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## Strengths and limitations

1. The study will investigate the impact of dexmedetomidine infusion during general anesthesia on the incidence of delirium in elderly patients after non-cardiac surgery.
2. The study designed will be a randomized, double-blinded and placebo-controlled, with a relative large sample size.
3. Anesthesia depth (Bispectral index) will be monitored to guide anesthesia maintenance. This will overcome an important limitation in a recent trial.
4. One limitation is that this is a single center trial, which will limit the generalizability of our results. Another limitation is that only early outcomes (up to 30 days after surgery) will be explored.

## Introduction

Delirium is a transient brain dysfunction which is characterized by altered consciousness, inattention, and changes in cognition or perception; it develops acutely with clinical manifestations fluctuate during the course of the day.<sup>1</sup> Prevalence of delirium varies from 12% to 51% in patients after non-cardiac surgery, and its prevalence increases with age.<sup>2-3</sup> The occurrence of postoperative delirium (POD) is associated with worsen outcomes including prolonged mechanical ventilation and intensive care unit stay, increased postoperative complications, high mortality rate and long-term cognitive decline.<sup>2-5</sup>

The etiology of POD is multifactorial and includes several intraoperative factors.<sup>6</sup> For example, exposure to general anesthetics (e.g., propofol or sevoflurane) might produce neurotoxicity,<sup>7-8</sup> whereas reduced anesthetic exposure by avoiding deep anesthesia reduced the occurrence of delirium in elderly patients undergoing major non-cardiac surgery.<sup>9</sup> Poor pain management is another risk factor of POD.<sup>6, 10</sup> It was reported that the risk of postoperative delirium was 1.2 times higher for every unit increment in visual analogue pain score (an 11-point pain scale where 0 indicates no pain and 10 the most severe pain).<sup>11</sup> Inflammation is also proposed to play an important role in the pathogenesis of POD.<sup>2</sup> Inflammatory responses induced by surgery and anesthesia are manifested by elevated levels of interleukins (IL), C-reactive protein (CRP) and tumor necrosis factor (TNF).<sup>2, 12</sup> Studies by our group and other found that higher levels of inflammatory mediators are associated with increased risk of POD.<sup>2, 13-16.</sup>

Dexmedetomidine is a highly selective  $\alpha_2$ -receptor agonist with sedative, analgesic and anxiolytic effects.<sup>17-21</sup> When used as a supplement during intraoperative

1 anesthesia, it reduces the consumption of general anesthetics.<sup>17</sup> Preclinical evidences  
2 suggest that use of dexmedetomidine might attenuate neurotoxicity induced by  
3 general anesthetics.<sup>18-19</sup> In a meta-analysis, intraoperative administration of  
4 dexmedetomidine lowers postoperative pain intensity and reduces opioid  
5 consumption.<sup>20</sup> Clinical evidence also showed that intraoperative dexmedetomidine  
6 significantly inhibits hyper-secretion of inflammatory cytokines during and after  
7 surgery.<sup>21-22</sup>

8       Use of dexmedetomidine during general anesthesia may reduce POD. In  
9 pediatric patients undergoing tonsillectomy and cardiac surgery, intraoperative  
10 infusion of dexmedetomidine lowered the incidence of emergence delirium.<sup>23,24</sup> In  
11 adult patients undergoing cardiac surgery and microvascular free flap surgery,  
12 intraoperative dexmedetomidine (comparison with normal saline) slightly decreased  
13 the incidence of delirium, although the differences were not statistically significant  
14 between two groups possibly due to underpowered sample size.<sup>25,26</sup> In a recent study  
15 of Deiner et al.,<sup>27</sup> use of dexmedetomidine during general anesthesia did not reduce  
16 delirium after major non-cardiac surgery in the elderly. However, in that study,  
17 anesthesia depth was not monitored and the consumption of anesthetics (such as  
18 propofol and fentanyl) was similar between the two groups. It was possible that  
19 patients in the dexmedetomidine group received deeper anesthesia which might have  
20 increased the risk of delirium.<sup>27</sup> Therefore, the effects of dexmedetomidine  
21 administered during general anesthesia on the occurrence of POD need to be  
22 evaluated further.

23       This study is designed to investigate whether dexmedetomidine use during  
24 general anesthesia can decrease the incidence of POD in elderly patients after major  
25 non-cardiac surgery.

## Method and analysis

### Study design

This randomized, double-blinded, and placebo-controlled trial with two parallel arms is designed to test the superiority of dexmedetomidine administered during general anesthesia on the incidence of delirium after surgery. Patients will be randomized into either the dexmedetomidine group or the control group (Figure 1). The study is conducted in the Department of Anesthesiology and Critical Care Medicine of Peking University First Hospital.

### Ethics approval

The study protocol (version 1.1, issue date Nov, 2015) was approved by Clinical Research Ethics Committee of Peking University First Hospital (2015-987). Any protocol modifications will be submitted for review and approval by the Ethics Committee. The trial was registered at Chinese Clinical Trial Registry (<http://www.chictr.org.cn>) on December 1, 2015, with identifier ChiCTR-IPR-15007654. Written informed consents are obtained from patients or their surrogate in law.

### Participants

Elderly (age  $\geq 60$  years) patients who are scheduled to undergo elective non-cardiac surgery with expected duration  $\geq 2$  hours under general anesthesia are screened for inclusion. Those who meet any of the following criteria will be excluded: (1) do not provide written informed consents; (2) previous history of schizophrenia, epilepsy or Parkinson disease; (3) visual, hearing, language or other barrier which impede communication and preoperative delirium assessment; (4) neurosurgery or traumatic brain injury; (5) severe bradycardia (heart rate less than 40 beats per minute), sick

1 sinus syndrome or atrioventricular block of degree 2 or above; (6) severe hepatic  
2 dysfunction (Child-Pugh grade C); or (7) renal failure (requirement of renal  
3 replacement therapy).

#### 4 **Patient recruitment and baseline data collection**

5 Potential participants are screened by qualified investigators the day before surgery  
6 (or on Friday for those who will undergo surgery next Monday). The study protocol  
7 including potential risks and benefits is explained in detail. Those who meet the  
8 inclusion and exclusion criteria are invited to participate in the study.

9 After obtaining written informed consents, the following baseline data are  
10 collected: demographic data, preoperative diagnosis, comorbidity, current medical  
11 therapy, previous surgery, and main results of physical and laboratory examinations.  
12 Barthel index is used for evaluation of activities of daily living.<sup>28</sup> Cognitive function  
13 is assessed with Mini-Mental State Examination (MMSE).<sup>29</sup> Preoperative delirium is  
14 assessed with confusion assessment method (CAM).<sup>30</sup>

#### 15 **Randomization, grouping and blinding**

16 Random numbers were created by an independent statistician using SAS statistical  
17 package version 9.3 (SAS Institute, Cary, NC, USA) in a 1:1 ratio and were sealed in  
18 sequentially numbered envelopes. A study coordinator, who has no knowledge of  
19 patients before randomization and does not participate in anesthesia and postoperative  
20 follow-up of enrolled patients, will open envelop for random numbers and prepare  
21 study drugs before induction of anesthesia. In this way, patients are randomly divided  
22 into either dexmedetomidine or control group.

23 Information regarding randomization, study drug preparation and group  
24 allocation will be masked from investigators who perform data collection and

1 postoperative follow-up, anesthesiologists, patients and other healthcare team  
2 members. Blinding will be maintained throughout the entire study period.

3 To ensure patients' safety, study group allocation can be unmasked in the  
4 following conditions, i.e., occurrence of severe adverse events or any unexpected  
5 deterioration in the patient's clinical status. These situations will be documented in  
6 the Case Report Forms (CRFs). The unmasked patients will be included in the  
7 intention-to-treat population but excluded from per-protocol analysis.

### 8 **Interventions, anesthesia and analgesia**

9 Intraoperative monitoring includes electrocardiogram, noninvasive blood pressure,  
10 pulse oxygen saturation, end-tidal carbon dioxide, nasopharyngeal temperature, urine  
11 output, and bispectral index (BIS). Intraarterial pressure (including derivative  
12 dynamic parameters such as stroke volume variation by FlowTrac system) and central  
13 venous pressure are monitored according to patients' conditions.

14 Study drugs, either 200 µg (2 ml) dexmedetomidine (Jiangsu Hengrui Medicine  
15 Co, Ltd, Jiangsu, China) or 2 ml normal saline, are diluted in 50 ml normal saline. All  
16 study drugs are colorless solution provided in syringes of the same size and brand.  
17 The regimen of study drug administration includes a loading dose of 0.15 ml/kg (i.e.,  
18 0.6 µg/kg dexmedetomidine for patients in the dexmedetomidine group) administered  
19 during a 10-minute period before anesthesia induction. This is followed by a  
20 continuous infusion at a rate of 0.125 ml/kg/h (i.e., a rate of 0.5 µg/kg/h  
21 dexmedetomidine for patients in the dexmedetomidine group) till 1 hour before the  
22 end of surgery. Study drug infusion is performed by an infusion pump specially  
23 designed for dexmedetomidine administration (Slgo® CP1000, Beijing Slgo medical  
24 technology Co., Ltd.).

1  
2  
3 1 Attending anesthesiologists can decrease or stop study drug infusion in the  
4  
5 2 following conditions: (1) severe bradycardia or hypotension which does not improve  
6  
7 3 after routine treatment; (2) new onset atrioventricular block which does not improve  
8  
9 4 after routine treatment; or (3) other conditions that anesthesiologists consider it  
10  
11 5 necessary. In these conditions the reasons that lead to any protocol deviations will be  
12  
13 6 recorded in the Case Report Forms. These patients will be included in the intention-  
14  
15 7 to-treat analysis but excluded from the per-protocol analysis.

18 8 Total intravenous anesthesia is performed for all patients. Anesthesia is induced  
19  
20 9 with intravenous sufentanil (target controlled infusion with effect-site concentration  
21  
22 10 from 0.2 to 0.5 ng/ml) and propofol (2-3 mg/kg) and maintained with sufentanil  
23  
24 11 (effect-site concentration from 0.2 to 0.5 ng/ml) and propofol (4-12 mg/kg/h).  
25  
26 12 Rocuronium and/or cisatracurium are administered for muscle relaxation. The mean  
27  
28 13 arterial pressure is maintained above 60 mmHg or within 20% from baseline. BIS is  
29  
30 14 maintained between 40 and 60. Body temperature is maintained with air-warming and  
31  
32 15 fluid heating systems. The target of nasopharyngeal temperature maintenance during  
33  
34 16 surgery is from 36.0 to 37°C. Mechanical ventilated is performed with a 1:1 nitrous  
35  
36 17 oxide-oxygen mixture.

39 18 All patients are transferred to post-anesthesia care unit (PACU) or intensive care  
40  
41 19 unit (ICU) before they are sent back to general wards. Patient-controlled intravenous  
42  
43 20 analgesia (PCIA) is provided for all patients, which is established with 0.5 mg/ml  
44  
45 21 morphine in 100 ml normal saline and programmed to deliver a 1 mg bolus with a  
46  
47 22 lock-out interval of 8 minutes and a background infusion at 0.5 mg/h. Supplemental  
48  
49 23 morphine at dose of 2 to 4 mg will be administered at 10-minute intervals if the  
50  
51 24 numeric rating scale (NRS) pain score (a 11-point scale where 0 indicates no pain and  
52  
53  
54  
55  
56  
57  
58  
59  
60

10 indicates the worst pain) remains above 4 after 3 consecutive PCIA boluses.<sup>31</sup>

Other postoperative managements were performed according to routine practice.

### **Outcome assessment**

Patients are followed up twice daily during the first 5 postoperative days and then weekly until 30 days after surgery. Investigators who are responsible for postoperative follow-up are not involved in anesthesia and perioperative care, and are not allowed to exchange patients' information with anesthesiologists who take care of patients in the operating room. Before the beginning of the study, investigators are trained to follow the study protocol and to perform delirium assessment and the training process is repeated at 4 to 6 months intervals during the study period.<sup>2-3, 31</sup>

The 4-hour training courses of delirium assessment include the following contents: (1) lectures regarding signs/symptoms, diagnosis and treatment of delirium by psychiatrists; (2) training courses of the use of CAM and CAM-ICU on patient-actors (trained ICU physicians or nurses who act as patients with or without delirium) conducted by psychiatrists. The process continued until 100% agreement is achieved in diagnosing delirium.

### **Primary endpoint**

The primary endpoint is the incidence of delirium during the first 5 days after surgery. Delirium is assessed twice daily (at 08:00-09:00 and 19:00-20:00, respectively) with CAM for non-intubated patients<sup>30</sup> or CAM for the Intensive Care Unit (CAM-ICU) for intubated patients.<sup>32</sup> These delirium assessment methods had been used in our previous studies.<sup>2-3, 31</sup> For patients who are discharged or died within 5 days after surgery, the results of last delirium assessment will be considered the results of the missing data. These patients will be excluded when calculating daily prevalence of delirium in a post-hoc analysis.



### 1 **Secondary endpoints**

2 Postoperative pain intensities at rest and with movement are assessed with NRS pain  
3 score at 24, 48 and 72 hours after surgery, respectively.<sup>31</sup> Cumulative morphine  
4 consumptions at these time points are recorded. Subjective sleep quality is assessed  
5 with NRS (an 11-point scale where 0 indicates the worst possible sleep and 10 the  
6 best possible sleep) at 08:00 on the first, second and third morning after surgery.<sup>3,31</sup>  
7 Other secondary endpoints include non-delirium complications within 30 days after  
8 surgery, length of stay in hospital after surgery and all-cause 30-day mortality. Non-  
9 delirium complications are generally defined as new-onset non-delirium conditions  
10 after surgery that are harmful to patients' recovery and require therapeutic  
11 intervention.

### 12 **Safety outcomes**

13 In the present study, adverse events are monitored from the start of study drug  
14 administration until PACU discharge or 2 hours after ICU admission. Hypotension is  
15 defined as systolic blood pressure of less than 90 mmHg or a decrement of more than  
16 30% from baseline. Hypertension is defined as systolic blood pressure of more than  
17 180 mmHg or an increment of more than 30% from baseline. Bradycardia is defined  
18 as heart rate of less than 40 beats per minute. Tachycardia is defined as heart rate of  
19 more than 100 beats per minute. Desaturation is defined as SpO<sub>2</sub> of less than 90%.  
20 Emergence agitation is defined as a Richmond Agitation-Sedation Scale (RASS)  
21 score of more than +2 within 30 minutes after extubation. Delayed extubation is  
22 defined when time to extubation is more than 2 hours (from the end of surgery) in  
23 PACU patients or more than 4 hours in ICU patients.<sup>2, 22-25</sup>

24 Severe adverse events, i.e., those that might result in patient's  
25 disability/deformity, prolong in-hospital stay, or life threatening events, will be

1 reported to Clinical Research Ethics Committee of Peking University First Hospital  
2 within 24 hours. For patients who suffered harm from present trial, medical treatment  
3 will be initiated as soon as possible and compensation will be completed according to  
4 local laws and regulations.

#### 5 **Data monitoring and management**

6 Original data will be recorded in the CRFs accordingly. ALL data will be kept  
7 confidentially. The completed CRFs will be checked by a study coordinator who is  
8 qualified by the principal investigator. Supplementations and corrections will be made  
9 when necessary. Data entry will be performed in a double-input and double-check  
10 way with the Data Management System (Fantastic Eight Tech. Co., Ltd, Beijing,  
11 China) of the Peking University First Hospital.

12 The conduct of the study and the quality of data will be monitored by the  
13 Clinical Research Ethics Committee of Peking University First Hospital. Data  
14 management and statistical analysis will be performed by the Department of  
15 Biostatistics of Peking University First Hospital. Considering that dexmedetomidine  
16 has been widely used during general anesthesia and its safety has been confirmed, no  
17 interim analysis will be performed and the trial will continue until the target sample  
18 size is achieved.

#### 19 **Statistical analysis**

##### 20 ***Sample size calculation***

21 In our previous study, the incidence of delirium was 14.8% in elderly patients after  
22 non-cardiac surgery.<sup>2</sup> Previous studies reported that intraoperative dexmedetomidine  
23 decreased the incidence of POD by 60-77% in comparison with placebo.<sup>22, 25</sup> We  
24 assumed that the incidence of POD will be reduced from 14.8% to 7.4% (i.e., a 50%  
25 reduction) in the present study. With the power set at 80% and significant level at

1 0.05, 564 patients are required to detect the difference. Considering a loss to follow-  
2 up rate of about 9%, we plan to enroll 620 patients in this study.

### 3 ***Outcome analysis***

4 Continuous data with normal distribution will be compared using independent sample  
5 T-test. Continuous data with asymmetric distribution will be compared using  
6 independent sample Mann-Whitney U test. Categorical data will be compared using  
7 Chi-squared test or continuity correction Chi-squared test. The difference (and 95%  
8 confidence interval of the difference) between two means or medians will be  
9 estimated using the methodology of Levene's test or Hodges-Lehmann estimator.  
10 Time-to-event data will be analyzed by survival analysis with differences between  
11 groups compared with log-rank test.

12 Statistical analyses will be performed with the SPSS 14.0 (SPSS, Inc., Chicago,  
13 IL) and SAS 9.3 (SAS Institute, Cary, NC, USA). All tests are two tailed, and P  
14 values of less than 0.05 are considered to be statistically significant. The Bonferroni  
15 adjustment is made to control type I error for multiple testing.

16

## Discussion

This randomized, double-blinded, placebo-controlled single center trial is designed to investigate if dexmedetomidine administration during general anesthesia can decrease the incidence of POD in elderly patients after major non-cardiac surgery.

In the present study, the dosing regimen of dexmedetomidine is similar to our previous study because it does not increase drug-related adverse events (such as severe bradycardia and hypotension).<sup>24</sup> Furthermore, the CAM and CAM-ICU are used to assess delirium in patients with or without intubation, respectively.<sup>30, 32</sup> Both CAM and CAM-ICU have been translated and validated in Chinese population.<sup>33-34</sup> Feasibility of these two assessment tools have been confirmed in our previous studies.<sup>2-3, 31</sup> To maintain the quality of delirium assessment, investigators in charge of postoperative follow-up are trained by a psychiatrist before the study and will be retrained at 4 to 6-month intervals.

Because of the hemodynamic and anesthetic-sparing effect of dexmedetomidine, it is not very difficult for the experienced anesthesiologists to guess which study drug is administered. This might weak the blinding to anesthesiologists. However, in the present study, investigators who are responsible for postoperative follow-up and delirium assessment are not involved in anesthesia and perioperative care; and they are not allowed to exchange patients' information with anesthesiologists who take care of patients in the operating room. In this way the blinding of investigators to study group assignment can be guaranteed.

The strengths of the present study include the following when compared with previous studies.<sup>22-24</sup> Firstly, a randomized, double-blind, and placebo-controlled study design with a relative large sample size (620 patients) is adopted. Results of the

1 study will provide high quality evidences. Secondly, BIS level is monitored in all  
2 enrolled patients, which will help us to avoid unnecessary and potentially harmful  
3 deep anesthesia. Thirdly, safety data will be recorded in detail. Our study also has  
4 some limitations. One is that this is a single center trial, which will limit the external  
5 validity of our results. Another one is that only early outcomes (up to 30 days after  
6 surgery) will be explored.

7  
8 **Trail status:**

9 This study is currently at stage of patient enrollment and data collection. The current  
10 version of study protocol is V1.1 and is approved on November 27, 2015. Patient  
11 recruitment started from December 2, 2015 and is expected to be finished by March  
12 31, 2018.

13

**List of abbreviations**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8

POD=postoperative delirium; IL=interleukins; CRP=C-reactive protein; TNF=tumor  
necrosis factor; IRB=Institutional Review Board; MMSE=Mini-Mental State  
Examination; CAM=confusion assessment method (CAM); BIS=Bispectral index;  
NRS=numeric rating scale; CAM-ICU=confusion assessment method-intensive care  
unit

## Declarations

**Ethics approval and consent to participate:** The study protocol was approved by Clinical Research Ethics Committee of Peking University First Hospital (2015-987). The trial was registered at Chinese Clinical Trial Registry (www.chictr.org.cn) with identifier ChiCTR-IPR-15007654 on December 1, 2015). Written informed consent will be obtained from all patients or their surrogate in law.

**Dissemination:** Results of the study will be presented at academic conferences and submitted to peer-reviewed journals.

**Competing interests:** DXW reports lecture fees and travel expenses for lectures given at academic meetings from Jiangsu Hengrui Medicine Co Ltd, China, and Yichang Humanwell Pharmaceutical Co Ltd, China. DLM is primary investigator of present study which was supported by Beijing Excellent Talent Support Program. Other authors reported no conflict of interests.

**Funding:** This trial was supported by Beijing Excellent Talent Support Program (No. 2014000020124G025). The program committee had no role in study design, data collection, management, analysis, interpretation of data, writing of the report and any other ultimate authority over any of these activities.

**Authors' contributions:** DXW and DLM designed this study. DLM draft the manuscript of protocol. DXW critically revised the manuscript. DLM, BJW, CJL, JH,

1 HJL, CG, ZHW and QCZ participate in the conduct of the study. All authors read and  
2 approved the final manuscript.

3  
4  
5  
6  
7  
8  
9  
10 **Acknowledgements:** We thank professor Xin-Yu Sun (Department of Psychiatrics,  
11 Peking University Sixth Hospital, Beijing, China) for her help in psychiatric  
12 consultation and personnel training. We also thank Dr. Xue-Ying Li (Department of  
13 Medical Statistic, Peking University First Hospital, Beijing, China) for her help in  
14 statistical analysis of this research.

15  
16  
17  
18  
19  
20  
21  
22 **Data sharing:** No plan  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) Washington, DC: American Psychiatric Association; 1994.
2. Inouye SK, Westendorp RG, Saczynski JS. Delirium in elderly people. *Lancet*. 2014 Mar 8;383(9920):911-22.
3. Su X, Meng ZT, Wu XH, et al. Dexmedetomidine for prevention of delirium in elderly patients after non-cardiac surgery: a randomised, double-blind, placebo-controlled trial. *Lancet*. 2016 Oct 15; 388(10054): 1893-1902.
4. Ely EW, Shintani A, Truman B, et al. Delirium as a predictor of mortality in mechanically ventilated patients in the intensive care unit. *JAMA*. 2004 Apr 14; 291(14): 1753-62.
5. Saczynski JS, Marcantonio ER, Quach L, et al. Cognitive trajectories after postoperative delirium. *N Engl J Med*. 2012 Jul 5; 367(1): 30-9.
6. Aldecoa C, Bettelli G, Bilotta F, et al. European Society of Anaesthesiology evidence-based and consensus-based guidelines on postoperative delirium. *Eur J Anaesthesiol*. 2017 Apr; 34(4):192-214.
7. Creeley C, Dikranian K, Dissen G, et al. Propofol-induced apoptosis of neurones and oligodendrocytes in fetal and neonatal rhesus macaque brain. *Br J Anaesth*. 2013 Jun; 110 Suppl 1: i29-38.
8. Xiong WX, Zhou GX, Wang B, et al. Impaired spatial learning and memory after sevoflurane-nitrous oxide anesthesia in aged rats is associated with down-regulated cAMP/CREB signaling. *PLoS One*. 2013 Nov 15;8(11): e79408.
9. Chan MT, Cheng BC, Lee TM, et al. BIS-guided anesthesia decreases

- 1  
2  
3 1 postoperative delirium and cognitive decline. *J Neurosurg Anesthesiol.* 2013 Jan;  
4  
5 2 25(1): 33-42.  
6  
7 3 10. Vaurio LE, Sands LP, Wang Y, et al. Postoperative delirium: the importance of  
8  
9 4 pain and pain management. *Anesth Analg.* 2006 Apr; 102(4): 1267-73.  
10  
11 5 11. Lynch EP, Lazor MA, Gellis JE, et al. The impact of postoperative pain on the  
12  
13 6 development of postoperative delirium. *Anesth Analg* 1998; 86:781-5  
14  
15 7 12. Hirsch J, Vacas S, Terrando N, et al. Perioperative cerebrospinal fluid and  
16  
17 8 plasma inflammatory markers after orthopedic surgery. *J Neuroinflammation.*  
18  
19 9 2016 Aug 30; 13(1): 211.  
20  
21 10 13. Cape E, Hall RJ, van Munster BC, et al. Cerebrospinal fluid markers of  
22  
23 11 neuroinflammation in delirium: a role for interleukin-1 $\beta$  in delirium after hip  
24  
25 12 fracture. *J Psychosom Res.* 2014 Sep; 77(3): 219-25.  
26  
27 13 14. van den Boogaard M, Kox M, Quinn KL, et al. Biomarkers associated with  
28  
29 14 delirium in critically ill patients and their relation with long-term subjective  
30  
31 15 cognitive dysfunction; indications for different pathways governing delirium in  
32  
33 16 inflamed and noninflamed patients. *Crit Care.* 2011; 15(6): R297.  
34  
35 17 15. McGrane S, Girard TD, Thompson JL, et al. Procalcitonin and C-reactive protein  
36  
37 18 levels at admission as predictors of duration of acute brain dysfunction in  
38  
39 19 critically ill patients. *Crit Care.* 2011; 15(2): R78.  
40  
41 20 16. Kazmierski J, Banys A, Latek J, et al. Raised IL-2 and TNF- $\alpha$  concentrations are  
42  
43 21 associated with postoperative delirium in patients undergoing coronary-artery  
44  
45 22 bypass graft surgery. *Int Psychogeriatr.* 2014 May; 26(5): 845-55.  
46  
47 23 17. Shin HW, Yoo HN, Kim DH, et al. Preanesthetic dexmedetomidine 1 $\mu$ g/kg  
48  
49 24 single infusion is a simple, easy, and economic adjuvant for general anesthesia.  
50  
51 25 Korean *J Anesthesiol.* 2013 Aug; 65(2): 114-20.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 18. Li J, Xiong M, Nadavaluru PR, et al. Dexmedetomidine attenuates neurotoxicity  
4 induced by prenatal propofol exposure. *J Neurosurg Anesthesiol.* 2016 Jan;  
5 28(1): 51-64.  
6  
7  
8  
9 19. Sanders RD, Xu J, Shu Y, et al. Dexmedetomidine attenuates isoflurane-induced  
10 neurocognitive impairment in neonatal rats. *Anesthesiology.* 2009 May; 110(5):  
11 1077-85.  
12  
13  
14  
15  
16 20. Le Bot A, Michelet D, Hilly J, et al. Efficacy of intraoperative dexmedetomidine  
17 compared with placebo for surgery in adults: a meta-analysis of published  
18 studies. *Minerva Anesthesiol.* 2015 Oct; 81(10): 1105-17.  
19  
20  
21  
22 21. Kang SH, Kim YS, Hong TH, et al. Effects of dexmedetomidine on  
23 inflammatory responses in patients undergoing laparoscopic cholecystectomy.  
24  
25  
26  
27  
28  
29 22. Li Y, Wang B, Zhang LL, et al. Dexmedetomidine Combined with General  
30 Anesthesia Provides Similar Intraoperative Stress Response Reduction When  
31 Compared with a Combined General and Epidural Anesthetic Technique. *Anesth*  
32  
33  
34  
35  
36  
37  
38 23. Cao JL, Pei YP, Wei JQ, et al. Effects of intraoperative dexmedetomidine with  
39 intravenous anesthesia on postoperative emergence agitation/delirium in  
40 pediatric patients undergoing tonsillectomy with or without adenoidectomy: A  
41  
42  
43  
44  
45  
46  
47  
48  
49 24. Sun Y, Liu J, Yuan X, et al. Effects of dexmedetomidine on emergence delirium  
50 in pediatric cardiac surgery. *Minerva Pediatr.* 2017 Jun; 69(3): 165-173.  
51  
52  
53 25. Li X, Yang J, Nie XL, et al. Impact of dexmedetomidine on the incidence of  
54 delirium in elderly patients after cardiac surgery: A randomized controlled trial.  
55  
56  
57  
58  
59  
60

- 1 PLoS One. 2017 Feb 9; 12(2): e0170757.
- 2 26. Yang X, Li Z, Gao C, et al. Effect of dexmedetomidine on preventing agitation  
3 and delirium after microvascular free flap surgery: a randomized, double-blind,  
4 control study. *J Oral Maxillofac Surg.* 2015 Jun; 73(6): 1065-72.
- 5 27. Deiner S, Luo X, Lin HM, et al. Intraoperative Infusion of Dexmedetomidine for  
6 Prevention of Postoperative Delirium and Cognitive Dysfunction in Elderly  
7 Patients Undergoing Major Elective Noncardiac Surgery: A Randomized Clinical  
8 Trial. *JAMA Surg.* 2017 Aug 16; 152(8): e171505.
- 9 28. Collin C, Wade DT, Davies S, et al. The Barthel ADL Index: a reliability study.  
10 *Int Disabil Stud.* 1988; 10: 61-3.
- 11 29. Katzman R, Zhang MY, Ouang-Ya-Qu, et al. A Chinese version of the Mini-  
12 Mental State Examination; impact of illiteracy in a Shanghai dementia survey. *J*  
13 *Clin Epidemiol.* 1988; 41: 971-8.
- 14 30. Inouye SK, van Dyck CH, Alessi CA, et al. Clarifying confusion: The Confusion  
15 Assessment Method. A new method for detecting delirium. *Ann Intern Med*  
16 1990; 113:941-8
- 17 31. Mu DL, Zhang DZ, Wang DX, et al. Parecoxib Supplementation to Morphine  
18 Analgesia Decreases Incidence of Delirium in Elderly Patients After Hip or Knee  
19 Replacement Surgery: A Randomized Controlled Trial. *Anesth Analg.* 2017 Jun;  
20 124(6):1992-2000.
- 21 32. Ely EW, Margolin R, Francis J, et al. Evaluation of delirium in critically ill  
22 patients: validation of the Confusion Assessment Method for the Intensive Care  
23 Unit (CAM-ICU). *Crit Care Med.* 2001 Jul; 29(7): 1370-9.
- 24 33. Ji L, Vc L, Leung CM, et al. Clinical utility and validation of two instruments  
25 (the Confusion Assessment Method Algorithm and the Chinese version of

1  
2  
3 1 Nursing Delirium Screening Scale) to detect delirium in geriatric inpatients. Gen  
4  
5 2 Hosp Psychiatry. 2008; 30: 171-6.  
6  
7 34. Wang C, Wu Y, Yue P, et al. Delirium assessment using Confusion Assessment  
8  
9 4 Method for the Intensive Care Unit in Chinese critically ill patients. J Crit Care.  
10  
11 5 2013; 28: 223-9.  
12  
13  
14 6  
15  
16 7  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

**Figure legend**

1

2

3 Figure 1. Flowchart of this study.

For peer review only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 1 Flow chart of this study




Figure1 Flow chart of this study

209x297mm (300 x 300 DPI)



SPIRIT 2013 Checklist: Recommended items to address in a clinical trial protocol and related documents\*

Section/item	Item No	Description	Addressed on page number
<b>Administrative information</b>			
Title	1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	1
Trial registration	2a	Trial identifier and registry name. If not yet registered, name of intended registry	4, 8, 19
	2b	All items from the World Health Organization Trial Registration Data Set	8
Protocol version	3	Date and version identifier	8
Funding	4	Sources and types of financial, material, and other support	19
Roles and responsibilities	5a	Names, affiliations, and roles of protocol contributors	19
	5b	Name and contact information for the trial sponsor	19
	5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities	19
	5d	Composition, roles, and responsibilities of the coordinating centre, steering committee, endpoint adjudication committee, data management team, and other individuals or groups overseeing the trial, if applicable (see Item 21a for data monitoring committee)	N/A

**Introduction**



1				
2				
3	Background and	6a	Description of research question and justification for undertaking the trial, including summary of relevant	6-7
4	rationale		studies (published and unpublished) examining benefits and harms for each intervention	
5				
6		6b	Explanation for choice of comparators	6-7, 10
7	Objectives	7	Specific objectives or hypotheses	7
8				
9	Trial design	8	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group),	
10			allocation ratio, and framework (eg, superiority, equivalence, noninferiority, exploratory)	8
11				
12				
13	<b>Methods: Participants, interventions, and outcomes</b>			
14				
15	Study setting	9	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will	8
16			be collected. Reference to where list of study sites can be obtained	
17				
18	Eligibility criteria	10	Inclusion and exclusion criteria for participants. If applicable, eligibility criteria for study centres and	8-9
19			individuals who will perform the interventions (eg, surgeons, psychotherapists)	
20				
21	Interventions	11a	Interventions for each group with sufficient detail to allow replication, including how and when they will be	10
22			administered	
23				
24		11b	Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose	10-11
25			change in response to harms, participant request, or improving/worsening disease)	
26				
27		11c	Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence	10-11
28			(eg, drug tablet return, laboratory tests)	
29				
30		11d	Relevant concomitant care and interventions that are permitted or prohibited during the trial	N/A
31				
32	Outcomes	12	Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood	12-15
33			pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg,	
34			median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen	
35			efficacy and harm outcomes is strongly recommended	
36				
37	Participant timeline	13	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for	Figure 1
38			participants. A schematic diagram is highly recommended (see Figure)	
39				
40				
41				
42				
43				
44				
45				
46				
47				

1				
2				
3	Sample size	14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	14-15
4				
5	Recruitment	15	Strategies for achieving adequate participant enrolment to reach target sample size	9
6				
7				

### 8 **Methods: Assignment of interventions (for controlled trials)**

#### 9 Allocation:

10				
11				
12	Sequence generation	16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	9
13				
14				
15				
16				
17	Allocation concealment mechanism	16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	9-10
18				
19				
20				
21	Implementation	16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	9-10
22				
23				
24	Blinding (masking)	17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	9-10, 12
25				
26				
27		17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	9-10
28				
29				
30				

### 31 **Methods: Data collection, management, and analysis**

32				
33	Data collection methods	18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known. Reference to where data collection forms can be found, if not in the protocol	12-15
34				
35				
36				
37				
38		18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols	12-15
39				
40				
41				
42				
43				
44				
45				
46				
47				

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

Data management	19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	14
Statistical methods	20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	14-15
	20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	14-15
	20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any statistical methods to handle missing data (eg, multiple imputation)	14-15

**Methods: Monitoring**

Data monitoring	21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent from the sponsor and competing interests; and reference to where further details about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not needed	14
	21b	Description of any interim analyses and stopping guidelines, including who will have access to these interim results and make the final decision to terminate the trial	14
Harms	22	Plans for collecting, assessing, reporting, and managing solicited and spontaneously reported adverse events and other unintended effects of trial interventions or trial conduct	13-14
Auditing	23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent from investigators and the sponsor	13-14

**Ethics and dissemination**

Research ethics approval	24	Plans for seeking research ethics committee/institutional review board (REC/IRB) approval	4, 8, 19
Protocol amendments	25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes, analyses) to relevant parties (eg, investigators, REC/IRBs, trial participants, trial registries, journals, regulators)	8



1				
2				
3	Consent or assent	26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	8
4				
5				
6		26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	N/A
7				
8				
9	Confidentiality	27	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	14
10				
11				
12	Declaration of interests	28	Financial and other competing interests for principal investigators for the overall trial and each study site	19
13				
14				
15	Access to data	29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	14
16				
17				
18	Ancillary and post-trial care	30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	14
19				
20				
21	Dissemination policy	31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	19
22				
23				
24				
25		31b	Authorship eligibility guidelines and any intended use of professional writers	19
26				
27		31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	20
28				
29	<b>Appendices</b>			
30				
31	Informed consent materials	32	Model consent form and other related documentation given to participants and authorised surrogates	N/A
32				
33				
34	Biological specimens	33	Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies, if applicable	N/A
35				
36				

37 \*It is strongly recommended that this checklist be read in conjunction with the SPIRIT 2013 Explanation & Elaboration for important clarification on the items.  
 38 Amendments to the protocol should be tracked and dated. The SPIRIT checklist is copyrighted by the SPIRIT Group under the Creative Commons  
 39 "[Attribution-NonCommercial-NoDerivs 3.0 Unported](https://creativecommons.org/licenses/by-nc-nd/3.0/)" license.  
 40

# BMJ Open

## Impact of dexmedetomidine infusion during general anesthesia on incidence of postoperative delirium in elderly patients after major non-cardiac surgery: study protocol of a randomized, double-blinded and placebo-controlled trial

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-019549.R2
Article Type:	Protocol
Date Submitted by the Author:	05-Feb-2018
Complete List of Authors:	Wang, Bojie; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Li, Chunjing; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Hu, Jian; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Li, Huaijin; Peking University First Hospital, Department of Anesthesiology and Critical Care Guo, Chao; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Wang, Zhenhan; Dongping People's Hospital, Department of anesthesiology Zhang, Qiaochu; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Mu, Dongliang; Department of Anesthesiology and Critical Care Medicine Wang, Dong-Xin; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine
<b>Primary Subject Heading</b>:	Anaesthesia
Secondary Subject Heading:	Anaesthesia
Keywords:	Elderly, Major non-cardiac surgery, Intraoperative dexmedetomidine, Postoperative delirium

SCHOLARONE™  
Manuscripts

1  
2  
3 **1 Impact of dexmedetomidine infusion during general anaesthesia on incidence of**  
4  
5 **2 postoperative delirium in elderly after major non-cardiac surgery: study**  
6  
7 **3 protocol of a randomized, double-blinded and placebo-controlled trial**  
8  
9  
10

11 5 Bo-Jie Wang<sup>1</sup>, Chun-Jing Li<sup>1</sup>, Jian Hu<sup>1</sup>, Huai-Jin Li<sup>1</sup>, Chao Guo<sup>1</sup>, Zhen-Han Wang<sup>2</sup>,  
12  
13 6 Qiao-Chu Zhang<sup>1</sup>, Dong-Liang Mu<sup>1,\*</sup>, Dong-Xin Wang<sup>1</sup>  
14  
15

16 7  
17  
18 8 <sup>1</sup>Department of Anesthesiology and Critical Care Medicine, Peking University First  
19  
20 9 Hospital, Beijing, China

21  
22 10 <sup>2</sup>Department of Anesthesiology, Dongping People's Hospital, Shandong, China  
23  
24

25  
26  
27 12 The first two authors (BJ Wang and CJ Li) contributed equally to this manuscript  
28  
29

30  
31 14 Corresponding author: Dong-Liang Mu, M.D., Department of Anesthesiology and  
32  
33 15 Critical Care Medicine, Peking University First Hospital, Beijing, China. Tel: +86  
34  
35 16 010 83572784 (O), +86 13810702725 (M); E-mail: mudongliang@icloud.com  
36  
37

38  
39 18 Bo-Jie Wang: ivyxbd@hotmail.com  
40

41  
42 19 Chun-Jing Li: gogolion@163.com  
43

44  
45 20 Jian Hu: hj19871204@163.com  
46

47  
48 21 Huai-Jin Li: sophie.lee.coffee@gmail.com  
49

50  
51 22 Chao Guo: guochao1988@yeah.com  
52

53  
54 23 Zhen-Han Wang: 1468432914@qq.com  
55

56  
57 24 Qiao-Chu Zhang: 372193745@qq.com  
58

59  
60 25 Dong-Liang Mu: mudongliang@icloud.com

1  
2  
3 1 Dong-Xin Wang: wangdongxin@hotmail.com  
4

5 2  
6

7 3 Word Count: 2845  
8

9 4  
10

11 5  
12

13 6  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

## Abstract for protocol

**Introduction:** Delirium is a common complication in elderly after surgery and is associated with worse outcomes. Multiple risk factors are related with postoperative delirium, such as exposure to general anaesthetics, pain and postoperative inflammatory response. Preclinical and clinical studies have shown that dexmedetomidine attenuated neurotoxicity induced by general anaesthetics, improved postoperative analgesia and inhibited inflammatory response after surgery. Several studies found that intraoperative use of dexmedetomidine can prevent postoperative delirium, but data were inconsistent. This study was designed to investigate the impact of dexmedetomidine administered during general anaesthesia in preventing delirium in elderly after major non-cardiac surgery.

**Method and analysis:** This is a randomized, double-blinded, and placebo-controlled trial. 620 elderly patients (age  $\geq 60$  years) who are scheduled to undertake elective major non-cardiac surgery (with an expected duration  $\geq 2$  hours) are randomly divided into two groups. For patients in the dexmedetomidine group, a loading dose dexmedetomidine (0.6  $\mu\text{g}/\text{kg}$ ) will be administered in 10 minutes before anaesthesia induction, followed by a continuous infusion at a rate of 0.5  $\mu\text{g}/\text{kg}/\text{h}$  till 1 hour before the end of surgery. For patients in the control group, normal saline will be administered with an identical rate as in the dexmedetomidine group. The primary endpoint is the incidence of delirium during the first five postoperative days. Secondary endpoints include pain intensity, cumulative opioid consumption and subjective sleep quality during the first 3 postoperative days as well as the incidence of non-delirium complications and all-cause mortality within 30 days after surgery.



1  
2  
3 1 **Ethics and dissemination:** The study protocol was approved by Clinical Research  
4  
5 2 Ethics Committee of Peking University First Hospital (2015-987) and registered at  
6  
7 3 Chinese Clinical Trial Registry (<http://www.chictr.org.cn>) with identifier ChiCTR-  
8  
9 4 IPR-15007654. The results of the study will be presented at academic conferences  
10  
11 5 and submitted to peer-reviewed journals.  
12  
13  
14 6

15  
16 7 **Key words:** Elderly; Major non-cardiac surgery; Intraoperative dexmedetomidine;  
17  
18 8 Postoperative delirium  
19  
20 9

## Strengths and limitations

### Strengths

1. The study design will be a randomized, double-blinded and placebo-controlled, with a relative large sample size.
2. Anaesthesia depth (Bispectral Index) will be monitored to guide anaesthesia maintenance to ensure patients in both groups received equal depth anaesthesia.

### Limitations

1. This is a single center trial, which will limit the generalizability of our results.
2. Only early outcomes (up to 30 days after surgery) are assessed in this trial.

## Introduction

Delirium is a transient brain dysfunction which is characterized by altered consciousness, inattention, and changes in cognition or perception; it develops acutely with clinical manifestations to be fluctuated with aggressive and depressive manner during developing course.<sup>1</sup> Prevalence of delirium varies from 12% to 51% in patients after non-cardiac surgery, and it is increased with age.<sup>2-3</sup> The occurrence of postoperative delirium (POD) is associated with worsen outcomes including prolonged mechanical ventilation and intensive care unit stay, increased postoperative complications, high mortality rate and long-term cognitive decline.<sup>2-5</sup>

The aetiology of POD is multifactorial and includes several intraoperative factors.<sup>6</sup> For example, exposure to general anaesthetics (e.g., propofol or sevoflurane) might produce neurotoxicity,<sup>7-8</sup> whereas reduced anaesthetic consumption by avoiding deep anaesthesia reduced the occurrence of delirium in elderly patients undergoing major non-cardiac surgery.<sup>9</sup> Poor pain management is another risk factor of POD.<sup>6,10</sup> It was reported that the risk of postoperative delirium was 1.2 times higher for every unit increment in visual analogue pain score (an 11-point pain scale where 0 indicates no pain and 10 the most severe pain).<sup>11</sup> Inflammation is also proposed to play an important role in the pathogenesis of POD.<sup>2</sup> Inflammatory responses induced by surgery and anaesthesia are manifested by elevated levels of interleukins (IL), C-reactive protein (CRP) and tumor necrosis factor (TNF).<sup>2,12</sup> Studies by our group and others found that higher levels of inflammatory mediators are associated with increased risk of POD.<sup>2,13-16.</sup>

Dexmedetomidine is a highly selective  $\alpha_2$ -receptor agonist with sedative, analgesic and anxiolytic effects.<sup>17-21</sup> When used as a supplement during intraoperative

1 anaesthesia, it reduces the consumption of general anaesthetics.<sup>17</sup> Preclinical evidence  
2 suggested that use of dexmedetomidine might attenuate neurotoxicity induced by  
3 general anaesthetics.<sup>18-19</sup> In a meta-analysis, intraoperative administration of  
4 dexmedetomidine lowers postoperative pain intensity and reduces opioid  
5 consumption.<sup>20</sup> Clinical evidence also showed that intraoperative dexmedetomidine  
6 significantly inhibits hyper-secretion of inflammatory cytokines during and after  
7 surgery.<sup>21-22</sup>

8       Use of dexmedetomidine during general anaesthesia may reduce POD. In  
9 pediatric patients undergoing tonsillectomy and cardiac surgery, intraoperative  
10 infusion of dexmedetomidine lowered the incidence of emergence delirium.<sup>23,24</sup> In  
11 adult patients undergoing cardiac surgery and microvascular free flap surgery,  
12 intraoperative dexmedetomidine (comparison with normal saline) slightly decreased  
13 the incidence of delirium, although the differences were not statistically significant  
14 between two groups possibly due to underpowered sample size.<sup>25,26</sup> In a recent study  
15 of Deiner et al.,<sup>27</sup> use of dexmedetomidine during general anaesthesia did not reduce  
16 delirium after major non-cardiac surgery in the elderly. However, in that study,  
17 anaesthesia depth was not monitored and the consumption of anaesthetics (such as  
18 propofol and fentanyl) was similar between the two groups. It was possible that  
19 patients in the dexmedetomidine group had deeper anaesthesia which might have  
20 increased the risk of delirium.<sup>27</sup> Therefore, the effects of dexmedetomidine  
21 administered during general anaesthesia on the occurrence of POD need to be  
22 evaluated further.

23       This study is designed to investigate whether dexmedetomidine use during  
24 general anaesthesia can decrease the incidence of POD in elderly after major non-  
25 cardiac surgery.

## Method and analysis

### Study design

This randomized, double-blinded, and placebo-controlled trial with two parallel arms was designed to test the superiority of dexmedetomidine administered during general anaesthesia on the incidence of delirium after surgery. Patients will be randomized into either the dexmedetomidine group or the control group (Figure 1). The study is conducted in the Department of Anesthesiology and Critical Care Medicine of Peking University First Hospital.

### Ethics approval

The study protocol (version 1.1, issue date Nov, 2015) was approved by Clinical Research Ethics Committee of Peking University First Hospital (2015-987). Any protocol modifications will be submitted for review and approval by the Ethics Committee. The trial was registered at Chinese Clinical Trial Registry (<http://www.chictr.org.cn>) on December 1, 2015, with identifier ChiCTR-IPR-15007654. Written informed consents are obtained from every patients or their surrogate in law.

### Participants

Elderly patients (age  $\geq 60$  years) who are scheduled to undergo elective non-cardiac surgery with expected duration  $\geq 2$  hours under general anaesthesia are screened for inclusion. Those who meet any of the following criteria will be excluded: (1) do not provide written informed consents; (2) previous history of schizophrenia, epilepsy or Parkinson disease; (3) visual, hearing, language or other barrier which impede communication and preoperative delirium assessment; (4) neurosurgery or traumatic brain injury; (5) severe bradycardia (heart rate less than 40 beats per minute), sick

1 sinus syndrome or atrioventricular block of degree 2 or above; (6) severe hepatic  
2 dysfunction (Child-Pugh grade C); or (7) renal failure (requirement of renal  
3 replacement therapy).

#### 4 **Patient recruitment and baseline data collection**

5 Potential participants are screened by investigators the day before surgery (or on  
6 Friday for those who will undergo surgery next Monday). The study protocol  
7 including potential risks and benefits will be explained to patients in person. Those  
8 who do not meet the exclusion criteria are invited to participate in the study.

9 After obtaining written informed consents, the following baseline data are  
10 collected: demographic data, preoperative diagnosis, comorbidity, current medical  
11 therapy, previous surgery, and main results of physical and laboratory examinations.  
12 Barthel index is used for evaluation of activities of daily living.<sup>28</sup> Cognitive function  
13 is assessed with Mini-Mental State Examination (MMSE).<sup>29</sup> Preoperative delirium is  
14 assessed with confusion assessment method (CAM).<sup>30</sup>

#### 15 **Randomization, grouping and blinding**

16 Random numbers were created by an independent statistician using SAS statistical  
17 package version 9.3 (SAS Institute, Cary, NC, USA) in a 1:1 ratio and were sealed in  
18 envelopes. A study coordinator, who has no knowledge of patients before  
19 randomization and does not participate in anaesthesia and postoperative follow-up of  
20 enrolled patients, will open envelop for random numbers and prepare study drugs  
21 before induction of anaesthesia.

22 Information regarding randomization, study drug preparation and group  
23 allocation will be masked from investigators who perform data collection and  
24 postoperative follow-up, anaesthesiologists, patients and other healthcare team  
25 members. Blinding will be maintained throughout the entire study period.

1 To ensure patients' safety, study group allocation can be unmasked in the  
2 following conditions, i.e., occurrence of severe adverse events or any unexpected  
3 deterioration of patient's clinical status. These situations will be documented in the  
4 Case Report Forms (CRFs). The unmasked patients will be included in the intention-  
5 to-treat population but excluded from per-protocol analysis.

#### 6 **Interventions, anaesthesia and analgesia**

7 Intraoperative monitoring includes electrocardiogram, noninvasive blood pressure,  
8 pulse oxygen saturation, end-tidal carbon dioxide, nasopharyngeal temperature, urine  
9 output, and bispectral index (BIS). Intraarterial pressure (including derivative  
10 dynamic parameters such as stroke volume variation by FlowTrac system) and central  
11 venous pressure are monitored according to patients' conditions.

12 Study drugs, either 200 µg (2 ml) dexmedetomidine (Jiangsu Hengrui Medicine  
13 Co, Ltd, Jiangsu, China) or 2 ml normal saline, are diluted into 50 ml normal saline.  
14 All study drugs are colorless solution provided in syringes of the same size and brand.  
15 The regimen of study drug administration includes a loading dose of 0.15 ml/kg (i.e.,  
16 0.6 µg/kg dexmedetomidine for patients in the dexmedetomidine group) administered  
17 during a 10-minute period before anaesthesia induction. This is followed by a  
18 continuous infusion at a rate of 0.125 ml/kg/h (i.e., a rate of 0.5 µg/kg/h  
19 dexmedetomidine for patients in the dexmedetomidine group) till 1 hour before the  
20 end of surgery. Study drug infusion is performed by an infusion pump specially  
21 designed for dexmedetomidine administration (Slgo<sup>®</sup> CP1000, Beijing Slgo medical  
22 technology Co., Ltd.).

23 Attending anaesthesiologists can decrease or stop study drug infusion in the  
24 following conditions: (1) severe bradycardia or hypotension which does not improve  
25 after routine treatment; (2) new onset atrioventricular block which does not improve

1 after routine treatment; or (3) other conditions that anaesthesiologists consider it  
2 necessary. In these conditions, the reasons that lead to any protocol deviations will be  
3 recorded in the case report forms (CRF). These patients will be included in the  
4 intention-to-treat analysis but excluded from the per-protocol analysis.

5 Anaesthesia is induced with intravenous sufentanil (target controlled infusion  
6 with effect-site concentration from 0.2 to 0.5 ng/ml) and propofol (2-3 mg/kg) and  
7 maintained with intravenous sufentanil (effect-site concentration from 0.2 to 0.5  
8 ng/ml) and propofol (4-12 mg/kg/h) and inhalation of a 1:1 nitrous oxide-oxygen  
9 mixture. Rocuronium and/or cisatracurium are administered for muscle relaxation.  
10 Patients will be mechanically ventilated with a tidal volume of 6-8 ml/kg and a  
11 positive end-expiratory pressure of 5 cm H<sub>2</sub>O. The mean arterial pressure is  
12 maintained above 60 mmHg or within 20% from baseline. BIS is maintained between  
13 40 and 60. Body temperature is maintained with air-warming and fluid heating  
14 systems. The target of nasopharyngeal temperature maintenance during surgery is  
15 from 36.0 to 37°C.

16 All patients are transferred to post-anaesthesia care unit (PACU) or intensive  
17 care unit (ICU) before they are sent back to general wards. Patient-controlled  
18 intravenous analgesia (PCIA) is provided for all patients, which is established with  
19 0.5 mg/ml morphine in 100 ml normal saline and programmed to deliver a 1 mg bolus  
20 with a lock-out interval of 8 minutes and a background infusion at 0.5 mg/h.  
21 Supplemental morphine at dose of 2 to 4 mg will be administered at 10-minute  
22 intervals if the numeric rating scale (NRS) pain score (a 11-point scale where 0  
23 indicates no pain and 10 indicates the worst pain) remains above 4 after 3 consecutive  
24 PCIA boluses.<sup>31</sup> Other postoperative managements were performed according to  
25 routine practice.



## 1 **Outcome assessment**

2 Patients are followed up twice daily during the first 5 postoperative days and then  
3 weekly until 30 days after surgery. Investigators who are responsible for  
4 postoperative follow-up are not involved in anaesthesia and perioperative care, and  
5 are not allowed to exchange patients' information with anaesthesiologists who take  
6 care of patients in the operating room. Before the beginning of the study, investigators  
7 are trained to follow the study protocol and to perform delirium assessment and the  
8 training process is repeated at 4 to 6 months intervals during the study period.<sup>2-3, 31</sup>

9 The 4-hour training courses of delirium assessment include the following contents: (1)  
10 lectures regarding signs/symptoms, diagnosis and treatment of delirium by  
11 psychiatrists; (2) training courses of the use of CAM and CAM-ICU on patient-actors  
12 (trained ICU physicians or nurses who act as patients with or without delirium)  
13 conducted by psychiatrists. The process continued until 100% agreement is achieved  
14 in diagnosing delirium.

### 15 **Primary endpoint**

16 The primary endpoint is the incidence of delirium during the first 5 days after surgery.  
17 Delirium is assessed twice daily (at 08:00-09:00 and 19:00-20:00, respectively) with  
18 CAM for non-intubated patients<sup>30</sup> or CAM for the Intensive Care Unit (CAM-ICU)  
19 for intubated patients.<sup>32</sup> These delirium assessment methods had been used in our  
20 previous studies.<sup>2-3, 31</sup> For patients who are discharged or died within 5 days after  
21 surgery, the results of last delirium assessment will be considered the results of the  
22 missing data. These patients will be excluded when calculating daily prevalence of  
23 delirium in a post-hoc analysis.

### 24 **Secondary endpoints**

1 Postoperative pain intensities at rest and with movement are assessed with NRS pain  
2 score at 24, 48 and 72 hours after surgery, respectively.<sup>31</sup> Cumulative morphine  
3 consumptions at these time points are recorded. Subjective sleep quality is assessed  
4 with NRS (an 11-point scale where 0 indicates the worst possible sleep and 10 the  
5 best possible sleep) at 08:00 on the first, second and third morning after surgery.<sup>3,31</sup>  
6 Other secondary endpoints include non-delirium complications within 30 days after  
7 surgery, length of stay in hospital after surgery and all-cause 30-day mortality. Non-  
8 delirium complications are generally defined as new-onset non-delirium conditions  
9 after surgery that are harmful to patients' recovery and require therapeutic  
10 intervention.

### 11 **Safety outcomes**

12 In the present study, adverse events are monitored from the start of study drug  
13 administration until PACU discharge or 2 hours after ICU admission. Hypotension is  
14 defined as systolic blood pressure of less than 90 mmHg or a decrement of more than  
15 30% from baseline. Hypertension is defined as systolic blood pressure of more than  
16 180 mmHg or an increment of more than 30% from baseline. Bradycardia is defined  
17 as heart rate of less than 40 beats per minute. Tachycardia is defined as heart rate of  
18 more than 100 beats per minute. Desaturation is defined as SpO<sub>2</sub> of less than 90%.  
19 Emergence agitation is defined as a Richmond Agitation-Sedation Scale (RASS)  
20 score of more than +2 within 30 minutes after extubation. Delayed extubation is  
21 defined when time to extubation is more than 2 hours (from the end of surgery) in  
22 PACU patients or more than 4 hours in ICU patients.<sup>2, 22-25</sup>

23 Severe adverse events, i.e., those that might result in patient's  
24 disability/deformity, prolong in-hospital stay, or life threatening events, will be  
25 reported to Clinical Research Ethics Committee of Peking University First Hospital

1 within 24 hours. For patients who suffered harm from present trial, medical treatment  
2 will be initiated as soon as possible and compensation will be completed according to  
3 local laws and regulations.

#### 4 **Data monitoring and management**

5 Original data will be recorded in the CRFs accordingly. All data will be kept  
6 confidentially. The completed CRFs will be checked by a study coordinator who is  
7 qualified by the principal investigator. Supplementations and corrections will be made  
8 when necessary. Data entry will be performed in a double-input and double-check  
9 way with the Data Management System (Fantastic Eight Tech. Co., Ltd, Beijing,  
10 China) of the Peking University First Hospital.

11 The conduct of the study and the quality of data will be monitored by the  
12 Clinical Research Ethics Committee of Peking University First Hospital. Data  
13 management and statistical analysis will be performed by the Department of  
14 Biostatistics of Peking University First Hospital. Considering that dexmedetomidine  
15 has been widely used during general anaesthesia and its safety has been confirmed, no  
16 interim analysis will be performed and the trial will continue until the target sample  
17 size is achieved.

#### 18 **Statistical analysis**

##### 19 ***Sample size calculation***

20 In our previous study, the incidence of delirium was 14.8% in elderly patients after  
21 non-cardiac surgery.<sup>33</sup> Previous studies reported that intraoperative dexmedetomidine  
22 decreased the incidence of POD by 60-77% in comparison with placebo.<sup>22, 25</sup> We  
23 assumed that the incidence of POD will be reduced from 14.8% to 7.4% (i.e., a 50%  
24 reduction) in the present study. With the power set at 80% and significant level at

1 0.05, 564 patients are required to detect the difference. Considering a loss to follow-  
2 up rate of about 9%, we plan to enroll 620 patients in this study.

### 3 ***Outcome analysis***

4 Continuous data with normal distribution will be compared using independent sample  
5 T-test. Continuous data with asymmetric distribution will be compared using  
6 independent sample Mann-Whitney U test. Categorical data will be compared using  
7 Chi-squared test or continuity correction Chi-squared test. The difference (and 95%  
8 confidence interval of the difference) between two means or medians will be  
9 estimated using the methodology of Levene's test or Hodges-Lehmann estimator.  
10 Time-to-event data will be analyzed by survival analysis with differences between  
11 groups compared with log-rank test.

12 Statistical analyses will be performed with the SPSS 14.0 (SPSS, Inc., Chicago,  
13 IL) and SAS 9.3 (SAS Institute, Cary, NC, USA). All tests are two tailed, and P  
14 values of less than 0.05 are considered to be statistically significant. The Bonferroni  
15 adjustment is made to control type I error for multiple testing.

16

## Discussion

This randomized, double-blinded, and placebo-controlled single center trial was designed to investigate if dexmedetomidine administration during general anaesthesia can decrease the incidence of POD in elderly patients after major non-cardiac surgery.

In the present study, the dosing regimen of dexmedetomidine is similar to our previous study because it does not increase drug-related adverse events (such as severe bradycardia and hypotension).<sup>25</sup> Furthermore, the CAM and CAM-ICU are used to assess delirium in patients with or without intubation, respectively.<sup>30, 32</sup> Both CAM and CAM-ICU have been validated in Chinese population.<sup>34, 35</sup> Feasibility of these two assessment tools have been confirmed in our previous studies.<sup>2-3, 31</sup> To maintain the quality of delirium assessment, investigators in charge of postoperative follow-up are trained by a psychiatrist before the study and will be retrained at 4 to 6-month intervals.

Because of the hemodynamic and anaesthetic-sparing effect of dexmedetomidine, it is not very difficult for the experienced anaesthesiologists to guess which study drug is administrated. This might weak the blinding to anaesthesiologists. However, in the present study, investigators who are responsible for postoperative follow-up and delirium assessment are not involved in anaesthesia and perioperative care; and they are not allowed to exchange patients' information with anaesthesiologists who take care of patients in the operating room. In this way the blinding of investigators to study group assignment can be guaranteed.

The strengths of the present study include the following when compared with previous studies.<sup>22-24</sup> Firstly, a randomized, double-blind, and placebo-controlled study design with a relative large sample size (620 patients) is adopted. Results of the

1 study will provide high quality evidences. Secondly, BIS level is monitored in all  
2 enrolled patients, which will help us to avoid unnecessary and potentially harmful  
3 deep anaesthesia. Thirdly, safety data will be recorded in detail. Our study also has  
4 some limitations. One is that this is a single center trial, which will limit the external  
5 validity of our results. Another one is that only early outcomes (up to 30 days after  
6 surgery) will be explored.

7  
8 **Trail status:**

9 This study is currently at stage of patient enrollment and data collection. The current  
10 version of study protocol is V1.1 and is approved on November 27, 2015. Patient  
11 recruitment started from December 2, 2015 and is expected to be finished by March  
12 31, 2018.

13

**List of abbreviations**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8

1  
2  
3  
4  
5  
6  
7  
8

POD=postoperative delirium; IL=interleukins; CRP=C-reactive protein; TNF=tumor  
necrosis factor; IRB=Institutional Review Board; MMSE=Mini-Mental State  
Examination; CAM=confusion assessment method (CAM); BIS=Bispectral index;  
NRS=numeric rating scale; CAM-ICU=confusion assessment method-intensive care  
unit

## Declarations

**Ethics approval and consent to participate:** The study protocol was approved by Clinical Research Ethics Committee of Peking University First Hospital (2015-987). The trial was registered at Chinese Clinical Trial Registry (www.chictr.org.cn) with identifier ChiCTR-IPR-15007654 on December 1, 2015). Written informed consent will be obtained from all patients or their surrogate in law.

**Dissemination:** Results of the study will be presented at academic conferences and submitted to peer-reviewed journals.

**Competing interests:** DXW reports lecture fees and travel expenses for lectures given at academic meetings from Jiangsu Hengrui Medicine Co Ltd, China, and Yichang Humanwell Pharmaceutical Co Ltd, China. DLM is primary investigator of present study which was supported by Beijing Excellent Talent Support Program. Other authors reported no conflict of interests.

**Funding:** This trial was supported by Beijing Excellent Talent Support Program (No. 2014000020124G025). The sponsors have no role in the study design and conduct; the collection, management, analysis, and interpretation of the data; or the preparation and approval of the manuscript.

**Authors' contributions:** DXW and DLM designed this study. DLM draft the manuscript of protocol. DXW critically revised the manuscript. DLM, BJW, CJL, JH,



1 HJL, CG, ZHW and QCZ participate in the conduct of the study. All authors read and  
2 approved the final manuscript.

3  
4  
5  
6  
7  
8  
9 **Acknowledgements:** We thank professor Xin-Yu Sun (Department of Psychiatrics,  
10 Peking University Sixth Hospital, Beijing, China) for her help in psychiatric  
11 consultation and personnel training and Dr. Xue-Ying Li (Department of Biostatistics,  
12 Peking University First Hospital, Beijing, China) for her help in statistical  
13 consultation. We also thank Professor Daqing Ma (Section of Anesthetics, Pain  
14 Medicine and Intensive Care, Department of Surgery and Cancer, Imperial College  
15 London, London, UK) for his help in revising the manuscript.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27 **Data sharing:** Will be provided on request.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) Washington, DC: American Psychiatric Association; 1994.
2. Inouye SK, Westendorp RG, Saczynski JS. Delirium in elderly people. *Lancet*. 2014 Mar 8;383(9920):911-22.
3. Su X, Meng ZT, Wu XH, et al. Dexmedetomidine for prevention of delirium in elderly patients after non-cardiac surgery: a randomised, double-blind, placebo-controlled trial. *Lancet*. 2016 Oct 15; 388(10054): 1893-1902.
4. Ely EW, Shintani A, Truman B, et al. Delirium as a predictor of mortality in mechanically ventilated patients in the intensive care unit. *JAMA*. 2004 Apr 14; 291(14): 1753-62.
5. Saczynski JS, Marcantonio ER, Quach L, et al. Cognitive trajectories after postoperative delirium. *N Engl J Med*. 2012 Jul 5; 367(1): 30-9.
6. Aldecoa C, Bettelli G, Bilotta F, et al. European Society of Anaesthesiology evidence-based and consensus-based guidelines on postoperative delirium. *Eur J Anaesthesiol*. 2017 Apr; 34(4):192-214.
7. Creeley C, Dikranian K, Dissen G, et al. Propofol-induced apoptosis of neurones and oligodendrocytes in fetal and neonatal rhesus macaque brain. *Br J Anaesth*. 2013 Jun; 110 Suppl 1: i29-38.
8. Xiong WX, Zhou GX, Wang B, et al. Impaired spatial learning and memory after sevoflurane-nitrous oxide anesthesia in aged rats is associated with down-regulated cAMP/CREB signaling. *PLoS One*. 2013 Nov 15;8(11): e79408.
9. Chan MT, Cheng BC, Lee TM, et al. BIS-guided anesthesia decreases

- 1  
2  
3 1 postoperative delirium and cognitive decline. *J Neurosurg Anesthesiol.* 2013 Jan;  
4  
5 2 25(1): 33-42.  
6  
7 3 10. Vaurio LE, Sands LP, Wang Y, et al. Postoperative delirium: the importance of  
8  
9 4 pain and pain management. *Anesth Analg.* 2006 Apr; 102(4): 1267-73.  
10  
11 5 11. Lynch EP, Lazor MA, Gellis JE, et al. The impact of postoperative pain on the  
12  
13 6 development of postoperative delirium. *Anesth Analg* 1998; 86:781-5  
14  
15 7 12. Hirsch J, Vacas S, Terrando N, et al. Perioperative cerebrospinal fluid and  
16  
17 8 plasma inflammatory markers after orthopedic surgery. *J Neuroinflammation.*  
18  
19 9 2016 Aug 30; 13(1): 211.  
20  
21 10 13. Cape E, Hall RJ, van Munster BC, et al. Cerebrospinal fluid markers of  
22  
23 11 neuroinflammation in delirium: a role for interleukin-1 $\beta$  in delirium after hip  
24  
25 12 fracture. *J Psychosom Res.* 2014 Sep; 77(3): 219-25.  
26  
27 13 14. van den Boogaard M, Kox M, Quinn KL, et al. Biomarkers associated with  
28  
29 14 delirium in critically ill patients and their relation with long-term subjective  
30  
31 15 cognitive dysfunction; indications for different pathways governing delirium in  
32  
33 16 inflamed and noninflamed patients. *Crit Care.* 2011; 15(6): R297.  
34  
35 17 15. McGrane S, Girard TD, Thompson JL, et al. Procalcitonin and C-reactive protein  
36  
37 18 levels at admission as predictors of duration of acute brain dysfunction in  
38  
39 19 critically ill patients. *Crit Care.* 2011; 15(2): R78.  
40  
41 20 16. Kazmierski J, Banys A, Latek J, et al. Raised IL-2 and TNF- $\alpha$  concentrations are  
42  
43 21 associated with postoperative delirium in patients undergoing coronary-artery  
44  
45 22 bypass graft surgery. *Int Psychogeriatr.* 2014 May; 26(5): 845-55.  
46  
47 23 17. Shin HW, Yoo HN, Kim DH, et al. Preanesthetic dexmedetomidine 1 $\mu$ g/kg  
48  
49 24 single infusion is a simple, easy, and economic adjuvant for general anesthesia.  
50  
51 25 Korean *J Anaesthesiol.* 2013 Aug; 65(2): 114-20.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1 18. Li J, Xiong M, Nadavaluru PR, et al. Dexmedetomidine attenuates neurotoxicity  
2 induced by prenatal propofol exposure. *J Neurosurg Anesthesiol.* 2016 Jan;  
3 28(1): 51-64.
- 4 19. Sanders RD, Xu J, Shu Y, et al. Dexmedetomidine attenuates isoflurane-induced  
5 neurocognitive impairment in neonatal rats. *Anesthesiology.* 2009 May; 110(5):  
6 1077-85.
- 7 20. Le Bot A, Michelet D, Hilly J, et al. Efficacy of intraoperative dexmedetomidine  
8 compared with placebo for surgery in adults: a meta-analysis of published  
9 studies. *Minerva Anesthesiol.* 2015 Oct; 81(10): 1105-17.
- 10 21. Kang SH, Kim YS, Hong TH, et al. Effects of dexmedetomidine on  
11 inflammatory responses in patients undergoing laparoscopic cholecystectomy.  
12 *Acta Anesthesiol Scand.* 2013 Apr; 57(4): 480-7.
- 13 22. Li Y, Wang B, Zhang LL, et al. Dexmedetomidine Combined with General  
14 Anesthesia Provides Similar Intraoperative Stress Response Reduction When  
15 Compared with a Combined General and Epidural Anesthetic Technique.  
16 *Anaesth Analg.* 2016 Apr; 122(4): 1202-10
- 17 23. Cao JL, Pei YP, Wei JQ, et al. Effects of intraoperative dexmedetomidine with  
18 intravenous anesthesia on postoperative emergence agitation/delirium in  
19 pediatric patients undergoing tonsillectomy with or without adenoidectomy: A  
20 CONSORT-prospective, randomized, controlled clinical trial. *Medicine*  
21 (Baltimore). 2016 Dec; 95(49): e5566.
- 22 24. Sun Y, Liu J, Yuan X, et al. Effects of dexmedetomidine on emergence delirium  
23 in pediatric cardiac surgery. *Minerva Pediatr.* 2017 Jun; 69(3): 165-173.
- 24 25. Li X, Yang J, Nie XL, et al. Impact of dexmedetomidine on the incidence of  
25 delirium in elderly patients after cardiac surgery: A randomized controlled trial.

- 1 PLoS One. 2017 Feb 9; 12(2): e0170757.
- 2 26. Yang X, Li Z, Gao C, et al. Effect of dexmedetomidine on preventing agitation  
3 and delirium after microvascular free flap surgery: a randomized, double-blind,  
4 control study. *J Oral Maxillofac Surg*. 2015 Jun; 73(6): 1065-72.
- 5 27. Deiner S, Luo X, Lin HM, et al. Intraoperative Infusion of Dexmedetomidine for  
6 Prevention of Postoperative Delirium and Cognitive Dysfunction in Elderly  
7 Patients Undergoing Major Elective Noncardiac Surgery: A Randomized Clinical  
8 Trial. *JAMA Surg*. 2017 Aug 16; 152(8): e171505.
- 9 28. Collin C, Wade DT, Davies S, et al. The Barthel ADL Index: a reliability study.  
10 *Int Disabil Stud*. 1988; 10: 61-3.
- 11 29. Katzman R, Zhang MY, Ouang-Ya-Qu, et al. A Chinese version of the Mini-  
12 Mental State Examination; impact of illiteracy in a Shanghai dementia survey. *J*  
13 *Clin Epidemiol*. 1988; 41: 971-8.
- 14 30. Inouye SK, van Dyck CH, Alessi CA, et al. Clarifying confusion: The Confusion  
15 Assessment Method. A new method for detecting delirium. *Ann Intern Med*  
16 1990; 113:941-8
- 17 31. Mu DL, Zhang DZ, Wang DX, et al. Parecoxib Supplementation to Morphine  
18 Analgesia Decreases Incidence of Delirium in Elderly Patients After Hip or Knee  
19 Replacement Surgery: A Randomized Controlled Trial. *Anaesth Analg*. 2017  
20 Jun; 124(6):1992-2000.
- 21 32. Ely EW, Margolin R, Francis J, et al. Evaluation of delirium in critically ill  
22 patients: validation of the Confusion Assessment Method for the Intensive Care  
23 Unit (CAM-ICU). *Crit Care Med*. 2001 Jul; 29(7): 1370-9.
- 24 33. Liu P, Li YW, Wang XS, et al. High serum interleukin-6 level is associated with  
25 increased risk of delirium in elderly patients after noncardiac surgery: a

- 1  
2  
3 1 prospective cohort study. *Chin Med J (Engl)*. 2013;126(19):3621-7.  
4  
5 2 34. Ji L, Vc L, Leung CM, et al. Clinical utility and validation of two instruments  
6  
7 3 (the Confusion Assessment Method Algorithm and the Chinese version of  
8  
9 4 Nursing Delirium Screening Scale) to detect delirium in geriatric inpatients. *Gen*  
10  
11 5 *Hosp Psychiatry*. 2008; 30: 171-6.  
12  
13 6 35. Wang C, Wu Y, Yue P, et al. Delirium assessment using Confusion Assessment  
14  
15 7 Method for the Intensive Care Unit in Chinese critically ill patients. *J Crit Care*.  
16  
17 8 2013; 28: 223-9.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Figure legend**

1

2

3 Figure 1. Flowchart of this study.

For peer review only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 1 Flow chart of this study




Figure1 Flow chart of this study

209x297mm (300 x 300 DPI)





SPIRIT 2013 Checklist: Recommended items to address in a clinical trial protocol and related documents\*

Section/item	Item No	Description	Addressed on page number
<b>Administrative information</b>			
Title	1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	P1/L1-3
Trial registration	2a	Trial identifier and registry name. If not yet registered, name of intended registry	P4/L7-8; P8/L14-16; P19/L4-7
	2b	All items from the World Health Organization Trial Registration Data Set	P8/L3-L17
Protocol version	3	Date and version identifier	P8/L11
Funding	4	Sources and types of financial, material, and other support	P19/L18-21
Roles and responsibilities	5a	Names, affiliations, and roles of protocol contributors	P1/L5-125; P2/L1
	5b	Name and contact information for the trial sponsor	P19/L18-21
	5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities	P19/L18-21
	5d	Composition, roles, and responsibilities of the coordinating centre, steering committee, endpoint adjudication committee, data management team, and other individuals or groups overseeing the trial, if applicable (see Item 21a for data monitoring committee)	N/A

**Introduction**

1				
2				
3	Background and	6a	Description of research question and justification for undertaking the trial, including summary of relevant	P6-7
4	rationale		studies (published and unpublished) examining benefits and harms for each intervention	
5				
6		6b	Explanation for choice of comparators	P7/L8-22; P10/L12-22
7				
8	Objectives	7	Specific objectives or hypotheses	P7/L23-25
9				
10	Trial design	8	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group),	
11			allocation ratio, and framework (eg, superiority, equivalence, noninferiority, exploratory)	P8/L3-9
12				
13				
14	<b>Methods: Participants, interventions, and outcomes</b>			
15				
16	Study setting	9	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will	P8/L7-9
17			be collected. Reference to where list of study sites can be obtained	
18				
19	Eligibility criteria	10	Inclusion and exclusion criteria for participants. If applicable, eligibility criteria for study centres and	P8/L18-P9/L3
20			individuals who will perform the interventions (eg, surgeons, psychotherapists)	
21				
22	Interventions	11a	Interventions for each group with sufficient detail to allow replication, including how and when they will be	P10/L12-22
23			administered	
24				
25		11b	Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose	P10/L23-P11/L4
26			change in response to harms, participant request, or improving/worsening disease)	
27				
28		11c	Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence	P10-11
29			(eg, drug tablet return, laboratory tests)	
30				
31		11d	Relevant concomitant care and interventions that are permitted or prohibited during the trial	N/A
32				
33	Outcomes	12	Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood	P12/L15-P14/L3
34			pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg,	
35			median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen	
36			efficacy and harm outcomes is strongly recommended	
37				
38	Participant timeline	13	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for	Figure 1
39			participants. A schematic diagram is highly recommended (see Figure)	
40				
41				
42				
43				
44				
45				
46				
47				

1				
2				
3	Sample size	14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	P14/L19-P15/L2
4				
5	Recruitment	15	Strategies for achieving adequate participant enrolment to reach target sample size	P9/L5-14
6				
7				

### 8 **Methods: Assignment of interventions (for controlled trials)**

#### 9 Allocation:

10				
11				
12	Sequence generation	16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	P9/L16-21
13				
14				
15				
16				
17	Allocation concealment mechanism	16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	P9/L22-P10/L5
18				
19				
20				
21	Implementation	16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	P9-/L18-21
22				
23				
24	Blinding (masking)	17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	P9/L22-25
25				
26				
27		17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	P10/L1-5
28				
29				
30				

### 31 **Methods: Data collection, management, and analysis**

32				
33	Data collection methods	18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known. Reference to where data collection forms can be found, if not in the protocol	P12/L1-P15/L15
34				
35				
36				
37				
38		18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols	P12/L20-23
39				
40				
41				
42				
43				
44				
45				
46				
47				

1				
2				
3	Data management	19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	P14/L5-17
4				
5				
6				
7	Statistical methods	20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	P14/L20-P15/L15
8				
9				
10		20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	P15/L4-11
11				
12		20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any statistical methods to handle missing data (eg, multiple imputation)	P11/L3-4; P12/L20-23
13				
14				
15				
16				
17	<b>Methods: Monitoring</b>			
18	Data monitoring	21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent from the sponsor and competing interests; and reference to where further details about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not needed	P14/L11-14
19				
20				
21				
22				
23				
24		21b	Description of any interim analyses and stopping guidelines, including who will have access to these interim results and make the final decision to terminate the trial	P14/L14-17
25				
26				
27	Harms	22	Plans for collecting, assessing, reporting, and managing solicited and spontaneously reported adverse events and other unintended effects of trial interventions or trial conduct	P13/L23-P14/L3
28				
29				
30	Auditing	23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent from investigators and the sponsor	P14/L11-14
31				
32				
33	<b>Ethics and dissemination</b>			
34				
35	Research ethics approval	24	Plans for seeking research ethics committee/institutional review board (REC/IRB) approval	P8/L11-14
36				
37				
38	Protocol amendments	25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes, analyses) to relevant parties (eg, investigators, REC/IRBs, trial participants, trial registries, journals, regulators)	P8/L11-14
39				
40				
41				
42				
43				
44				
45				
46				
47				

1				
2				
3	Consent or assent	26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	P8/L16-17 P19/L3-7
4				
5				
6		26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	N/A
7				
8	Confidentiality	27	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	P14/L5-7
9				
10				
11	Declaration of interests	28	Financial and other competing interests for principal investigators for the overall trial and each study site	P19/L12-16
12				
13				
14	Access to data	29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	P14/L11-14; P20/L12
15				
16				
17	Ancillary and post-trial care	30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	P14/L1-3
18				
19				
20	Dissemination policy	31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	P4/L4-5; P19/L9-10
21				
22				
23				
24				
25		31b	Authorship eligibility guidelines and any intended use of professional writers	P19/L23-P20/L2
26				
27		31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	P20/L12
28				
29	<b>Appendices</b>			
30				
31	Informed consent materials	32	Model consent form and other related documentation given to participants and authorised surrogates	N/A
32				
33				
34	Biological specimens	33	Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies, if applicable	N/A
35				
36				

37 \*It is strongly recommended that this checklist be read in conjunction with the SPIRIT 2013 Explanation & Elaboration for important clarification on the items.  
 38 Amendments to the protocol should be tracked and dated. The SPIRIT checklist is copyrighted by the SPIRIT Group under the Creative Commons  
 39 "[Attribution-NonCommercial-NoDerivs 3.0 Unported](https://creativecommons.org/licenses/by-nc-nd/3.0/)" license.  
 40

# BMJ Open

## Impact of dexmedetomidine infusion during general anaesthesia on incidence of postoperative delirium in elderly patients after major non-cardiac surgery: study protocol of a randomized, double-blinded and placebo-controlled trial

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-019549.R3
Article Type:	Protocol
Date Submitted by the Author:	26-Mar-2018
Complete List of Authors:	Wang, Bojie; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Li, Chunjing; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Hu, Jian; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Li, Huaijin; Peking University First Hospital, Department of Anesthesiology and Critical Care Guo, Chao; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Wang, Zhenhan; Dongping People's Hospital, Department of anesthesiology Zhang, Qiaochu; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine Mu, Dongliang; Department of Anesthesiology and Critical Care Medicine Wang, Dong-Xin; Peking University First Hospital, Department of Anesthesiology and Critical Care Medicine
<b>Primary Subject Heading</b>:	Anaesthesia
Secondary Subject Heading:	Anaesthesia
Keywords:	Elderly, Major non-cardiac surgery, Intraoperative dexmedetomidine, Postoperative delirium

SCHOLARONE™  
Manuscripts

1  
2  
3 **1 Impact of dexmedetomidine infusion during general anaesthesia on incidence of**  
4  
5 **2 postoperative delirium in elderly patients after major non-cardiac surgery: study**  
6  
7 **3 protocol of a randomized, double-blinded and placebo-controlled trial**  
8  
9  
10

11 Bo-Jie Wang<sup>1</sup>, Chun-Jing Li<sup>1</sup>, Jian Hu<sup>1</sup>, Huai-Jin Li<sup>1</sup>, Chao Guo<sup>1</sup>, Zhen-Han Wang<sup>2</sup>,  
12  
13 Qiao-Chu Zhang<sup>1</sup>, Dong-Liang Mu<sup>1,\*</sup>, Dong-Xin Wang<sup>1</sup>  
14  
15

16  
17  
18 <sup>1</sup>Department of Anesthesiology and Critical Care Medicine, Peking University First  
19  
20 Hospital, Beijing, China

21  
22 <sup>2</sup>Department of Anesthesiology, Dongping People's Hospital, Shandong, China  
23  
24

25  
26  
27 The first two authors (BJ Wang and CJ Li) contributed equally to this manuscript  
28  
29

30  
31 Corresponding author: Dong-Liang Mu, M.D., Department of Anesthesiology and  
32  
33 Critical Care Medicine, Peking University First Hospital, Beijing, China. Tel: +86  
34  
35 010 83572784 (O), +86 13810702725 (M); E-mail: mudongliang@icloud.com  
36  
37

38  
39  
40 Bo-Jie Wang: ivyxbd@hotmail.com

41  
42 Chun-Jing Li: gogolion@163.com

43  
44 Jian Hu: hj19871204@163.com

45  
46 Huai-Jin Li: sophie.lee.coffee@gmail.com

47  
48 Chao Guo: guochao1988@yeah.com

49  
50 Zhen-Han Wang: 1468432914@qq.com

51  
52 Qiao-Chu Zhang: 372193745@qq.com

53  
54 Dong-Liang Mu: mudongliang@icloud.com  
55  
56  
57  
58  
59  
60

1  
2  
3 1 Dong-Xin Wang: wangdongxin@hotmail.com  
4

5 2  
6

7 3 Word Count: 2888  
8

9 4  
10

11 5  
12

13 6  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only



## Abstract for protocol

**Introduction:** Delirium is a common complication in elderly after surgery and is associated with worse outcomes. Multiple risk factors are related with postoperative delirium, such as exposure to general anaesthetics, pain and postoperative inflammatory response. Preclinical and clinical studies have shown that dexmedetomidine attenuated neurotoxicity induced by general anaesthetics, improved postoperative analgesia and inhibited inflammatory response after surgery. Several studies found that intraoperative use of dexmedetomidine can prevent postoperative delirium, but data were inconsistent. This study was designed to investigate the impact of dexmedetomidine administered during general anaesthesia in preventing delirium in elderly after major non-cardiac surgery.

**Method and analysis:** This is a randomized, double-blinded, and placebo-controlled trial. 620 elderly patients (age  $\geq 60$  years) who are scheduled to undertake elective major non-cardiac surgery (with an expected duration  $\geq 2$  hours) are randomly divided into two groups. For patients in the dexmedetomidine group, a loading dose dexmedetomidine (0.6  $\mu\text{g}/\text{kg}$ ) will be administered in 10 minutes before anaesthesia induction, followed by a continuous infusion at a rate of 0.5  $\mu\text{g}/\text{kg}/\text{h}$  till 1 hour before the end of surgery. For patients in the control group, normal saline will be administered with an identical rate as in the dexmedetomidine group. The primary endpoint is the incidence of delirium during the first five postoperative days. Secondary endpoints include pain intensity, cumulative opioid consumption and subjective sleep quality during the first 3 postoperative days as well as the incidence of non-delirium complications and all-cause mortality within 30 days after surgery.

1  
2  
3 1 **Ethics and dissemination:** The study protocol was approved by Clinical Research  
4  
5 2 Ethics Committee of Peking University First Hospital (2015-987) and registered at  
6  
7 3 Chinese Clinical Trial Registry (<http://www.chictr.org.cn>) with identifier ChiCTR-  
8  
9 4 IPR-15007654. The results of the study will be presented at academic conferences  
10  
11 5 and submitted to peer-reviewed journals.  
12  
13  
14 6

15  
16 7 **Key words:** Elderly; Major non-cardiac surgery; Intraoperative dexmedetomidine;  
17  
18 8 Postoperative delirium  
19  
20 9

## Strengths and limitations

### Strengths

1. The study design will be a randomized, double-blinded and placebo-controlled, with a relative large sample size.
2. Anaesthesia depth (Bispectral Index) will be monitored to guide anaesthesia maintenance to ensure patients in both groups received equal depth anaesthesia.

### Limitations

1. This is a single center trial, which will limit the generalizability of our results.
2. Only early outcomes (up to 30 days after surgery) are assessed in this trial.
3. The hemodynamic and anaesthetic-sparing effects of dexmedetomidine might weaken the efficiency of blindness to the treating anaesthesiologist.

## Introduction

Delirium is a transient brain dysfunction which is characterized by altered consciousness, inattention, and changes in cognition or perception; it develops acutely with clinical manifestations to be fluctuated with aggressive and depressive manner during developing course.<sup>1</sup> Prevalence of delirium varies from 12% to 51% in patients after non-cardiac surgery, and it is increased with age.<sup>2-3</sup> The occurrence of postoperative delirium (POD) is associated with worsen outcomes including prolonged mechanical ventilation and intensive care unit stay, increased postoperative complications, high mortality rate and long-term cognitive decline.<sup>2-5</sup>

The aetiology of POD is multifactorial and includes several intraoperative factors.<sup>6</sup> For example, exposure to general anaesthetics (e.g., propofol or sevoflurane) might produce neurotoxicity,<sup>7-8</sup> whereas reduced anaesthetic consumption by avoiding deep anaesthesia reduced the occurrence of delirium in elderly patients undergoing major non-cardiac surgery.<sup>9</sup> Poor pain management is another risk factor of POD.<sup>6,10</sup> It was reported that the risk of postoperative delirium was 1.2 times higher for every unit increment in visual analogue pain score (an 11-point pain scale where 0 indicates no pain and 10 the most severe pain).<sup>11</sup> Inflammation is also proposed to play an important role in the pathogenesis of POD.<sup>2</sup> Inflammatory responses induced by surgery and anaesthesia are manifested by elevated levels of interleukins (IL), C-reactive protein (CRP) and tumor necrosis factor (TNF).<sup>2,12</sup> Studies by our group and others found that higher levels of inflammatory mediators are associated with increased risk of POD.<sup>2,13-16.</sup>

Dexmedetomidine is a highly selective  $\alpha_2$ -receptor agonist with sedative, analgesic and anxiolytic effects.<sup>17-21</sup> When used as a supplement during intraoperative

1 anaesthesia, it reduces the consumption of general anaesthetics.<sup>17</sup> Preclinical evidence  
2 suggested that use of dexmedetomidine might attenuate neurotoxicity induced by  
3 general anaesthetics.<sup>18-19</sup> In a meta-analysis, intraoperative administration of  
4 dexmedetomidine lowers postoperative pain intensity and reduces opioid  
5 consumption.<sup>20</sup> Clinical evidence also showed that intraoperative dexmedetomidine  
6 significantly inhibits hyper-secretion of inflammatory cytokines during and after  
7 surgery.<sup>21-22</sup>

8       Use of dexmedetomidine during general anaesthesia may reduce POD. In  
9 pediatric patients undergoing tonsillectomy and cardiac surgery, intraoperative  
10 infusion of dexmedetomidine lowered the incidence of emergence delirium.<sup>23,24</sup> In  
11 adult patients undergoing cardiac surgery and microvascular free flap surgery,  
12 intraoperative dexmedetomidine (comparison with normal saline) slightly decreased  
13 the incidence of delirium, although the differences were not statistically significant  
14 between two groups possibly due to underpowered sample size.<sup>25,26</sup> In a recent study  
15 of Deiner et al.,<sup>27</sup> use of dexmedetomidine during general anaesthesia did not reduce  
16 delirium after major non-cardiac surgery in the elderly. However, in that study,  
17 anaesthesia depth was not monitored and the consumption of anaesthetics (such as  
18 propofol and fentanyl) was similar between the two groups. It was possible that  
19 patients in the dexmedetomidine group had deeper anaesthesia which might have  
20 increased the risk of delirium.<sup>27</sup> Therefore, the effect of dexmedetomidine  
21 administered during general anaesthesia on the occurrence of POD needs to be  
22 evaluated further.

23       This study is designed to investigate whether dexmedetomidine use during  
24 general anaesthesia can decrease the incidence of POD in elderly after major non-  
25 cardiac surgery.

## Method and analysis

### Study design

This randomized, double-blinded, and placebo-controlled trial with two parallel arms was designed to test the superiority of dexmedetomidine administered during general anaesthesia on the incidence of delirium after surgery. Patients will be randomized into either the dexmedetomidine group or the control group (Figure 1). The study is conducted in the Department of Anesthesiology and Critical Care Medicine of Peking University First Hospital.

### Ethics approval

The study protocol (version 1.1, issue date Nov, 2015) was approved by Clinical Research Ethics Committee of Peking University First Hospital (2015-987). Any protocol modification will be submitted for review and approval by the Ethics Committee. The trial was registered at Chinese Clinical Trial Registry (<http://www.chictr.org.cn>) on December 1, 2015, with identifier ChiCTR-IPR-15007654. Written informed consents are obtained from every patient or his/her surrogate in law.

### Patient and public involvement

Patients and public were not involved in study design or conduct of study. There is no plan to disseminate the results to study participants.

### Participants

Elderly patients (age  $\geq 60$  years) who are scheduled to undergo elective non-cardiac surgery with expected duration  $\geq 2$  hours under general anaesthesia are screened for inclusion. Those who meet any of the following criteria will be excluded: (1) do not provide written informed consents; (2) previous history of schizophrenia, epilepsy or

1  
2  
3 1 Parkinson disease; (3) visual, hearing, language or other barrier which impede  
4  
5 2 communication and preoperative delirium assessment; (4) neurosurgery or traumatic  
6  
7 3 brain injury; (5) severe bradycardia (heart rate less than 40 beats per minute), sick  
8  
9 4 sinus syndrome or atrioventricular block of degree 2 or above; (6) severe hepatic  
10  
11 5 dysfunction (Child-Pugh grade C); or (7) renal failure (requirement of renal  
12  
13 6 replacement therapy).

### 7 **Patient recruitment and baseline data collection**

8 Potential participants are screened by investigators the day before surgery (or on  
9  
10 Friday for those who will undergo surgery next Monday). The study protocol  
11  
12 including potential risks and benefits will be explained to patients in person. Those  
13  
14 who do not meet the exclusion criteria are invited to participate in the study.

15 After obtaining written informed consents, the following baseline data are  
16  
17 collected: demographic data, preoperative diagnosis, comorbidity, current medical  
18  
19 therapy, previous surgery, and main results of physical and laboratory examinations.  
20  
21 Barthel index is used for evaluation of activities of daily living.<sup>28</sup> Cognitive function  
22  
23 is assessed with Mini-Mental State Examination (MMSE).<sup>29</sup> Preoperative delirium is  
24  
25 assessed with confusion assessment method (CAM).<sup>30</sup>

### 26 **Randomization, grouping and blinding**

27  
28 Random numbers were created by an independent statistician using SAS statistical  
29  
30 package version 9.3 (SAS Institute, Cary, NC, USA) in a 1:1 ratio and were sealed in  
31  
32 envelopes. A study coordinator, who has no knowledge of patients before  
33  
34 randomization and does not participate in anaesthesia and postoperative follow-up of  
35  
36 enrolled patients, will open envelop for random numbers and prepare study drugs  
37  
38 before induction of anaesthesia.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 Information regarding randomization, study drug preparation and group  
2 allocation will be masked from investigators who perform data collection and  
3 postoperative follow-up, anaesthesiologists, patients and other healthcare team  
4 members. Blinding will be maintained throughout the entire study period.

5 To ensure patients' safety, study group allocation can be unmasked in the  
6 following conditions, i.e., occurrence of severe adverse events or any unexpected  
7 deterioration of patient's clinical status. These situations will be documented in the  
8 Case Report Forms (CRFs). The unmasked patients will be included in the intention-  
9 to-treat population but excluded from per-protocol analysis.

#### 10 **Interventions, anaesthesia and analgesia**

11 Intraoperative monitoring includes electrocardiogram, noninvasive blood pressure,  
12 pulse oxygen saturation, end-tidal carbon dioxide, nasopharyngeal temperature, urine  
13 output, and bispectral index (BIS). Intra-arterial pressure (including derivative  
14 dynamic parameters such as stroke volume variation by FlowTrac system) and central  
15 venous pressure are monitored according to patients' conditions.

16 Study drugs, either 200 µg (2 ml) dexmedetomidine (Jiangsu Hengrui Medicine  
17 Co, Ltd, Jiangsu, China) or 2 ml normal saline, are diluted into 50 ml normal saline.  
18 All study drugs are colorless solution provided in syringes of the same size and brand.  
19 The regimen of study drug administration includes a loading dose of 0.15 ml/kg (i.e.,  
20 0.6 µg/kg dexmedetomidine for patients in the dexmedetomidine group) administered  
21 during a 10-minute period before anaesthesia induction. This is followed by a  
22 continuous infusion at a rate of 0.125 ml/kg/h (i.e., a rate of 0.5 µg/kg/h  
23 dexmedetomidine for patients in the dexmedetomidine group) till 1 hour before the  
24 end of surgery. Study drug infusion is performed by an infusion pump specially



1  
2  
3 1 designed for dexmedetomidine administration (Slgo<sup>®</sup> CP1000, Beijing Slgo medical  
4  
5 2 technology Co., Ltd.).  
6

7 3 Attending anaesthesiologists can decrease or stop study drug infusion in the  
8  
9 4 following conditions: (1) severe bradycardia or hypotension which does not improve  
10  
11 5 after routine treatment; (2) new onset atrioventricular block which does not improve  
12  
13 6 after routine treatment; or (3) other conditions that anaesthesiologists consider it  
14  
15 7 necessary. In these conditions, the reasons that lead to any protocol deviations will be  
16  
17 8 recorded in the case report forms (CRF). These patients will be included in the  
18  
19 9 intention-to-treat analysis but excluded from the per-protocol analysis.  
20  
21

22 10 Anaesthesia is induced with intravenous sufentanil (target controlled infusion  
23  
24 11 with effect-site concentration from 0.2 to 0.5 ng/ml) and propofol (2-3 mg/kg) and  
25  
26 12 maintained with intravenous sufentanil (effect-site concentration from 0.2 to 0.5  
27  
28 13 ng/ml) and propofol (4-12 mg/kg/h) and inhalation of a 1:1 nitrous oxide-oxygen  
29  
30 14 mixture. Rocuronium and/or cisatracurium are administered for muscle relaxation.  
31  
32 15 Patients will be mechanically ventilated with a tidal volume of 6-8 ml/kg and a  
33  
34 16 positive end-expiratory pressure of 5 cm H<sub>2</sub>O. The mean arterial pressure is  
35  
36 17 maintained above 60 mmHg or within 20% from baseline. BIS is maintained between  
37  
38 18 40 and 60. Body temperature is maintained with air-warming and fluid heating  
39  
40 19 systems. The target of nasopharyngeal temperature maintenance during surgery is  
41  
42 20 from 36.0 to 37°C.  
43  
44  
45

46 21 All patients are transferred to post-anaesthesia care unit (PACU) or intensive  
47  
48 22 care unit (ICU) before they are sent back to general wards. Patient-controlled  
49  
50 23 intravenous analgesia (PCIA) is provided for all patients, which is established with  
51  
52 24 0.5 mg/ml morphine in 100 ml normal saline and programmed to deliver a 1 mg bolus  
53  
54 25 with a lock-out interval of 8 minutes and a background infusion at 0.5 mg/h.  
55  
56  
57  
58  
59  
60

1 Supplemental morphine at dose of 2 to 4 mg will be administered at 10-minute  
2 intervals if the numeric rating scale (NRS) pain score (a 11-point scale where 0  
3 indicates no pain and 10 indicates the worst pain) remains above 4 after 3 consecutive  
4 PCIA boluses.<sup>31</sup> Other postoperative managements were performed according to  
5 routine practice.

## 6 **Outcome assessment**

7 Patients are followed up twice daily during the first 5 postoperative days and then  
8 weekly until 30 days after surgery. Investigators who are responsible for  
9 postoperative follow-up are not involved in anaesthesia and perioperative care, and  
10 are not allowed to exchange patients' information with anaesthesiologists who take  
11 care of patients in the operating room. Before the beginning of the study, investigators  
12 are trained to follow the study protocol and to perform delirium assessment and the  
13 training process is repeated at 4 to 6 months intervals during the study period.<sup>2-3, 31</sup>  
14 The 4-hour training courses of delirium assessment include the following contents: (1)  
15 lectures regarding signs/symptoms, diagnosis and treatment of delirium by  
16 psychiatrists; (2) training courses of the use of CAM and CAM-ICU on patient-actors  
17 (trained ICU physicians or nurses who act as patients with or without delirium)  
18 conducted by psychiatrists. The process continued until 100% agreement is achieved  
19 in diagnosing delirium.

## 20 **Primary endpoint**

21 The primary endpoint is the incidence of delirium during the first 5 days after surgery.  
22 Delirium is assessed twice daily (at 08:00-09:00 and 19:00-20:00, respectively) with  
23 CAM for non-intubated patients<sup>30</sup> or CAM for the Intensive Care Unit (CAM-ICU)  
24 for intubated patients.<sup>32</sup> These delirium assessment methods had been used in our  
25 previous studies.<sup>2-3, 31</sup> For patients who are discharged or died within 5 days after

1 surgery, the results of last delirium assessment will be considered the results of the  
2 missing data. These patients will be excluded when calculating daily prevalence of  
3 delirium in a post-hoc analysis.

#### 4 ***Secondary endpoints***

5 Postoperative pain intensities at rest and with movement are assessed with NRS pain  
6 score at 24, 48 and 72 hours after surgery, respectively.<sup>31</sup> Cumulative morphine  
7 consumptions at these time points are recorded. Subjective sleep quality is assessed  
8 with NRS (an 11-point scale where 0 indicates the worst possible sleep and 10 the  
9 best possible sleep) at 08:00 on the first, second and third morning after surgery.<sup>3,31</sup>  
10 Other secondary endpoints include non-delirium complications within 30 days after  
11 surgery, length of stay in hospital after surgery and all-cause 30-day mortality. Non-  
12 delirium complications are generally defined as new-onset non-delirium conditions  
13 after surgery that are harmful to patients' recovery and require therapeutic  
14 intervention.

#### 15 **Safety outcomes**

16 In the present study, adverse events are monitored from the start of study drug  
17 administration until PACU discharge or 2 hours after ICU admission. Hypotension is  
18 defined as systolic blood pressure of less than 90 mmHg or a decrement of more than  
19 30% from baseline. Hypertension is defined as systolic blood pressure of more than  
20 180 mmHg or an increment of more than 30% from baseline. Bradycardia is defined  
21 as heart rate of less than 40 beats per minute. Tachycardia is defined as heart rate of  
22 more than 100 beats per minute. Desaturation is defined as SpO<sub>2</sub> of less than 90%.  
23 Emergence agitation is defined as a Richmond Agitation-Sedation Scale (RASS)  
24 score of more than +2 within 30 minutes after extubation. Delayed extubation is

1 defined when time to extubation is more than 2 hours (from the end of surgery) in  
2 PACU patients or more than 4 hours in ICU patients.<sup>2, 22-25</sup>

3 Severe adverse events, i.e., those that might result in patient's  
4 disability/deformity, prolong in-hospital stay, or life threatening events, will be  
5 reported to Clinical Research Ethics Committee of Peking University First Hospital  
6 within 24 hours. For patients who suffered harm from present trial, medical treatment  
7 will be initiated as soon as possible and compensation will be completed according to  
8 local laws and regulations.

### 9 **Data monitoring and management**

10 Original data will be recorded in the CRFs accordingly. All data will be kept  
11 confidentially. The completed CRFs will be checked by a study coordinator who is  
12 qualified by the principal investigator. Supplementations and corrections will be made  
13 when necessary. Data entry will be performed in a double-input and double-check  
14 way with the Data Management System (Fantastic Eight Tech. Co., Ltd, Beijing,  
15 China) of the Peking University First Hospital.

16 The conduct of the study and the quality of data will be monitored by the  
17 Clinical Research Ethics Committee of Peking University First Hospital. Data  
18 management and statistical analysis will be performed by the Department of  
19 Biostatistics of Peking University First Hospital. Considering that dexmedetomidine  
20 has been widely used during general anaesthesia and its safety has been confirmed, no  
21 interim analysis will be performed and the trial will continue until the target sample  
22 size is achieved.

### 23 **Statistical analysis**

#### 24 *Sample size calculation*

1 In our previous study, the incidence of delirium was 14.8% in elderly patients after  
2 non-cardiac surgery.<sup>33</sup> Previous studies reported that intraoperative dexmedetomidine  
3 decreased the incidence of POD by 60-77% in comparison with placebo.<sup>22, 25</sup> We  
4 assumed that the incidence of POD will be reduced from 14.8% to 7.4% (i.e., a 50%  
5 reduction) in the present study. With the power set at 80% and significant level at  
6 0.05, 564 patients are required to detect the difference. Considering a loss to follow-  
7 up rate of about 9%, we plan to enroll 620 patients in this study.

### 8 ***Outcome analysis***

9 Continuous data with normal distribution will be compared using independent sample  
10 T-test. Continuous data with asymmetric distribution will be compared using  
11 independent sample Mann-Whitney U test. Categorical data will be compared using  
12 Chi-squared test or continuity correction Chi-squared test. The difference (and 95%  
13 confidence interval of the difference) between two means or medians will be  
14 estimated using the methodology of Levene's test or Hodges-Lehmann estimator.  
15 Time-to-event data will be analyzed by survival analysis with differences between  
16 groups compared with log-rank test.

17 Statistical analyses will be performed with the SPSS 14.0 (SPSS, Inc., Chicago,  
18 IL) and SAS 9.3 (SAS Institute, Cary, NC, USA). All tests are two tailed, and P  
19 values of less than 0.05 are considered to be statistically significant. The Bonferroni  
20 adjustment is made to control type I error for multiple testing.

21

## Discussion

This randomized, double-blinded, and placebo-controlled single center trial was designed to investigate if dexmedetomidine administration during general anaesthesia can decrease the incidence of POD in elderly patients after major non-cardiac surgery.

In the present study, the dosing regimen of dexmedetomidine is similar to our previous study because it does not increase drug-related adverse events (such as severe bradycardia and hypotension).<sup>25</sup> Furthermore, the CAM and CAM-ICU are used to assess delirium in patients with or without intubation, respectively.<sup>30, 32</sup> Both CAM and CAM-ICU have been validated in Chinese population.<sup>34, 35</sup> Feasibility of these two assessment tools has been confirmed in our previous studies.<sup>2-3, 31</sup> To maintain the quality of delirium assessment, investigators in charge of postoperative follow-up are trained by a psychiatrist before the study and will be retrained at 4 to 6-month intervals.

Because of the hemodynamic and anaesthetic-sparing effects of dexmedetomidine, it is not very difficult for the experienced anaesthesiologists to guess which study drug is administered. This might weaken the blinding to anaesthesiologists. However, in the present study, investigators who are responsible for postoperative follow-up and delirium assessment are not involved in anaesthesia and perioperative care; and they are not allowed to exchange patients' information with anaesthesiologists who take care of patients in the operating room. In this way, the blinding of investigators to study group assignment can be guaranteed.

The strengths of the present study include the following when compared with previous studies.<sup>22-24</sup> Firstly, a randomized, double-blind, and placebo-controlled study design with a relative large sample size (620 patients) is adopted. Results of the

1 study will provide high quality evidences. Secondly, BIS level is monitored in all  
2 enrolled patients, which will help us to avoid unnecessary and potentially harmful  
3 deep anaesthesia. Thirdly, safety data will be recorded in detail. Our study also has  
4 some limitations. One is that this is a single center trial, which will limit the external  
5 validity of our results. Second, only early outcomes (up to 30 days after surgery) will  
6 be explored. Third, the hemodynamic and anaesthetic-sparing effects of  
7 dexmedetomidine might weaken the efficiency of blindness to the treating  
8 anaesthesiologist.

9  
10

11 **Trail status:**

12 This study is currently at stage of patient enrollment and data collection. The current  
13 version of study protocol is V1.1 and is approved on November 27, 2015. Patient  
14 recruitment started from December 2, 2015 and is expected to be finished by March  
15 31, 2018.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60





## Declarations

**Ethics approval and consent to participate:** The study protocol was approved by Clinical Research Ethics Committee of Peking University First Hospital (2015-987). The trial was registered at Chinese Clinical Trial Registry (www.chictr.org.cn) with identifier ChiCTR-IPR-15007654 on December 1, 2015). Written informed consent will be obtained from all patients or their surrogate in law.

**Dissemination:** Results of the study will be presented at academic conferences and submitted to peer-reviewed journals.

**Competing interests:** DXW reports lecture fees and travel expenses for lectures given at academic meetings from Jiangsu Hengrui Medicine Co Ltd, China, and Yichang Humanwell Pharmaceutical Co Ltd, China. DLM is primary investigator of present study which was supported by Beijing Excellent Talent Support Program. Other authors reported no conflict of interests.

**Funding:** This trial was supported by Beijing Excellent Talent Support Program (No. 2014000020124G025). The sponsors have no role in the study design and conduct; the collection, management, analysis, and interpretation of the data; or the preparation and approval of the manuscript.

**Authors' contributions:** DXW and DLM designed this study. DLM draft the manuscript of protocol. DXW critically revised the manuscript. DLM, BJW, CJL, JH,

1 HJL, CG, ZHW and QCZ participate in the conduct of the study. All authors read and  
2 approved the final manuscript.

3  
4  
5  
6  
7  
8  
9 **Acknowledgements:** We thank professor Xin-Yu Sun (Department of Psychiatrics,  
10 Peking University Sixth Hospital, Beijing, China) for her help in psychiatric  
11 consultation and personnel training and Dr. Xue-Ying Li (Department of Biostatistics,  
12 Peking University First Hospital, Beijing, China) for her help in statistical  
13 consultation. We also thank Professor Daqing Ma (Section of Anesthetics, Pain  
14 Medicine and Intensive Care, Department of Surgery and Cancer, Imperial College  
15 London, London, UK) for his help in revising the manuscript.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27 **Data sharing:** Will be provided on request.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) Washington, DC: American Psychiatric Association; 1994.
2. Inouye SK, Westendorp RG, Saczynski JS. Delirium in elderly people. *Lancet*. 2014 Mar 8;383(9920):911-22.
3. Su X, Meng ZT, Wu XH, et al. Dexmedetomidine for prevention of delirium in elderly patients after non-cardiac surgery: a randomised, double-blind, placebo-controlled trial. *Lancet*. 2016 Oct 15; 388(10054): 1893-1902.
4. Ely EW, Shintani A, Truman B, et al. Delirium as a predictor of mortality in mechanically ventilated patients in the intensive care unit. *JAMA*. 2004 Apr 14; 291(14): 1753-62.
5. Saczynski JS, Marcantonio ER, Quach L, et al. Cognitive trajectories after postoperative delirium. *N Engl J Med*. 2012 Jul 5; 367(1): 30-9.
6. Aldecoa C, Bettelli G, Bilotta F, et al. European Society of Anaesthesiology evidence-based and consensus-based guidelines on postoperative delirium. *Eur J Anaesthesiol*. 2017 Apr; 34(4):192-214.
7. Creeley C, Dikranian K, Dissen G, et al. Propofol-induced apoptosis of neurones and oligodendrocytes in fetal and neonatal rhesus macaque brain. *Br J Anaesth*. 2013 Jun; 110 Suppl 1: i29-38.
8. Xiong WX, Zhou GX, Wang B, et al. Impaired spatial learning and memory after sevoflurane-nitrous oxide anesthesia in aged rats is associated with down-regulated cAMP/CREB signaling. *PLoS One*. 2013 Nov 15;8(11): e79408.
9. Chan MT, Cheng BC, Lee TM, et al. BIS-guided anesthesia decreases

- 1  
2  
3 1 postoperative delirium and cognitive decline. *J Neurosurg Anesthesiol.* 2013 Jan;  
4  
5 2 25(1): 33-42.  
6  
7 3 10. Vaurio LE, Sands LP, Wang Y, et al. Postoperative delirium: the importance of  
8  
9 4 pain and pain management. *Anesth Analg.* 2006 Apr; 102(4): 1267-73.  
10  
11 5 11. Lynch EP, Lazor MA, Gellis JE, et al. The impact of postoperative pain on the  
12  
13 6 development of postoperative delirium. *Anesth Analg* 1998; 86:781-5  
14  
15 7 12. Hirsch J, Vacas S, Terrando N, et al. Perioperative cerebrospinal fluid and  
16  
17 8 plasma inflammatory markers after orthopedic surgery. *J Neuroinflammation.*  
18  
19 9 2016 Aug 30; 13(1): 211.  
20  
21 10 13. Cape E, Hall RJ, van Munster BC, et al. Cerebrospinal fluid markers of  
22  
23 11 neuroinflammation in delirium: a role for interleukin-1 $\beta$  in delirium after hip  
24  
25 12 fracture. *J Psychosom Res.* 2014 Sep; 77(3): 219-25.  
26  
27 13 14. van den Boogaard M, Kox M, Quinn KL, et al. Biomarkers associated with  
28  
29 14 delirium in critically ill patients and their relation with long-term subjective  
30  
31 15 cognitive dysfunction; indications for different pathways governing delirium in  
32  
33 16 inflamed and noninflamed patients. *Crit Care.* 2011; 15(6): R297.  
34  
35 17 15. McGrane S, Girard TD, Thompson JL, et al. Procalcitonin and C-reactive protein  
36  
37 18 levels at admission as predictors of duration of acute brain dysfunction in  
38  
39 19 critically ill patients. *Crit Care.* 2011; 15(2): R78.  
40  
41 20 16. Kazmierski J, Banys A, Latek J, et al. Raised IL-2 and TNF- $\alpha$  concentrations are  
42  
43 21 associated with postoperative delirium in patients undergoing coronary-artery  
44  
45 22 bypass graft surgery. *Int Psychogeriatr.* 2014 May; 26(5): 845-55.  
46  
47 23 17. Shin HW, Yoo HN, Kim DH, et al. Preanesthetic dexmedetomidine 1 $\mu$ g/kg  
48  
49 24 single infusion is a simple, easy, and economic adjuvant for general anesthesia.  
50  
51 25 Korean *J Anaesthesiol.* 2013 Aug; 65(2): 114-20.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 18. Li J, Xiong M, Nadavaluru PR, et al. Dexmedetomidine attenuates neurotoxicity  
4  
5 2 induced by prenatal propofol exposure. *J Neurosurg Anesthesiol.* 2016 Jan;  
6  
7 3 28(1): 51-64.  
8  
9 4 19. Sanders RD, Xu J, Shu Y, et al. Dexmedetomidine attenuates isoflurane-induced  
10  
11 5 neurocognitive impairment in neonatal rats. *Anesthesiology.* 2009 May; 110(5):  
12  
13 6 1077-85.  
14  
15 7 20. Le Bot A, Michelet D, Hilly J, et al. Efficacy of intraoperative dexmedetomidine  
16  
17 8 compared with placebo for surgery in adults: a meta-analysis of published  
18  
19 9 studies. *Minerva Anesthesiol.* 2015 Oct; 81(10): 1105-17.  
20  
21 10 21. Kang SH, Kim YS, Hong TH, et al. Effects of dexmedetomidine on  
22  
23 11 inflammatory responses in patients undergoing laparoscopic cholecystectomy.  
24  
25 12 *Acta Anesthesiol Scand.* 2013 Apr; 57(4): 480-7.  
26  
27 13 22. Li Y, Wang B, Zhang LL, et al. Dexmedetomidine Combined with General  
28  
29 14 Anesthesia Provides Similar Intraoperative Stress Response Reduction When  
30  
31 15 Compared with a Combined General and Epidural Anesthetic Technique.  
32  
33 16 *Anaesth Analg.* 2016 Apr; 122(4): 1202-10  
34  
35 17 23. Cao JL, Pei YP, Wei JQ, et al. Effects of intraoperative dexmedetomidine with  
36  
37 18 intravenous anesthesia on postoperative emergence agitation/delirium in  
38  
39 19 pediatric patients undergoing tonsillectomy with or without adenoidectomy: A  
40  
41 20 CONSORT-prospective, randomized, controlled clinical trial. *Medicine*  
42  
43 21 (Baltimore). 2016 Dec; 95(49): e5566.  
44  
45 22 24. Sun Y, Liu J, Yuan X, et al. Effects of dexmedetomidine on emergence delirium  
46  
47 23 in pediatric cardiac surgery. *Minerva Pediatr.* 2017 Jun; 69(3): 165-173.  
48  
49 24 25. Li X, Yang J, Nie XL, et al. Impact of dexmedetomidine on the incidence of  
50  
51 25 delirium in elderly patients after cardiac surgery: A randomized controlled trial.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1 PLoS One. 2017 Feb 9; 12(2): e0170757.
- 2 26. Yang X, Li Z, Gao C, et al. Effect of dexmedetomidine on preventing agitation  
3 and delirium after microvascular free flap surgery: a randomized, double-blind,  
4 control study. *J Oral Maxillofac Surg.* 2015 Jun; 73(6): 1065-72.
- 5 27. Deiner S, Luo X, Lin HM, et al. Intraoperative Infusion of Dexmedetomidine for  
6 Prevention of Postoperative Delirium and Cognitive Dysfunction in Elderly  
7 Patients Undergoing Major Elective Noncardiac Surgery: A Randomized Clinical  
8 Trial. *JAMA Surg.* 2017 Aug 16; 152(8): e171505.
- 9 28. Collin C, Wade DT, Davies S, et al. The Barthel ADL Index: a reliability study.  
10 *Int Disabil Stud.* 1988; 10: 61-3.
- 11 29. Katzman R, Zhang MY, Ouang-Ya-Qu, et al. A Chinese version of the Mini-  
12 Mental State Examination; impact of illiteracy in a Shanghai dementia survey. *J*  
13 *Clin Epidemiol.* 1988; 41: 971-8.
- 14 30. Inouye SK, van Dyck CH, Alessi CA, et al. Clarifying confusion: The Confusion  
15 Assessment Method. A new method for detecting delirium. *Ann Intern Med*  
16 1990; 113:941-8
- 17 31. Mu DL, Zhang DZ, Wang DX, et al. Parecoxib Supplementation to Morphine  
18 Analgesia Decreases Incidence of Delirium in Elderly Patients After Hip or Knee  
19 Replacement Surgery: A Randomized Controlled Trial. *Anaesth Analg.* 2017  
20 Jun; 124(6):1992-2000.
- 21 32. Ely EW, Margolin R, Francis J, et al. Evaluation of delirium in critically ill  
22 patients: validation of the Confusion Assessment Method for the Intensive Care  
23 Unit (CAM-ICU). *Crit Care Med.* 2001 Jul; 29(7): 1370-9.
- 24 33. Liu P, Li YW, Wang XS, et al. High serum interleukin-6 level is associated with  
25 increased risk of delirium in elderly patients after noncardiac surgery: a

- 1  
2  
3 1 prospective cohort study. *Chin Med J (Engl)*. 2013;126(19):3621-7.  
4  
5 2 34. Ji L, Ve L, Leung CM, et al. Clinical utility and validation of two instruments  
6  
7 3 (the Confusion Assessment Method Algorithm and the Chinese version of  
8  
9 4 Nursing Delirium Screening Scale) to detect delirium in geriatric inpatients. *Gen*  
10  
11 5 *Hosp Psychiatry*. 2008; 30: 171-6.  
12  
13 6 35. Wang C, Wu Y, Yue P, et al. Delirium assessment using Confusion Assessment  
14  
15 7 Method for the Intensive Care Unit in Chinese critically ill patients. *J Crit Care*.  
16  
17 8 2013; 28: 223-9.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60





1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 1 Flow chart of this study




Figure1 Flow chart of this study

209x297mm (300 x 300 DPI)



SPIRIT 2013 Checklist: Recommended items to address in a clinical trial protocol and related documents\*

Section/item	Item No	Description	Addressed on page number
<b>Administrative information</b>			
Title	1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	P1/L1-3
Trial registration	2a	Trial identifier and registry name. If not yet registered, name of intended registry	P4/L7-8; P8/L14-16; P19/L4-7
	2b	All items from the World Health Organization Trial Registration Data Set	P8/L3-L17
Protocol version	3	Date and version identifier	P8/L11
Funding	4	Sources and types of financial, material, and other support	P19/L18-21
Roles and responsibilities	5a	Names, affiliations, and roles of protocol contributors	P1/L5-16, P19/L23-P20/L2
	5b	Name and contact information for the trial sponsor	P19/L18-21
	5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities	P19/L18-21
	5d	Composition, roles, and responsibilities of the coordinating centre, steering committee, endpoint adjudication committee, data management team, and other individuals or groups overseeing the trial, if applicable (see Item 21a for data monitoring committee)	N/A

**Introduction**

1				
2				
3	Background and	6a	Description of research question and justification for undertaking the trial, including summary of relevant	P6-7
4	rationale		studies (published and unpublished) examining benefits and harms for each intervention	
5				
6		6b	Explanation for choice of comparators	P7/L8-22;
7				P10/L12-22
8	Objectives	7	Specific objectives or hypotheses	P7/L23-25
9				
10	Trial design	8	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group),	
11			allocation ratio, and framework (eg, superiority, equivalence, noninferiority, exploratory)	P8/L3-9
12				
13				
14	<b>Methods: Participants, interventions, and outcomes</b>			
15				
16	Study setting	9	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will	P8/L7-9
17			be collected. Reference to where list of study sites can be obtained	
18				
19	Eligibility criteria	10	Inclusion and exclusion criteria for participants. If applicable, eligibility criteria for study centres and	P8/L21-P9/L6
20			individuals who will perform the interventions (eg, surgeons, psychotherapists)	
21				
22	Interventions	11a	Interventions for each group with sufficient detail to allow replication, including how and when they will be	P10/L12-22
23			administered	
24				
25		11b	Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose	P10/L11-24,
26			change in response to harms, participant request, or improving/worsening disease)	P11/L1-2
27				
28		11c	Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence	P10-11
29			(eg, drug tablet return, laboratory tests)	
30				
31		11d	Relevant concomitant care and interventions that are permitted or prohibited during the trial	N/A
32				
33	Outcomes	12	Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood	P12/L20-P14/L8
34			pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg,	
35			median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen	
36			efficacy and harm outcomes is strongly recommended	
37				
38	Participant timeline	13	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for	Figure 1
39			participants. A schematic diagram is highly recommended (see Figure)	
40				
41				
42				
43				
44				
45				
46				
47				

1				
2				
3	Sample size	14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	P14/L24-P15/L7
4				
5	Recruitment	15	Strategies for achieving adequate participant enrolment to reach target sample size	P9/L7-17
6				
7				

### 8 **Methods: Assignment of interventions (for controlled trials)**

#### 9 Allocation:

10				
11				
12	Sequence generation	16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	P9/L18-24
13				
14				
15				
16				
17	Allocation concealment mechanism	16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	P9/L18-P10/L9
18				
19				
20				
21	Implementation	16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	P9/L21-24
22				
23				
24	Blinding (masking)	17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	P10/L1-4
25				
26				
27		17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	P10/L5-9
28				
29				
30				

### 31 **Methods: Data collection, management, and analysis**

32				
33	Data collection methods	18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known. Reference to where data collection forms can be found, if not in the protocol	P12/L6-P15/L20
34				
35				
36				
37				
38		18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols	P12/L25-P13/L3
39				
40				
41				
42				
43				
44				
45				
46				
47				

1				
2				
3	Data management	19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	P14/L9-15
4				
5				
6				
7	Statistical methods	20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	P14/L23-P15/L20
8				
9				
10		20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	P15/L9-16
11				
12		20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any statistical methods to handle missing data (eg, multiple imputation)	P11/L7-9; P12/L25-P13/L3
13				
14				
15				
16				
17	<b>Methods: Monitoring</b>			
18	Data monitoring	21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent from the sponsor and competing interests; and reference to where further details about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not needed	P14/L9-22
19				
20				
21				
22				
23				
24		21b	Description of any interim analyses and stopping guidelines, including who will have access to these interim results and make the final decision to terminate the trial	P14/L19-22
25				
26				
27	Harms	22	Plans for collecting, assessing, reporting, and managing solicited and spontaneously reported adverse events and other unintended effects of trial interventions or trial conduct	P13/L16-P14/L8
28				
29				
30	Auditing	23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent from investigators and the sponsor	P14/L16-19
31				
32				
33	<b>Ethics and dissemination</b>			
34				
35	Research ethics approval	24	Plans for seeking research ethics committee/institutional review board (REC/IRB) approval	P8/L11-14
36				
37				
38	Protocol amendments	25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes, analyses) to relevant parties (eg, investigators, REC/IRBs, trial participants, trial registries, journals, regulators)	P8/L11-14
39				
40				
41				
42				
43				
44				
45				
46				
47				

1				
2				
3	Consent or assent	26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	P8/L16-17 P19/L3-7
4				
5				
6		26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	N/A
7				
8	Confidentiality	27	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	P14/L10-11
9				
10				
11	Declaration of interests	28	Financial and other competing interests for principal investigators for the overall trial and each study site	P19/L12-16
12				
13				
14	Access to data	29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	P14/L11-14; P20/L12
15				
16				
17	Ancillary and post-trial care	30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	P14/L6-8
18				
19				
20	Dissemination policy	31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	P4/L4-5; P19/L9-10
21				
22				
23				
24				
25		31b	Authorship eligibility guidelines and any intended use of professional writers	P19/L23-P20/L2
26				
27		31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	P20/L12
28				
29	<b>Appendices</b>			
30				
31	Informed consent materials	32	Model consent form and other related documentation given to participants and authorised surrogates	N/A
32				
33				
34	Biological specimens	33	Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies, if applicable	N/A
35				
36				

37 \*It is strongly recommended that this checklist be read in conjunction with the SPIRIT 2013 Explanation & Elaboration for important clarification on the items.  
 38 Amendments to the protocol should be tracked and dated. The SPIRIT checklist is copyrighted by the SPIRIT Group under the Creative Commons  
 39 "[Attribution-NonCommercial-NoDerivs 3.0 Unported](https://creativecommons.org/licenses/by-nc-nd/3.0/)" license.  
 40