

**eTable 2.** Odds ratios and confidence intervals of being lost to follow-up – injury factors: Results from the multivariable model for each post-injury year<sup>a</sup>

		Post-injury year							
		1	5	10	15	20	25	30	35
Injury cohort	1972-1985	<b>2.30</b> (2.08, 2.55)	<b>2.11</b> (1.90, 2.35)	<b>2.23</b> (1.91, 2.61)	0.88 (0.05, 14.93)	<b>1.56</b> (1.37, 1.77)	<b>1.40</b> (1.16, 1.69)		
	1986-2000	<b>1.33</b> (1.21, 1.47)	<b>1.64</b> (1.48, 1.82)	<b>1.83</b> (1.57, 2.13)	0.49 (0.03, 8.38)	Ref	Ref		
	2001-2013	Ref	Ref	Ref	Ref	Ref	Ref		
Etiology	Vehicular					Ref			
	Violence	<b>1.12</b> (1.02, 1.23)	1.09 (0.99, 1.21)	1.09 (0.97, 1.23)	1.07 (0.93, 1.23)	1.05 (0.89, 1.24)	1.14 (0.93, 1.39)	0.92 (0.70, 1.20)	0.79 (0.51, 1.25)
	Sports	0.95 (0.85, 1.06)	0.93 (0.83, 1.04)	0.91 (0.80, 1.04)	0.90 (0.78, 1.05)	0.93 (0.79, 1.10)	0.98 (0.80, 1.18)	0.90 (0.72, 1.13)	1.03 (0.71, 1.49)
	Falls	1.02 (0.93, 1.11)	1.11 (1.00, 1.22)	<b>1.14</b> (1.02, 1.28)	1.04 (0.91, 1.2)	1.03 (0.88, 1.22)	0.97 (0.80, 1.19)	1.00 (0.78, 1.28)	1.14 (0.77, 1.71)
	Other/Unknown	1.07 (0.95, 1.21)	1.13 (1.00, 1.29)	1.00 (0.86, 1.17)	1.04 (0.86, 1.24)	0.98 (0.79, 1.21)	0.95 (0.73, 1.24)	0.87 (0.63, 1.19)	1.40 (0.84, 2.35)
Neurological category	Ventilator dependent	0.74 (0.58, 0.94)	<b>0.75</b> (0.59, 0.97)	0.81 (0.60, 1.10)	<b>0.50</b> (0.33, 0.75)	0.69 (0.39, 1.21)	0.48 (0.24, 0.94)	0.65 (0.26, 1.67)	2.77 (0.38, 20.36)
	C1-C4 AIS ABC	0.86 (0.76, 0.97)	<b>0.62</b> (0.54, 0.71)	<b>0.75</b> (0.64, 0.87)	<b>0.63</b> (0.52, 0.76)	<b>0.73</b> (0.58, 0.91)	<b>0.58</b> (0.45, 0.76)	<b>0.54</b> (0.39, 0.76)	0.72 (0.40, 1.27)
	C5-C8 AIS ABC	<b>0.89</b> (0.80, 0.98)	<b>0.69</b> (0.63, 0.77)	<b>0.67</b> (0.60, 0.76)	<b>0.75</b> (0.65, 0.86)	<b>0.69</b> (0.59, 0.81)	<b>0.58</b> (0.48, 0.69)	<b>0.72</b> (0.57, 0.90)	0.94 (0.66, 1.34)
	T1-S3 AIS ABC	0.98 (0.89, 1.07)	<b>0.77</b> (0.70, 0.84)	<b>0.79</b> (0.71, 0.87)	0.90 (0.79, 1.01)	<b>0.82</b> (0.71, 0.94)	<b>0.66</b> (0.56, 0.77)	<b>0.75</b> (0.62, 0.92)	0.91 (0.67, 1.26)
	AIS D					Ref			
	Unknown	1.04 (0.89, 1.21)	0.86 (0.72, 1.02)	0.96 (0.77, 1.20)	1.21 (0.91, 1.62)	0.77 (0.54, 1.11)	0.60 (0.36, 0.87)	1.17 (0.68, 2.00)	1.44 (0.37, 5.65)

(Continued)

**eTable 2.** Odds ratios and confidence intervals of being lost to follow-up – injury factors: Results from the multivariable model for each post-injury year<sup>a</sup> (CONT.)

Bladder management	Catheter-free/Normal	Post-injury year						
		1	5	10	15	20	25	30
ICP	0.90 (0.83, 0.98)	0.99 (0.89, 1.10)	<b>0.82</b> (0.71, <b>0.93</b> )	0.86 (0.73, 1.02)	0.93 (0.76, 1.13)	1.01 (0.79, 1.29)	0.89 (0.66, 1.21)	0.82 (0.49, 1.36)
Indwelling	0.94 (0.85, 1.04)	<u>0.87</u> (0.78, <u>0.98</u> )	0.88 (0.76, 1.01)	1.07 (0.90, 1.26)	0.85 (0.70, 1.05)	1.00 (0.800, 1.27)	0.87 (0.66, 1.15)	<u>0.57</u> (0.36, <u>0.9</u> )
Other/Unknown	1.03 (0.86, 1.22)	1.12 (0.94, 1.34)	1.20 (0.97, 1.49)	<b>1.43</b> (1.10, <b>1.85</b> )	1.35 (0.98, 1.86)	1.46 (0.93, 2.23)	0.76 (0.44, 1.32)	0.54 (0.25, 1.20)

Note: Odds ratios >1 indicate more likely to be lost to follow-up. Odds ratios with underline indicate .01 ≤ p < .05; bold font indicates p < .01.

AIS = American Spinal Injury Association Impairment Scale; C = cervical; Gov = government; ICP = intermittent catheterization program; T = thoracic; S = sacral.

<sup>a</sup>Controlling for Spinal Cord Injury Model Systems centers, listed variables, and demographic factors.