

**Table S2.** Baseline characteristics of those included and excluded from mortality analysis in CRIC and AASK.

Baseline characteristics at first CRIC visit of those included and excluded from mortality analysis in CRIC.

Characteristics during CKD	Included (N=770)	Excluded (N=101)	P-value
Mean age (yrs) ±SD	56.5 ± 11.0	51.3 ± 14.1	<0.001
Men N (%)	455 (59.1)	67 (66.3)	0.16
Black N (%)	418 (54.3)	50 (49.5)	0.37
Annual household income N (%)			0.60
≤\$20,000	337 (43.8)	52 (51.5)	
\$20,001-\$50,000	179 (23.2)	21 (20.8)	
≥\$50,001	135 (17.5)	17 (16.7)	
Decline to answer	119 (15.5)	11 (10.9)	
Smoked > 100 cigarettes over lifetime N (%)	448 (58.2)	59 (58.4)	0.96
Diabetes N (%)	518 (67.3)	67 (66.3)	0.85
Hypertension N (%)	735 (95.5)	92 (91.1)	0.06
Prevalent CVD N (%)	322 (41.8)	35 (34.7)	0.17
Mean serum albumin (g/dL) ± SD	3.7 ± 0.5	3.6 ± 0.6	0.69
COPD N (%)	20 (2.6%)	8 (8.0)	0.005
Cancer N (%)	82 (10.8)	6 (5.9)	0.13
Systolic BP (mm Hg)	139.4 ± 24.1	135.4 ± 23.5	0.09
Diastolic BP (mm Hg)	72.9 ± 13.9	76.4 ± 14.5	0.14
Median hemoglobin A1C (%)	6.7 [5.8-8.1]	6.4[5.7-8.3]	0.36

Baseline characteristics of those included and excluded from mortality analysis in AASK.

Characteristics during CKD	Included (N=333)	Excluded (N=64)	P-value
Mean age (yrs) ±SD	51.7 ± 11.3	49.1 ± 11.7	0.10
Men N (%)	199 (59.8)	38 (59.4)	0.95
Annual household income determined at baseline AASK enrollment N (%)			<0.001
<\$15,000	165 (49.6)	25 (39.1)	
\$15,000-39,999	117 (35.1)	19 (29.7)	
>\$40,000	51 (15.3)	20 (31.3)	
Current or past smoker determined at time of baseline AASK enrollment	179 (53.8)	38 (59.4)	0.41
Prevalent CVD N (%)	168 (50.5)	32 (50.0)	0.95
Mean serum albumin (g/dL) ±SD	4.1 ± 0.4	4.2 ± 0.4	0.18
Systolic BP (mm Hg)	147.1 ± 24.0	150.4 ± 24.1	0.33
Diastolic BP (mm Hg)	91.5 ± 16.3	97.5 ± 14.1	<0.001