

Hsp90 inhibition as a means to inhibit activation of the NLRP3 inflammasome

Niina Piippo¹, Eveliina Korhonen¹, Maria Hytti¹, Heli Skottman², Kati Kinnunen³, Natasha Josifovska⁴, Goran Petrovski^{4,5}, Kai Kaarniranta^{3,6}, Anu Kauppinen^{1,*}

¹School of Pharmacy, University of Eastern Finland, Kuopio, 70211, Finland

²Faculty of Medicine and Life Sciences, BioMediTech, University of Tampere, Tampere, 33014, Finland


³Department of Ophthalmology, Kuopio University Hospital, Kuopio, 70211, Finland

⁴Stem Cells and Eye Research Laboratory, Department of Ophthalmology, Faculty of Medicine, Albert Szent-Györgyi Clinical Center, University of Szeged, Szeged, Hungary

⁵Center for Eye Research, Department of Ophthalmology, Oslo University Hospital, University of Oslo, Oslo, Norway

⁶Department of Ophthalmology, Institute of Clinical Medicine, University of Eastern Finland, Kuopio, 70211, Finland

***Corresponding author:** Anu Kauppinen, School of Pharmacy, Faculty of Health Sciences, University of Eastern Finland, P.O. Box 1627, FIN-70211 Kuopio, Finland, Fax: +358 17 162048, Tel.: +358 40 3553216, e-mail: anu.kauppinen@uef.fi



Supplementary Figure 1. Scan of the complete blot represented in Figure 5. NLRP3 and α -tubulin, as an internal control, were detected from the same blot. Boxes indicate the cropped lanes shown in Figure 5.