



SUPPLEMENTAL FIGURE 1. ROC space plot for caregiver-reported stool consistency with and without the modified Bristol Stool Form Scale for children.

SUPPLEMENTAL TABLE 1

K-by-K matrix of caregiver-reported “loose or watery” stool versus laboratory technician–graded stool consistency using modified Bristol Stool Form Scale for children

Caregiver-reported stool consistency (index test 1)	First laboratory technician–graded stool consistency (reference standard)					Total (index test)
	Type 1: pellets	Type 2: lumpy	Type 3: snake	Type 4: loose	Type 5: watery	
Not loose or watery	290	479	404	332	35	1,540
Loose or watery	0	11	8	35	33	87
Total (reference standard)	290	490	412	367	68	1,628

SUPPLEMENTAL TABLE 2

K-by-K matrix of caregiver-reported stool consistency using modified Bristol Stool Form Scale for children (mBSFS-C) vs. laboratory technician–graded stool consistency using mBSFS-C

Caregiver-reported stool consistency using mBSFS-C (index test 2)	Laboratory technician–graded stool consistency (reference standard)					Total (index test)
	Type 1: pellets	Type 2: lumpy	Type 3: snake	Type 4: loose	Type 5: watery	
Type 1: pellets	136	68	10	57	2	273
Type 2: lumpy	98	237	99	36	5	475
Type 3: snake	35	76	267	36	1	415
Type 4: loose	10	51	26	117	16	220
Type 5: watery	11	58	10	121	44	244
Total (reference standard)	290	489	412	367	68	1,627

Bold indicates the total number of grades that agree in each category.