

Table S1

Variables	Amerindian community-based urban groups			Mestizos	<i>p value</i>	
	Low	Medium	High	Mestizo	Amer-indians (a)	Amer-indians high vs. Mestizos
N subjects	24	28	30	29	0.710 (a)	0.890 (a)
Age, Mean [min, max]	28.3 [12-46]	30.2 [18-42]	31.2 [17-45]	26.0 [17-53]	0.500 (b)	0.025 (b)
Educational level % (n/N); [CI 95%]					0.001*	
No studies	64.7 (11/17); [41-86]	16.0 (4/25); [5.3-37]	16.0 (4/25); [5.3-37]	4.1 (1/24); [0.2-23]	(a)	0.350 (a)
Finished elementary school	23.5 (4/17); [7.8-50]	40.0 (10/25); [22-61]	32.0 (8/25); [16-54]	12.5 (3/24); [3.3-33]	0.570 (a)	0.170 (a)
Finished high school	11.8 (2/17); [2-39]	64.8 (11/25); [37-85]	52.0 (13/25); [32-72]	83.3 (20/24); [62-95]	0.020 (a)	0.220 (a)
	0.0	3.6	10.0	34.5		
Hormonal contraceptive** use currently % (n/N); [CI 95%]					0.220 (a)	0.050 (a)
Parity, mean [IQR]	4.3 [0-11]	4.0 [0-13]	2.9 [0-12]	1.8 [0-8]	0.340 (b)	0.090 (b)
Breastfeeding currently % (n/N); [CI 95%]	62.5 (15/24); [41-80]	62.9 (17/28); [41-78]	36.7 (11/30); [21-56]	70.0 (20/29); [49-84]	0.094 (b)	0.026 (b)
Sexual partner history number, Median [min- max]	2.0 [1.0-10]	2.0 [1.0-6.0]	3.0 [1.0-15]	2.0 [1.0-25]	0.750 (c)	0.100 (c)
Number of sexual partners in last 60 days % (n/N); [CI 95%]						
None	25.0 (6/24); [11-47]	21.4 (6/28); [9-42]	30.0 (9/30); [15-50]	13.8 (4/29); [4.5-33]	0.750 (a)	0.260 (a)
1	75.0 (18/24); [53-89]	78.6 (22/28); [59-91]	70.0 (21/30); [50-85]	82.8 (24/29); [64-94]		
Sexual intercourse weekly frequency % (n/N); [CI 95%]						
≤ 1 time	95.8 (23/24); [77-100]	75.0 (21/28); [55-89]	70.0 (21/30); [50-85]	41.4 (12/29); [24-61]	0.060 (a)	0.050 (a)
≥ 2 times	4.2 (1/24); [0.2-23]	25.0 (7/28); [11-45]	30.0 (9/30); [15-50]	58.6 (17/29); [64-94]		
Sexual contact with mestizo % (n/N); [CI 95%]	8.3 (2/24); [15-29]	10.7 (3/28); [28-29]	46.7 (14/30); [29-65]	100.0 (29/29); [85-100]	6x10 ⁻⁴ * (a)	2x10 ⁻⁵ * (a)
Currently smoking*** % (n/N); [CI 95%]	0.0 (0/24); [0-17]	0.0 (0/28); [0-15]	10.0 (3/30); [2.6-27]	13.8 (4/29); [4.5-33]	0.034 (a)	0.699 (a)

(a) χ^2 test or Fisher's Exact Test

(b) T-test and ANOVA for 2 or more than two groups

(c) Kruskal-Wallis

*Significant differences reached ($p < 0.05$) after Holm correction for multiple comparisons.

**For non hormonal contraceptive use, for Amerindians, Low group = 0 cases, Medium group = 1 sterilization, High group = 4 sterilizations and 2 condom use cases; for Mestizo = 3 condom use cases.

***Smoking frequency from 1 to 10 cigarettes daily during 1 or more years