

S1 Table Literature data overview on wild and cultivated plants associated with *Hyalesthes obsoletus* feeding, aggregation and/or development resulting in polyphagous perception of this vector.

Country	Plant	Note	Reference
Russia	<i>Convolvulus arvensis</i>	nymphs and adults	Sukhov & Razvyazkina 1955
	<i>Lepidium draba</i>		
	<i>Solanum tuberosum</i> (potato)	adults	Imo <i>et al.</i> 2013
Turkey	<i>Cistus</i> sp.	adults	Hoch & Remane 1985
	<i>Crataegus</i> sp.		
	<i>Myrtus</i> sp.		
	<i>Paliurus</i> sp.		
	<i>Quercus coccifera</i>		
	<i>Quercus ilex</i>		
	<i>Rubus</i> sp.		
	<i>Sesamum indicum</i>		
	<i>Vitex</i> sp.		
	<i>Capsicum annuum</i> (paprika)		
	<i>Convolvulus arvensis</i>	overwintering hosts for the	Güclü & Ozbek 1988
	<i>Crambe orientalis</i>	nymphs; adults aggregate	
	<i>Falcaria vulgaris</i>	mainly on <i>Medicago sativa</i>	
Israel	<i>Geranium tuberosum</i>	but also found on diverse	
	<i>Isatis glauca</i>	crops (potato, sugar beet,	
	<i>Lepidium draba</i>	sainfoin, red clover, bean	
	<i>Melilotus officinalis</i>	and rarely sunflower and	
	<i>Onosma armeneum</i>	squash)	
	<i>Rumex crispus</i>		
	<i>Taraxacum</i> sp.		
	<i>Medicago sativa</i> (alfalfa)		
	<i>Onobrichis sativa</i> (sainfoin)		
	<i>Tamarix</i> sp.	adults	Hoch & Remane 1985
	<i>Gossypium</i> sp. (cotton)		
	<i>Solanum tuberosum</i> (potato)		
	<i>Zea mays</i> (corn)		
	<i>Amaranthus retroflexus</i>	adults;	Sharon <i>et al.</i> 2005
	<i>Convolvulus arvensis</i>	nymphs were found on the	
	<i>Myrtus communis</i>	<i>V. agnus-castus</i> roots	
	<i>Olea europaea</i> ,		

	<i>Vitex agnus-castus</i>		
	<i>Vitis vinifera</i> (grapevine)		
Greece	<i>Acer campestre</i>	adults;	Hoch & Remane 1985
	<i>Arbutus</i> sp.	the most common host	
	<i>Arundo</i> sp.	plant is <i>Vitex</i> sp. in	
	<i>Artemisia</i> sp.	coastline habitats	
	<i>Carpinus betulus</i>		
	<i>Cistus</i> sp.		
	<i>Convolvulus</i> sp.		
	<i>Crataegus</i> sp.		
	<i>Cynodon</i> sp.		
	<i>Erica arborea</i>		
	<i>Euphorbia dendroides</i>		
	<i>Galium</i> sp.		
	<i>Hipparhenia hirta</i>		
	<i>Imperata cylindrica</i>		
	<i>Juncus</i> sp.		
	<i>Ligustrum vulgare</i>		
	<i>Myrtus</i> sp.		
	<i>Oreganum</i> sp.		
	<i>Quercus coccifera</i>		
	<i>Quercus cerris</i>		
	<i>Quercus ilex</i>		
	<i>Quercus pubescens</i>		
	<i>Paliurus</i> sp.		
	<i>Phragmites</i> sp.		
	<i>Platanus</i> sp.		
	<i>Populus</i> sp.		
	<i>Prunus spinosa</i>		
	<i>Pteridium</i> sp.		
	<i>Rubus</i> sp.		
	<i>Salix</i> sp.		
	<i>Tamarix</i> sp.		
	<i>Ulmus</i> sp.		
	<i>Vitex</i> sp.		
	<i>Capsicum annuum</i> (paprika)		
	<i>Gossypium</i> sp. (cotton)		
	<i>Sesamum indicum</i> (sesame)		

	<i>Solanum tuberosum</i> (potato) <i>Zea mays</i> (corn)		
Cyprus and Greek islands (Crete, Lesbos, Ikaria, Rhodes)	<i>Arbutus</i> sp.	adults;	Hoch & Remane 1985
	<i>Arundo</i> sp.	the most common host	
	<i>Erica arborea</i>	plant is <i>Vitex</i> sp. in	
	<i>Euphorbia dendroides</i>	coastline habitats	
	<i>Imperata cylindrica</i>		
	<i>Juncus</i> sp.		
	<i>Phragmites</i> sp.		
	<i>Platanus</i> sp.		
	<i>Pteridium</i> sp.		
	<i>Tamarix</i> sp		
Romania	<i>Quercus pubescens</i>		
	<i>Vitex</i> sp.		
	<i>Solanum tuberosum</i> (potato)	adults	Imo <i>et al.</i> 2013
	<i>Convolvulus arvensis</i>	adults collected using	Atanasova <i>et al.</i> 2015
	<i>Urtica dioica</i>	sweep net and mouth aspirator	
	<i>Convolvulus arvensis</i>	adults collected using	Kosovac <i>et al.</i> 2016
	<i>Urtica dioica</i>	sweep net and mouth aspirator	
	<i>Vitex agnus-castus</i>		
	<i>Convolvulus arvensis</i>		Aleksić <i>et al.</i> 1967
	various vineyard weeds	first data of <i>Crepis foetida</i> as a member of weed community where adults were non selectively captured by sweeping net	Cvrković Tatjana, PhD thesis, 2010
Serbia	<i>Convolvulus arvensis</i>	adults collected using	Kosovac <i>et al.</i> 2013
	<i>Urtica dioica</i>	sweep net and mouth aspirator	
	<i>Crepis foetida</i>		
Bosnia and Herzegovina	<i>Convolvulus arvensis</i>	adults collected using	Đurić <i>et al.</i> 2017
	<i>Vitex agnus-castus</i>	sweep net and mouth aspirator	
Croatia	<i>Vitex agnus-castus</i>	very often found on <i>V. agnus-castus</i>	Horváth 1891
Slovenia	<i>Convolvulus arvensis</i>	adults collected using	Mehle <i>et al.</i> 2011
	<i>Urtica dioica</i>	yellow sticky traps or by sweep net sampling	

	<i>Convolvulus arvensis</i>	adults collected on a single plant species with an exhauster or with sweep nets in monoculture plant stands	Imo <i>et al.</i> 2013
Hungary	<i>Urtica dioica</i>	adults collected with D-Vac aspirators	Palermo <i>et al.</i> 2004
Austria	<i>Urtica dioica</i> <i>Convolvulus arvensis</i>	association with <i>U. dioica</i> is dominant	Aryan <i>et al.</i> 2014
Switzerland	<i>Urtica dioica</i> <i>Convolvulus arvensis</i>	adults collected with mouth aspirators mainly on <i>U. dioica</i> ; nymphs present on the root in the field and confirmed as laboratory plant for nymphs development	Kessler <i>et al.</i> 2011
	<i>Calystegia sepium</i>	adults were captured on yellow sticky traps positioned above the specific plant	
	<i>Lepidium draba</i>		
	<i>Cardaria draba</i>		
	<i>Clematis vitalba</i>		
	<i>Polygonum aviculare</i>	adults were taken with a D-vac vacuum insect collector	
	<i>Calystegia sepium</i>		
	<i>Taraxatum officinale</i>		
	<i>Plantago lanceolata</i>		
	<i>Calystegia sepium</i>		
Germany	<i>Convolvulus arvensis</i>	on limestone	Hoch & Remane 1985
	<i>Festuca</i> sp.		
	<i>Allium</i> sp.		
	<i>Convolvulus arvensis</i>		Maixner 1994
	<i>Solanum nigrum</i>		
	<i>Vitis vinifera</i> (grapevine)		
	<i>Ranunculus bulbosus</i>		Weber & Maixner 1998
	<i>Convolvulus arvensis</i>		Langer & Maixner 2004
	<i>Urtica dioica</i>		
	<i>Calystegia sepium</i>		
France	<i>Calystegia sepium</i>	adults collected by sweeping net	Sforza <i>et al.</i> 1999
	<i>Cardaria draba</i>		
	<i>Convolvulus arvensis</i>		

	<i>Galium verum</i>		
	<i>Lavandula angustifolia</i>		
	<i>Linaria striata</i>		
	<i>Plantago cynopsis</i>		
	<i>Satureia montana</i>		
	<i>Convolvulus arvensis</i>	adults collected with D-Vac	Bressan <i>et al.</i> 2008
	<i>Urtica dioica</i>	aspirators	
	<i>Salvia sclarea</i>	adults collected with D-Vac aspirators and yellow sticky traps; nymphs collected on uprooted individual plants	Chuche <i>et al.</i> 2018
	<i>Convolvulus arvensis</i>	adults collected with D-Vac	
	<i>Urtica dioica</i>	aspirators and yellow sticky traps	
	<i>Lavandula angustifolia</i> (lavander)	adults were captured on yellow sticky traps in fields	Yvin <i>et al.</i> 2017
	<i>Lavandula x intermedia</i> (lavandin)	planted with diverse lavender and lavandin cultivars	
	<i>Salvia sclarea</i>	nymphs counted on uprooted plants; adults collected with emergence traps	Hossard <i>et al.</i> 2018
	<i>Convolvulus arvensis</i>	adults collected with	
	<i>Lavandula hybrida</i>	emergence traps	
Spain	<i>Trifolium pratense</i>		Hoch & Remane 1985
	<i>Convolvulus arvensis</i>		Sabate <i>et al.</i> 2014
Italy	<i>Urtica dioica</i>	nymphs found on <i>U. dioica</i>	Lessio <i>et al.</i> 2007
	<i>Convolvulus arvensis</i>	roots	
Italy	<i>Dactylis</i> sp.		Hoch & Remane 1985
(Sicily)	<i>Juncus</i> sp.		
	fam. <i>Lamiaceae</i>		
	<i>Ulmus</i> sp.		
	<i>Urtica</i> sp.		
Morocco	<i>Clematis</i> sp.		Hoch & Remane 1985
	<i>Pistacia</i> sp.		
	<i>Tamarix</i> sp		
	<i>Quercus coccifera</i>		

References

- Atanasova B, Jakovljević M, Spasov D, Jović J, Mitrović M, Toševski I, Cvrković T (2015) The molecular epidemiology of bois noir grapevine yellows caused by '*Candidatus Phytoplasma solani*' in the Republic of Macedonia. *European Journal of Plant Pathology*, **142**, 759-770.
- Aleksić Ž, Šutić D, Aleksić D (1967) Transmission intensity of stolbur virus by means of *Hyalesthes obsoletus* Sign. on some host plants. *Plant Protection*, **93–95**, 67–73.
- Aryan A, Brader G, Mörtel J, Pastar M, Riedle-Bauer M (2014) An abundant '*Candidatus Phytoplasma solani*' Stolbur tuf b phytoplasma strain is associated with grapevine, stinging nettle and *Hyalesthes obsoletus*. *European Journal of Plant Pathology*, **140**, 213-227.
- Bressan A, Sémétey O, Nusillard B, Clair D, Boudon-Padieu E (2008) Insect vectors (Hemiptera: Cixiidae) and pathogens associated with the disease syndrome “basses richesses” of sugar beet in France. *Plant Disease*, **92**(1), 113-119.
- Chuche J, Danet JL, Rivoal JB, Arricau-Bouvery N, Thiéry D (2018) Minor cultures as hosts for vectors of extensive crop diseases: Does *Salvia sclarea* act as a pathogen and vector reservoir for lavender decline? *Journal of Pest Science*, **91**(1), 145-155.
- Cvrković T (2010) Diverzitet faune cikada u vinogradima Srbije i njihova uloga u prenošenju Bois Noir fitoplazme. PhD thesis, Faculty of Agriculture, University of Belgrade, 1-101.
- Đurić Z, Hrnčić S, Delić D (2017) Morphological and molecular identification of *Hyalesthes obsoletus* Signoret (Auchenorrhyncha: Cixiidae) in Herzegovina vineyard. *Mitteilungen Klosterneuburg* **67**, 177-181
- Güclü S, Ozbek H (1988) Some studies on the biology of *Hyalesthes obsoletus* Signoret (Homoptera: Cixidae) in the conditions of Erzurum. *Turkiye-Entomoloji-Dergisi*, **12**, 103-111.
- Hoch H, Remane R (1985) Evolution und Speziation der Zikaden-Gattung *Hyalesthes* Signoret, 1865 (Homoptera Auchenorrhyncha Fulgoroidea Cixiidae). *Marburger Entomologische Publikationen*, **2**, 1-427.
- Horváth G (1891) Hemipterološki izlet u Primorje i Plitvička jezera. *Glasnik Hrvatskoga Naravoslovnoga Drustva (Societas historico-naturalis Croatica)*, VI (1-5), 29-49.
- Hossard L, Guimier S, Vinatier F, Barbier JM, Delmotte S, Fontaine M, Rivoal JB (2018) Source of *Hyalesthes obsoletus* Signoret planthopper (Hemiptera: Cixiidae) in southern France and potential effects of landscape. *Bulletin of Entomological Research*, **108**, 213-222.

- Imo M, Maixner M, Johannesen J (2013) Sympatric diversification vs. immigration: deciphering host-plant specialization in a polyphagous insect, the stolbur phytoplasma vector *Hyalesthes obsoletus* (Cixiidae). *Molecular Ecology*, **22**, 2188-2203.
- Kessler S, Schaefer S, Delabays N, Turlings TC, Trivellone V, Kehrli P (2011) Host plant preferences of *Hyalesthes obsoletus*, the vector of the grapevine yellows disease ‘bois noir’, in Switzerland. *Entomologia Experimentalis et Applicata*, **139**, 60-67.
- Kosovac A, Johannesen J, Krstić O, Mitrović M, Cvrković T, Maixner M, Toševski I, Jović J (2013) Microsatellite and mtDNA evidence of genetic differentiation in *Hyalesthes obsoletus* populations associated with a new major host, stinking hawk’s-beard (*Crepis foetida*), in southeast Europe. *Proceedings of the 3rd European Bois Noir Workshop*, Barcelona, 18-19.
- Kosovac A, Radonjić S, Hrnčić S, Krstić O, Toševski I, Jović J (2016) Molecular tracing of the transmission routes of bois noir in Mediterranean vineyards of Montenegro and experimental evidence for the epidemiological role of *Vitex agnus-castus* (Lamiaceae) and associated *Hyalesthes obsoletus* (Cixiidae). *Plant Pathology*, **65**, 285-298.
- Langer M, Maixner M (2004) Molecular characterisation of grapevine yellows associated phytoplasmas of the stolbur group based on RFLP analysis of non-ribosomal DNA. *Vitis*, **43**, 191–199.
- Lessio F, Tedeschi R, Alma A (2007) Population dynamics, host plants and infection rate with Stolbur phytoplasma of *Hyalesthes obsoletus* Signoret in north-western Italy. *Journal of Plant Pathology*, **89**, 97-102.
- Maixner M (1994) Research note: Transmission of German grapevine yellows (Vergilbungskrankheit) by the planthopper *Hyalesthes obsoletus* (Auchenorrhyncha: Cixiidae). *Vitis*, **33**, 103-104.
- Palermo S, Elekes M, Botti S, Ember I, Alma A, Orosz A, Bertaccini A, Kölber M (2004). Presence of stolbur phytoplasma in Cixiidae in Hungarian vineyards. *Vitis*, **43**(4), 201-203.
- Sabaté J, Laviña A, Batlle A (2014) Incidence of Bois Noir phytoplasma in different viticulture regions of Spain and Stolbur isolates distribution in plants and vectors. *European Journal of Plant Pathology*, **139**, 185-193.
- Sforza R, Bourgoin T, Wilson SW, Boudon-Padieu E (1999) Field observations, laboratory rearing and descriptions of immatures of the planthopper *Hyalesthes obsoletus* (Hemiptera: Cixiidae). *European Journal of Entomology*, **96**, 409–418.

- Sharon R, Soroker V, Wesley SD, Zahavi T, Harari A, Weintraub PG (2005) *Vitex agnus-castus* is a preferred host plant for *Hyalesthes obsoletus*. *Journal of Chemical Ecology*, **31**, 1051-1063.
- Suhov KS, Vovk AM (1949) Stolbur of solanaceous plants. USSR Academy of Sciences Publishers. 7-11.
- Weber A, Maixner M (1998) Survey of populations of the planthopper *Hyalesthes obsoletus* Sign.(Auchenorrhyncha, Cixiidae) for infection with the phytoplasma causing grapevine yellows in Germany. *Journal of Applied Entomology*, **122**, 375-381.
- Yvin C, Chaisse E, Fontaine M, Rivoal JB, Verdin E, Nicolè F, Jourdan P, Savarit P, Gallois P (2017) Dépérissage de la lavande et du lavandin: mise en oeuvre d'un programme de recherches appliquées afin d'apporter des solutions de lutte aux producteurs. *Innovations Agronomiques*, **55**, 107-120.