

Perlecan, a heparan sulfate proteoglycan, regulates systemic metabolism with dynamic changes in adipose tissue and skeletal muscle

Yuri Yamashita^{1,2}, Satoshi Nakada³, Toshinori Yoshihara⁴, Takeshi Nara⁵, Norihiko Furuya², Takashi Miida⁶, Nobutaka Hattori², and Eri Arikawa-Hirasawa^{1,2,*}

¹Aging Biology in Health and Disease, ²Department of Neurology, and ⁶Department of Clinical Laboratory medicine, Juntendo University Graduate School of Medicine, Tokyo 113-8421, Japan. ³Japanese Center for Research on Women in Sport and ⁴Department of Exercise Physiology, Juntendo University Graduate School of Health and Sports Science, Chiba 270-1695, Japan. ⁵Faculty of Pharmacy, Iwaki Meisei University, Fukushima 970-8551, Japan.

*Correspondence and requests for materials should be addressed to E.AH.

Eri Arikawa-Hirasawa: at: Juntendo University Faculty of Medicine, Research Institute for Disease of Old Age, Building 10, Room 606, 2-1-1 Hongo, Bunkyo-ku, Tokyo 113-8421, Japan (email: ehirasaw@juntendo.ac.jp)

SUPPLEMENTARY INFORMATION

Supplementary information contains


Supplementary Figure legend & Supplementary Figure 1, 2

Supplementary Figure legend



Supplementary Figure 1: Full length gels and western blot displayed in Figure 5a and 5f.

Supplementary Figure 2: The size of each adipocyte in Figure 2c and 2d was re-plotted (mean \pm S.D., n = 3,000-4,000 cells). Data were analyzed by two-way ANOVA with Sidak's multiple comparison. **** $p < 0.0001$.

Full length gel of MHC isoforms electrophoresis in Figure 5a



Full length blots in Figure 5f



Adipocyte Area

