Supplemental Table 2: Among current female contraceptive users ages 15-44, adjusted odds ratios from logistic regression assessing associations between user characteristics and use of individual methods, 2014

	pill		other hormonal methods		condom	
	aOR	p-value	aOR	p-value	aOR	p-value
Age						
15-24	ref		ref		ref	
25-34	1.0	.92	0.8	.32	1.2	.35
35-44	0.7	.18	0.5	.12	1.1	.62
Race/ethnicity						
White, non-Hispanic	ref		ref		ref	
Black, non-Hispanic	0.6	< 0.01	1.9	< 0.01	1.5	.02
Other or multiple races, non-Hispanic	0.6	.02	1.0	.91	1.4	.16
Hispanic	0.7	.09	1.5	.09	1.4	.06
Poverty status						
<100% federal poverty level	ref		ref		ref	
100-199%	1.2	.24	1.5	.09	0.8	.27
200-299%	1.4	.15	0.6	.19	0.8	.40
300% or higher	1.6	.02	0.9	.61	0.7	.15
Born outside US						
No					ref	
Yes					1.5	.03
Relationship status						
Married	ref		ref		ref	
Cohabitating	1.2	.50	1.5	.26	0.6	.08
Not married or cohabitating	1.5	.03	2.2	.02	0.7	.02
Education						
No high school diploma			ref		ref	
High school or GED			0.5	.01	1.3	.23
Some college			0.4	<0.01	0.9	.77
College graduate			0.4	.05	1.9	<0.01
Insurance coverage						
No	ref		ref		ref	
Yes (private, Medicaid, other)	1.9	<0.01	1.6	.09	0.6	<0.01
Parity						
0	ref		ref		ref	
1-2	0.3	<0.01	1.0	.97	0.8	.12
3 or more	0.2	<0.01	0.5	.06	0.5	.01
Number of future births expected						
0	ref				ref	
1-2	1.9	<0.01			2.5	<0.01
3 or more	1.8	.03			3.6	<0.01

Notes: 2014 represents the midpoint of the data collection years for the 2013-2015 NSFG survey. Other hormonal methods include the patch, ring and injectables.