

Population^a	2-3cM^b	3-5cM	5-10cM	10+cM	Total^c
Colombia	1.9	1.5	2.4	1.7	7.6
Cuba	1.1	0.9	1.1	0.7	3.7
Dominican Republic	1.2	1.6	3.3	3.1	9.2
Ecuador	1.9	1.4	1.8	1.4	6.5
Guatemala	1.4	1.4	2.1	1.6	6.6
Honduras	1.7	1.9	3.5	2.6	9.7
Mexico	1.2	0.8	0.7	0.3	3.0
Nicaragua	2.4	1.9	2.8	2.1	9.2
Puerto Rico	3.6	6.7	14.8	10.0	35.1
Memphis AA	1.5	0.4	0.2	0.2	2.3
Memphis EA	2.0	0.6	0.3	0.1	2.9
Pittsburgh AA	1.3	0.3	0.2	0.2	2.0
Pittsburgh EA	2.7	2.6	2.0	0.3	7.5

^a Population is country of origin of grandparents for the HCHS/SOL populations (first nine populations in table). For the Health ABC populations (last four populations in table), EA is European American, AA is African American.

^b Average total length of detected IBD in cM for segments with inferred length in the specified range, per pair of unrelated individuals in the sample from the population.

^c Average total length of detected IBD in cM for all segments with inferred length ≥ 2 cM, per pair of unrelated individuals in the sample from the population.