

Supplemental Table 1. Clinical characteristics of nonsmokers

Nonsmokers (n = 80)			
Age (years)	65	±	10
Male/Female	19	/	61
BMI (kg/m ²)	23.0	[19.8, 26.5]	
SBP (mmHg)	123	±	18
DBP (mmHg)	70	±	11
HbA1c (%)	5.8	[5.6, 6.1]	
HDL-C (mg/dL)	66	±	13
LDL-C (mg/dL)	127	±	39
MCP-1 (pg/mL)	310	[264, 380]	

Data are presented as mean ± SD or median [interquartile range].

BMI, body mass index; SBP, systolic blood pressures; DBP, diastolic blood pressures; HbA1c, hemoglobin A1c; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; MCP-1, Monocyte chemoattractant protein-1.

Supplemental Table 2. Gender-adjusted analysis on correlation between serum MCP-1 levels and clinical parameters in nonsmokers

	Univariate		Multivariate	
	β value	p value	β value	p value
Age (years)	0.303	0.006	0.418	0.001
BMI (kg/m^2)	0.106	0.367	0.285	0.029
SBP (mmHg)	0.184	0.105	-	-
DBP (mmHg)	0.087	0.463	-	-
HbA1c (%)	-0.066	0.590	-	-
HDL-C (mg/dL)	0.002	0.983	-	-
LDL-C (mg/dL)	-0.038	0.754	-	-

β value: Correlation coefficients, $R^2 = 0.162$

BMI, body mass index; SBP, systolic blood pressures; DBP, diastolic blood pressures; HbA1c, hemoglobin A1c; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol.