Supplementary data 4. Average cumulative dispensing of agents used in the treatment of inflammatory bowel disease in disease modifying medication in defined daily dose (DDD) in Norwegian patients with ulcerative colitis two years before and after first dispense of low dose naltrexone (LDN). Three groups based on number of LDN dispenses: LDN x 1 (N=54) collected LDN once, LDN x 2-3 (N=43) two or three times and LDN x 4+ (N=81) four or more times. Other immunosuppressants include azathioprine, mercaptopurine, methotrexate and tacrolimus.

| Ulcerative colitis | Group | Average DDD per patient | | Difference of mean | | |
|-------------------------------------|-----------|-------------------------|-----------|--------------------|-----------------------------------|-------|
| | | Before LDN | After LDN | DDD | (95% CI) | р |
| All outcome drugs | LDN x1 | 970.8 | 1026.3 | 55.5 | -198.5 to 309.5 | 0.663 |
| | LDN x 2-3 | 856.1 | 841.6 | -14.5 | -171.1 to 142.0 | 0.852 |
| | LDN x 4+ | 996.0 | 983.7 | -12.4 | -169.3 to 144.6 | 0.876 |
| Systemic immunosuppressants | LDN x1 | 338.1 | 395.6 | 57.6 | -0,74.1 to 189.2 | 0.384 |
| | LDN x 2-3 | 234.6 | 276.8 | 42.2 | -34.9 to 119.3 | 0.275 |
| | LDN x 4+ | 269.7 | 279.5 | 9.8 | -64.2 to 83.9 | 0.792 |
| Systemic glucocorticoids | LDN x1 | 180.8 | 159.4 | -21.4 | -91.8 to 48.9 | 0.544 |
| | LDN x 2-3 | 129.9 | 136.6 | 6.7 | -60.1 to 73.5 | 0.840 |
| | LDN x 4+ | 152.1 | 118.6 | -33.4 | -83.2 to 16.3 | 0.185 |
| TNF-α inhibitors | LDN x1 | 76.5 | 161.1 | 84.6 | -6.8 to 175.9 | 0.069 |
| | LDN x 2-3 | 55.2 | 93.1 | 38.0 | -24.8 to 100.7 | 0.229 |
| | LDN x 4+ | 55.8 | 66.8 | 11.0 | -18.3 to 40.2 | 0.458 |
| Other immunosuppressants | LDN x1 | 80.7 | 75.2 | -5.6 | -45.4 to 34.2 | 0.779 |
| | LDN x 2-3 | 49.5 | 47.0 | -2.4 | -16.2 to 11.3 | 0.722 |
| | LDN x 4+ | 61.8 | 94.1 | 32.3 | -10.4 to 75.0 | 0.136 |
| Intestinal anti-inflammatory agents | LDN x1 | 632.7 | 630.7 | -2.1 | -178.0 to 173.9 | 0.981 |
| | LDN x 2-3 | 621.5 | 564.8 | -56.8 | -170.0 to 56.5 | 0.318 |
| | LDN x 4+ | 726.4 | 704.2 | -22.2 | -158.6 to 114.2 | 0.747 |
| Intestinal corticosteroids | LDN x1 | 31.6 | 37.0 | 5.4 | -20.3 to 31.1 | 0.677 |
| | LDN x 2-3 | 50.5 | 27.4 | -23.0 | -59.9 to 13.8 | 0.214 |
| | LDN x 4+ | 37.1 | 15.6 | -21.5 | -55.8 to 12.8 | 0.214 |
| Aminosalisylates | LDN x1 | 601.1 | 593.7 | -7.4 | -177.9 to 163.0 | 0.930 |
| | LDN x 2-3 | 571.1 | 537.3 | -7.4 | -177.9 to 163.0 | 0.561 |
| | LDN x 4+ | 689.3 | 688.6 | -0.7 | -149.9 to 82.3 -122.6 to 121.3 | 0.561 |