

Electronic Supplementary Information

Site-specific characterization and quantitation of N-glycopeptides in PKM2 knockout breast cancer cells using DiLeu isobaric tags enabled by electron-transfer/higher-energy collision dissociation (ETHcD)

Zhengwei Chen,^{1#} Qing Yu,^{2#} Ling Hao,² Fabao Liu,³ Jillian Johnson,² Zichuan Tian,¹ W. John Kao,² Wei Xu,^{3,*} Lingjun Li^{1,2,4,*}

¹Department of Chemistry, University of Wisconsin, Madison, WI 53705, USA

²School of Pharmacy, University of Wisconsin, Madison, WI 53705, USA

³McArdle Laboratory for Cancer Research, University of Wisconsin, Madison, WI 53705, USA

⁴School of Life Sciences, Tianjin University, Tianjin, 300072, P.R. China

***Correspondence:** Professor Lingjun Li, School of Pharmacy and Department of Chemistry, University of Wisconsin-Madison, 777 Highland Avenue, Madison, Wisconsin 53705-2222

E-mail: lingjun.li@wisc.edu

Fax: +1-608-262-5345

Phone: +1-608-265-8491

and Professor Wei Xu, McArdle Laboratory for Cancer Research, University of Wisconsin, Madison, WI 53705, USA

Email: wxu@oncology.wisc.edu

These authors contributed equally.

Table of Contents:

Figure S1. The reporter ions intensity (A), precursor ions intensity (B) and reporter ion yield (C) of glycopeptides and nonglycosylated peptides in the same run.

Figure S2. The Venn diagram analysis of (A) intact N-glycopeptides (A) and (B) N-glycosylation sites identified with HILIC and MLA enrichment methods. (C) The comparison of the percentage of sialylated N-glycopeptides enriched between two methods.

Table S1 The identified N-glycopeptides enriched by HILIC alone, lectin alone and sequential HILIC and lectin enrichment.

Table S2 Glycoproteomics results between PKM2 knockout cells vs. parental breast cancer cells.

Table S3 Deglycoproteomics results between PKM2 knockout cells vs. parental breast cancer cells.

Table S4 Gene ontology analysis results.

Figure S1. Chen et al.




Figure S1. The reporter ions intensity (A), precursor ions intensity (B) and reporter ion yield (C) of glycopeptides and nonglycosylated peptides in the same run.

Figure S2. Chen et al.




Figure S2. The Venn diagram analysis of (A) intact N-glycopeptides (A) and (B) N-glycosylation sites identified with HILIC and MLA enrichment methods. (C) The comparison of the percentage of sialylated N-glycopeptides enriched between two methods.