

Supplemental Table S1. Correlation data between clinical outcomes and admission DOGiBAT scores at individual anatomic sites in all enrolled dogs (n=61). Spearman correlation coefficients (r_s) and p-values considered significant ($p < 0.05$) are in bolded text.

* Correlation data not available for intracranial bleeding due to lack of reported events.

	Blood product administration		Hospitalization length (days)		Survival to discharge	
	r_s	p-value	r_s	p-value	r_s	p-value
Skin	0.336	0.008	0.303	0.018	0.01	0.94
Catheter, etc.	0.252	0.049	0.232	0.072	-0.044	0.736
Oral cavity	0.265	0.04	0.195	0.135	-0.159	0.226
Ocular	0.07	0.6	-0.055	0.681	-0.314	0.016
Epistaxis	-0.034	0.797	0.152	0.243	-0.013	0.92
Gastrointestinal	0.438	0.002	0.248	0.096	-0.185	0.218
Urinary	0.337	0.015	0.157	0.265	-0.114	0.422
Pulmonary	-0.074	0.572	0.041	0.755	0.039	0.768
Intracranial*	--	--	--	--	--	--
Total DOGiBAT	0.512	<0.0001	0.35	0.006	-0.168	0.197

Supplemental Table S2. Correlation data between clinical outcomes and admission DOGiBAT scores at individual anatomic sites in only dogs categorized as having primary immune thrombocytopenia (n=34). Spearman correlation coefficients (r_s) and p-values considered significant ($p < 0.05$) in bolded text.

* Correlation data not available for intracranial bleeding due to lack of reported events.

DOGiBAT site	Blood product administration		Hospitalization length (days)		Survival to discharge	
	r_s	p-value	r_s	p-value	r_s	p-value
Skin	0.263	0.133	0.377	0.028	0.04	0.823
Catheter, etc.	0.183	0.3	0.145	0.411	-0.166	0.349
Oral cavity	0.345	0.049	0.364	0.038	-0.166	0.356
Ocular	0.096	0.601	0.012	0.948	-0.222	0.221
Epistaxis	-0.099	0.576	0.137	0.441	0.114	0.522
Gastrointestinal	0.39	0.044	0.199	0.32	-0.246	0.215
Urinary	0.331	0.074	0.143	0.453	0.062	0.744
Pulmonary	-0.137	0.44	0.063	0.724	0.054	0.761
Intracranial *	--	--	--	--	--	--
Total DOGiBAT	0.541	0.001	0.452	0.007	-0.123	0.49