

VERSION 2

Table 2: Characteristics of participants with baseline disability who demonstrated further functional decline over the one year period, versus those who remained stable or improved

	Progressive Functional Decline (n = 11)	Functional Stability or Improvement (n = 21)	P Value
Age at follow-up (mean \pm SD)	72.0 \pm 5.8	71.2 \pm 4.6	0.87
Male (%)	8/11 (73%)	13/21 (62%)	0.54
Time since transplant (yrs) (median, IQR)	6.9 (2.5,8.9)	3.8 (2.0,7.7)	0.12
No. of comorbidities at baseline (median, IQR)	4.7 (3,6)	3.0 (2,5)	0.79
TUG >15s at baseline (%)	6/10 (60%)	5/20 (25%)	0.06
Dominant handgrip <10 th percentile at baseline (%)	6/11 (55%)	9/21 (43%)	0.40
Median disabled ADLs at baseline (IQR)	3 (1,5)	2 (1,4)	1.00
No disability in instrumental ADLs at baseline (%)	8/11 (73%)	16/21 (76%)	0.51
Fall during year of follow-up (%)	6/11 (55%)	3/21 (14%)	0.02
Hospitalized during year of follow-up (%)	2/11 (18%)	9/21 (43%)	0.45