

# De Novo Synthesis of Elastin by Exogenous Delivery of Synthetic Modified mRNA into Skin and Elastin-Deficient Cells

Mario Lescan,<sup>1</sup> Regine Mariette Perl,<sup>1</sup> Sonia Golombek,<sup>1</sup> Martin Pilz,<sup>1</sup> Ludmilla Hann,<sup>1</sup> Mahua Yasmin,<sup>1</sup> Andreas Behring,<sup>1</sup> Timea Keller,<sup>1</sup> Andrea Nolte,<sup>1</sup> Franziska Gruhn,<sup>1</sup> Efrat Kochba,<sup>2</sup> Yotam Levin,<sup>2</sup> Christian Schlensak,<sup>1</sup> Hans Peter Wendel,<sup>1</sup> and Meltem Avci-Adali<sup>1</sup>

<sup>1</sup>Department of Thoracic and Cardiovascular Surgery, University Hospital Tübingen, Calwerstraße 7/1, 72076 Tübingen, Germany; <sup>2</sup>NanoPass Technologies, Ltd., 3 Golda Meir, 7403648 Nes Ziona, Israel

**Elastin is one of the most important and abundant extracellular matrix (ECM) proteins that provide elasticity and resilience to tissues and organs, including vascular walls, ligaments, skin, and lung. Besides hereditary diseases, such as Williams-Beuren syndrome (WBS), which results in reduced elastin synthesis, injuries, aging, or acquired diseases can lead to the degradation of existing elastin fibers. Thus, the *de novo* synthesis of elastin is required in several medical conditions to restore the elasticity of affected tissues. Here, we applied synthetic modified mRNA encoding tropoelastin (TE) for the *de novo* synthesis of elastin and determined the mRNA-mediated elastin synthesis in cells, as well as *ex vivo* in porcine skin. EA.hy926 cells, human fibroblasts, and mesenchymal stem cells (MSCs) isolated from a patient with WBS were transfected with 2.5 µg TE mRNA. After 24 hr, the production of elastin was analyzed by Fastin assay and dot blot analyses. Compared with untreated cells, significantly enhanced elastin amounts were detected in TE mRNA transfected cells. The delivered synthetic TE mRNA was even able to significantly increase the elastin production in elastin-deficient MSCs. In porcine skin, approximately 20% higher elastin amount was detected after the intradermal delivery of synthetic mRNA by microinjection. In this study, we demonstrated the successful applicability of synthetic TE encoding mRNA to produce elastin in elastin-deficient cells as well as in skin. Thus, this auspicious mRNA-based integration-free method has a huge potential in the field of regenerative medicine to induce *de novo* elastin synthesis, e.g., in skin, blood vessels, or alveoli.**

## INTRODUCTION

Elastin is an extracellular matrix (ECM) protein that provides elasticity and resilience to tissues and organs.<sup>1</sup> Thus, this protein is most abundant in organs where elasticity and flexibility are of major importance. Elastin constitutes about 28%–32% of major blood vessels, 30%–57% of the aorta, 50% of elastic ligaments, 3%–7% of lung, and 2%–3% of the dry weight of skin.<sup>2–4</sup> The expression of elastin primarily occurs during late and early neonatal periods,<sup>5</sup>

and after the first few years of life. From a young age, elastin synthesis decreases and ceases in adults.<sup>6</sup> During development, elastin is produced and secreted by different types of cells, such as smooth muscle cells (SMCs), fibroblasts, and endothelial cells.<sup>7</sup>

Although the half-life of elastin is about 74 years<sup>8</sup> and it is one of the most stable proteins known, the persistence of elastin, which is produced during development, is important for proper functioning of elastic connective tissues. Abnormalities in the elastin synthesis due to genetic disorders such as Williams-Beuren syndrome (WBS), also called elfin facies syndrome, autosomal dominant cutis laxa (ADCL), or supravalvular aortic stenosis (SVAS), result in several elastinopathies. Furthermore, in the skin of adults, aging, injuries, or sun damage also leads to the loss of elasticity, because the low level of elastin synthesis by the cells is not sufficient to efficiently repair the damage.


WBS occurs in 1 in 7,500–20,000 births<sup>9,10</sup> and is caused by the hemizygous microdeletion of approximately 26 to 28 genes on chromosome 7q11.23, which includes the elastin gene, resulting in less than 50% of normal elastin expression.<sup>11,12</sup> The condition is associated with connective tissue abnormalities and cardiovascular diseases, specifically SVAS and supravalvular pulmonary stenosis.<sup>13</sup> Homozygous elastin null mutants (ELN<sup>-/-</sup>) result in increased proliferation of arterial SMCs, and that leads to thickening of the arterial wall, narrowing of the arteries, and death within a few days of birth.<sup>14</sup> Cutis laxa is caused by mutations of elastin or fibulin-5 (FBLN5). It occurs in less than 1 in 1,000,000 births (<http://www.orpha.net>) and results in wrinkled, redundant, and sagging inelastic skin.<sup>15</sup> It is associated with hernias, cardiac valve anomalies, and cardiovascular diseases, e.g., pulmonary stenosis, aortic and arterial dilatation, and emphysema.

Received 23 November 2017; accepted 27 March 2018;  
<https://doi.org/10.1016/j.omtn.2018.03.013>

**Correspondence:** Meltem Avci-Adali, PhD, Department of Thoracic and Cardiovascular Surgery, University Hospital Tübingen, Calwerstr. 7/1, 72076 Tübingen, Germany.

**E-mail:** [meltem.avci-adali@uni-tuebingen.de](mailto:meltem.avci-adali@uni-tuebingen.de)





**Figure 1. Analysis of the Generated Synthetic TE mRNA and Cy3-Labeled TE mRNA by 1% Agarose Gel Electrophoresis**

(Lane 1) RNA marker, 400 ng of (lane 2) Cy3-labeled TE mRNA, (lane 3) modified, or (lane 4) unmodified TE mRNA was loaded on 1% agarose gel. A single band at around 2,500 bases confirmed the purity and specific length of the synthetic mRNA. First, (A) Cy3-labeled TE mRNA was detected using an UV-transilluminator, and then (B) all nucleic acids were detected by GelRed staining.

Elastin is synthesized as a soluble monomeric precursor, called tropoelastin (TE).<sup>16</sup> After the translation, TE is transported in association with elastin binding protein (EBP) to the extracellular space. Subsequently, TE is assembled on microfibrils and covalently cross-linked by lysyl oxidase (LOX) to form mature elastic fibers.

In recent years, the use of synthetic mRNAs as a new therapeutic drug for the production of desired proteins in cells gained in importance.<sup>17–21</sup> Synthetic mRNAs with or without modified nucleotides are generated by *in vitro* transcription (IVT) using RNA polymerases. Furthermore, a poly A-tail is added at the 3' end and a cap analog is incorporated at the 5' end of the mRNA to improve the stability and the translation of the generated synthetic mRNA. Compared with viral vectors and DNA plasmids, the application of synthetic mRNAs has several advantages: Synthetic mRNAs do not need to enter the cell nucleus, thereby insertional mutagenesis is prevented and non-dividing cells can be transfected to produce the desired protein. Additionally, mRNAs are smaller than plasmids and viral vectors, which allows improved delivery of synthetic mRNAs into the cells. After the release of mRNA into the cytosol, the mRNA is immediately translated by ribosomes into proteins. Furthermore, permanent protein overexpression-related complications are prevented, because the synthetic mRNA is transiently present due to natural degradation in cells.

Here, we generated synthetic modified TE encoding mRNA and analyzed the ability to produce elastin in EA.hy926 cells, human fibroblasts, and mesenchymal stem cells (MSCs) isolated from a patient with WBS. Afterward, the synthetic mRNA-mediated production of elastin in skin was analyzed using an *ex vivo* porcine skin model.

## RESULTS

### Synthesis of Modified TE mRNA and Analysis of Transfection Efficiency in EA.hy926 Cells and Fibroblasts

Modified TE mRNA containing 5mCTP and  $\Psi$  instead of cytidine triphosphate (CTP) and uridine triphosphate (UTP) was obtained

after the IVT. The agarose gel electrophoresis showed that the generated mRNA has the expected length of approximately 2,500 nt (Figure 1). Additionally, an unmodified TE mRNA was generated and the product was also analyzed. Due to Cy3 conjugation, only the specific band for Cy3-labeled TE mRNA could be detected using UV-transilluminator (Figure 1A). After staining with GelRed, all mRNAs could be detected, thereby successful labeling of TE mRNA with Cy3 was demonstrated.

The generated Cy3-labeled TE mRNA was used to analyze the transfection efficiency. Therefore,  $3 \times 10^5$  EA.hy926 cells or fibroblasts were transfected with lipoplexes containing 2.5  $\mu\text{g}$  Cy3-labeled TE mRNA. The fluorescence microscopy analyses revealed a high transfection efficiency, which was also confirmed by flow cytometry measurements (Figure 2). After the incubation of cells for 4 hr at 37°C with lipoplexes,  $98.16\% \pm 1.1\%$  of EA.hy926 cells and  $96.14\% \pm 1.9\%$  of human fibroblasts were transfected with Cy3-labeled TE mRNA.

### Characterization of WBS\_MSCs

The isolated MSCs from a patient with WBS (WBS\_MSCs) were characterized by staining with antibodies and performing of flow cytometry. The WBS\_MSCs were negative for the expression of CD31 and CD45, but expressed the characteristic marker of MSCs, CD90, and CD105 (Figure 3).


### Analysis of Cell Viability after the Delivery of Synthetic TE mRNA in EA.hy926 Cells

EA.hy926 cells were transfected with 2.5  $\mu\text{g}$  of modified or unmodified TE mRNA. After 24 hr, the cell viability was determined using PrestoBlue assay. As shown in Figure 4, the transfection with unmodified TE mRNA led to a strong cytotoxicity. In contrast, the viability of cells after the transfection with modified mRNA was not significantly different from the cells incubated with OptiMEM or only with transfection reagent. The modified TE mRNA treatment revealed with a detected cell viability of  $74\% \pm 11\%$ , a good cytocompatibility.

### Production of Elastin after the Transfection of Cells with Synthetic Modified TE mRNA

The production of elastin after the transfection of EA.hy926 cells, human fibroblasts, and WBS\_MSCs with 2.5  $\mu\text{g}$  synthetic modified TE mRNA was analyzed by using two independent methods, Fastin assay and dot blot analysis. Fastin assay detects elastin by binding to the basic and non-polar amino acid sequences, and dot blot analysis detects the elastin by using a specific antibody. Compared with control cells treated with OptiMEM containing Lipofectamine 2000, a significantly increased amount of elastin was detected after the delivery of synthetic modified TE mRNA into the cells using Fastin assay (Figure 5), as well as dot blot assay (Figure 6). In Table 1, the determined elastin amounts are compared.

Using Fastin assay, an approximately 16-fold higher elastin amount was detected (Figure 5) after the transfection of 2.5  $\mu\text{g}$  modified TE mRNA in EA.hy926 cells compared with the cells without mRNA



**Figure 2. Analysis of TE mRNA Transfection Efficiency Using Fluorescence Microscopy and Flow Cytometry**



$3 \times 10^5$  EA.hy926 cells and human fibroblasts were transfected with 2.5  $\mu$ g Cy3-labeled TE mRNA using Lipofectamine 2000. Cells incubated only with the transfection reagent were used as negative control. (A) Fluorescence microscopy and (B) flow cytometry analysis was performed 4 hr after the incubation of cells with lipoplexes. Black line represents cells treated only with the transfection reagent, and red line represents cells treated with Cy3 TE mRNA. BF, bright field.

treatment. In human fibroblasts about a 10-fold and in WBS-MSCs about a 13-fold higher amount of elastin was detected. Although statistically significant, the detected increase of elastin amount in dot blot assay (Figure S1) was less than the detected amounts in Fastin assay. Here, a 1.3-fold higher elastin amount was detected in TE mRNA transfected EA.hy926 cells (Figure 6) compared with control cells. In human fibroblasts, a 1.1-fold higher and 1.33-fold higher elastin amount was detected.

**Intradermal Delivery of Synthetic Modified TE mRNA by Microinjection Results in Increased Elastin Synthesis**

The ability of synthetic TE mRNA to mediate production of elastin was analyzed by microinjection of 2.5  $\mu$ g synthetic modified TE

**WBS\_MSCs**



**Figure 3. Characterization of MSCs Isolated from the Thymus of a WBS Patient**


(Top panel) Phase-contrast micrograph of MSCs at passage 1. (Bottom panels) Flow cytometry analysis of WBS\_MSCs after the staining with mouse anti-human antibodies against CD90, CD105, CD31, and CD45. Black line: negative control; red line: WBS\_MSCs stained with specific antibodies.

mRNA complexed with Lipofectamine 2000 in a total volume of 35  $\mu$ L into porcine skin. Using Fastin assay, we detected a 1.2-fold increased amount of elastin in the skin treated with synthetic modified TE mRNA compared with the skin without mRNA treatment (Figure 7).

**DISCUSSION**

Elastin is one of the most important components of the blood vessels, lungs, and skin. It provides elasticity and resilience to skin and has major effects on elastic properties of the lungs. Loss of elastin fibers in the aortic wall due to aging leads to functional and morphologic changes as observed in the impairment of the “Windkessel” function and aneurysm formation.<sup>22</sup> Degradation of elastin in the skin due to aging, UV light exposure, or injury leads to impairment of elasticity and results in formation of wrinkles and scarring. Furthermore, diseases caused by mutations or deletion of the elastin gene, such as cutis laxa or WBS, can affect the functionality of various organs and especially blood vessels.

In this study, for the first time, we successfully demonstrated that the elastin synthesis can be significantly increased by the delivery of



**Figure 4. Analysis of the Influence of TE mRNA Transfection on the Viability of EA.hy293 Cells Using PrestoBlue Assay**


$4 \times 10^5$  EA.hy293 cells were transfected with 2.5  $\mu$ g unmodified TE mRNA (unmod TE mRNA) or 2.5  $\mu$ g modified TE mRNA (mod TE mRNA). Furthermore, cells were incubated only with OptiMEM or with transfection reagent (TR) Lipofectamine 2000. PrestoBlue cell viability assay was performed 24 hr after the transfection of cells. The viability of cells incubated with OptiMEM was set to 100%, and the viability relative to these cells was expressed. Data are shown as mean  $\pm$  SEM (n = 3). Statistical differences were determined using one-way ANOVA with Bonferroni's multiple comparisons test. \*\*\*p < 0.001; \*\*\*\*p < 0.0001.

synthetic modified TE mRNA into cells as well as in porcine skin via microinjection. This approach is a new and promising method for counteracting pathological changes caused by loss of elastin in adult dermis. Previous studies with transforming growth factor  $\beta$ 1 (TGF- $\beta$ 1) demonstrated positive effects on elastin synthesis in human fibroblasts,<sup>23–25</sup> rat aortic SMCs,<sup>26</sup> and transgenic mice.<sup>27,28</sup> In further studies, Mizuno et al.<sup>29</sup> transfected rat endothelial cells with plasmids to express recombinant elastin. The transplantation of these cells after a myocardial infarction into the myocardial infarct scar reduced scar expansion and prevented left ventricular enlargement. Later, Xiong and colleagues<sup>30</sup> demonstrated that aortic vascular SMCs can be efficiently transfected *in vitro* and *in vivo* with an adenoviral vector carrying a recombinant TE gene. The application of this vector in elastase-induced abdominal aortic aneurysm in rats resulted in accumulation of elastic fibers leading to a reduction of aortic diameter in the medial layer *in vivo*, which indicates that the expression of recombinant elastin and the reconstruction of elastic fibers within the aneurysmal tissue could be used to prevent or reverse the aneurysm dilatation. In a recent study, Xie and colleagues<sup>31</sup> subcutaneously injected recombinant human TE into wound bed of a severe burn injury and demonstrated formation of new elastin fibers in a swine model. Despite the ability of TGF- $\beta$ 1 to induce elastin synthesis, it should be also mentioned that TGF- $\beta$ 1 can act as a significant stimulator

of tumor progression, invasion, and metastasis.<sup>32</sup> Welch et al.<sup>33</sup> demonstrated that the *in vitro* pretreatment of adenocarcinoma cells with TGF- $\beta$ 1 enhanced the metastatic ability of these cells. Furthermore, the use of adenovirus-based vectors has the disadvantage that these vectors are highly immunogenic and often have high levels of pre-existing immune responses in the host, which can restrict the delivery of vectors into desired cells.<sup>34</sup> The delivery of genes by using retroviral vectors can lead to the permanent expression of proteins, which can have adverse effects over time. Furthermore, these vectors transduce only replicating cells and they randomly incorporate into the genome of host cells, and this can result in insertional mutagenesis.<sup>35</sup> In contrast, synthetic mRNA is shorter than plasmid DNA and does not need to enter the nucleus. After the delivery of mRNA into the cytosol, it is immediately translated by ribosomes into desired protein. Thus, compared with other gene delivery methods, the use of synthetic TE mRNA has huge potential in the field of regenerative medicine.

The successful detection of elastin after the transfection of cells with modified TE mRNA in the supernatant indicates that the synthesized TE could successfully bind to EBP and be transported to the cell surface. After the delivery of TE to the cell surface, the binding of galactosugars to EBP leads to the dramatic decrease of the affinity of the elastin site for its ligands and results in dissociation of TE from EBP.<sup>36</sup> EBP is a spliced variant of lysosomal  $\beta$ -galactosidase, and it is also a part of the elastin receptor complex (ERC), which is a heterotrimer composed of EBP and two membrane-associated proteins, protective protein/cathepsin A and neuraminidase-1 (Neu-1). Especially, vascular SMCs express ERC, which directly interacts with elastin and mediates elastin-induced cellular activities.<sup>37,38</sup> Thus, in a previous study, the incorporation of  $\alpha$ -elastin into type 1 collagen gel resulted in significant inhibition of SMC proliferation and migration.<sup>39</sup>

Using the transfection reagent Lipofectamine 2000, a highly efficient transfection of TE mRNA could be achieved. After 4 hr of transfection, 98% of EA.hy926 cells and 96% of human fibroblasts were transfected with the Cy3-labeled TE mRNA. Furthermore, the transfection of EAHy.926 cells with modified TE mRNA demonstrated no significant influence on cell viability. In contrast, the unmodified TE mRNA led to significantly reduced cell viability. We additionally analyzed the quality of the unmodified and modified TE mRNA by Agilent Bioanalyzer and because the quality of the generated mRNAs was high and similar in both samples, high-performance liquid chromatography (HPLC) purification, which is recommended by Karikó et al.<sup>40</sup> and Weissman et al.<sup>41</sup> to reduce immune reactions to synthetic mRNAs, was not performed in this study. Therefore, we assume that the use of unmodified nucleotides led to a stronger immune activation than modified nucleotides. In a previous study, Thess and colleagues<sup>42</sup> successfully demonstrated that sequence engineering and the purification of synthetic mRNA with HPLC are able to avoid immune activation and to produce high levels of encoded protein *in vitro* and *in vivo*. Thus, we suggest that sequence engineering (codon optimization) of the unmodified mRNA could further reduce immune stimulation.



**Figure 5. Detection of Elastin Amount after the Transfection of EA.hy293 Cells, Human Fibroblasts, and MSCs Derived from WBS Patient with Synthetic Modified TE mRNA Using Fastin Assay**

$3 \times 10^5$  EA.hy926 cells, human fibroblasts, or WBS\_MSCs were transfected with 2.5 µg synthetic modified TE mRNA. After 24 hr, the elastin amount was detected using Fastin assay. As a control, cells were incubated with OptiMEM containing Lipofectamine 2000 without TE mRNA. Results are shown as mean  $\pm$  SEM (EA.hy926 cells:  $n = 4$ , fibroblasts:  $n = 6$ , WBS\_MSCs:  $n = 4$ ). Statistical differences were determined using paired t test. \* $p < 0.05$ ; \*\* $p < 0.01$ .

In a supplementary experiment, we analyzed using ELISA whether the translation of TE mRNA can be increased using m1Ψ or m1Ψ/5mCTP compared with the used modification Ψ/5mCTP (Figure S2). Thereby, we could show that by using m1Ψ or m1Ψ/5mCTP modifications, the translation of the TE mRNA could be significantly increased in EA.hy293 cells. The highest elastin concentration was detected after the transfection of cells with m1Ψ-modified TE mRNA.

Using dot blot assay, the detected elastin amount differences were less than using Fastin assay. The reason, therefore, could be the quick coverage of the nitrocellulose membrane with other proteins, such as albumin, that are present in high amounts in the medium. The detected differences of elastin amount by using Fastin assay were approximately 10-fold higher than in the dot blot assay. However, using the dot blot assay, the detected elastin amount in the synthetic modified TE mRNA transfected samples was also significantly higher than in the controls.

Importantly, we also demonstrated that synthetic modified TE mRNA can be locally delivered into the skin by using hollow micro-needles. Using this method, the consistency and reliability of intradermal injections, which are difficult to conduct with standard needles, can be improved.<sup>43</sup> Thus, this method can warrant a reliable intradermal delivery of mRNA and a targeted localization of mRNA in the skin. In our study, the microinjected modified synthetic TE mRNA was able to mediate elastin synthesis in the skin. The detected increase in elastin amount was approximately 1.2-fold, which corresponds to an increase of 20%.

In an additional experiment, human fibroblasts transfected with modified TE mRNA were seeded on gelatin scaffolds and after 21 days, the production of elastin fibers was analyzed using two-photon laser scanning microscopy (LSM). The scaffolds seeded

with TE mRNA transfected cells showed at an excitation wavelength of 760 nm an increased autofluorescence signal compared with the control scaffolds seeded with cells without TE mRNA transfection (Figure S3). Elastin emits autofluorescent light from 403–531 nm when excited at 760 nm.<sup>44</sup> Besides promising *ex vivo* skin results, these results indicate the generation of elastin fibers. However, particularly, extensive *in vivo* studies will allow us to obtain detailed information regarding the generation of functional elastic fibers, and currently, pig *in vivo* experiments are in progress.

## Conclusions


The ECM protein elastin gives connective tissues its elasticity and thereby enables the proper functioning of several organs and tissues, such as the blood vessels, elastic ligaments, lung, and skin. Because this protein is physiologically synthesized only to a limited extent after the neonatal period, elastin deficiencies caused by degenerative changes or genetic defects lead to significant functional limitations in the affected organs or tissues. So far, these deficiencies can be only treated symptomatically. Thus, the presented approach, using synthetic modified TE mRNA for *de novo* synthesis of elastin, represents an enormous potential for the therapy of congenital or acquired elastin deficiencies to restore the lost elasticity. Furthermore, because elastin has a long half-life, a few treatments or even only one treatment could be enough to obtain an improvement of elasticity in the affected tissues.

## MATERIALS AND METHODS

### Ethics Statement

The study was performed in accordance with the Federation of European Laboratory Animal Science Associations (FELASA) and American Association for Laboratory Animal Science (AALAS) recommendations on the care and use of laboratory animals. The Animal Care and Welfare Commissioner of the University of Tübingen approved the protocols and procedures.

MSCs were isolated from the thymus tissue of a 1-year-old child suffering from WBS who had undergone cardiovascular surgery. The guardians of the patient gave informed consent to participate



**Figure 6. Detection of Elastin Amount after the Transfection of EA.hy9293 Cells, Human Fibroblasts, and MSCs Derived from WBS Patient with Synthetic Modified TE mRNA Using Dot Blot Assay**

$3 \times 10^5$  EA.hy926 cells, human fibroblasts, or WBS\_MSCs were transfected with 2.5 µg synthetic modified TE mRNA. As a control, cells were incubated with OptiMEM containing Lipofectamine 2000 without TE mRNA. After 24 hr, 500 µL supernatant was blotted on nitrocellulose membrane, and elastin was detected using polyclonal antibody to human aortic elastin and the subsequent detection of alkaline-phosphatase-conjugated secondary antibody. Results are shown as mean  $\pm$  SEM (EA.hy926 cells:  $n = 3$ , fibroblasts:  $n = 3$ , WBS\_MSCs:  $n = 4$ ). Statistical differences were determined using paired t test. \* $p < 0.05$ ; \*\*\* $p < 0.001$ .

and for the publication of this report. The study was approved by the Ethical Committee of the Faculty of Medicine of the University of Tübingen.

#### Animals

German Landrace pigs ( $70 \pm 10$  kg, female, 4–6 months old) were obtained from a local, specific pathogen-free breeding facility (Benz, Germany). After the euthanasia of pigs by intravenous injection of a lethal dose of potassium chloride ( $1 \text{ mmol kg}^{-1}$ ; B. Braun Melsungen AG, Melsungen, Germany), the ears were explanted to obtain pig skin from the outer side of the ear.

#### Synthesis of TE mRNA

The plasmid pcDNA 3.3-containing human TE encoding sequence was produced by Aldevron (Fargo, ND, USA). The synthesis of TE mRNA was performed by IVT according to our previously established protocol.<sup>45</sup> In brief, the HotStar HiFidelity Polymerase Kit (QIAGEN, Hilden, Germany) and 0.7 µM of each forward (5'-TTGGACCCTCGTACAGAAGCTAATACG-3') and reverse primer (5'-T<sub>120</sub>-CTTCCTACTCAGGCTTTATTCAAAGACCA-3') were used to amplify the plasmid insert by PCR. Primers were purchased from ELLA Biotech (Martinsried, Germany) in HPLC grade. PCR was run using the following cycling protocol: initial activation step at 94°C for 3 min, followed by 25 cycles of denaturation at 94°C for 45 s, annealing at 60°C for 1 min, extension at 72°C for 1 min, and final extension at 72°C for 5 min. During the PCR, a poly T-tail of 120 thymidines (T) was added to the insert. PCR product was purified using the QIAquick PCR purification kit (QIAGEN, Hilden, Germany) according to the manufacturer's instructions.

The DNA is then *in vitro* transcribed into mRNA using the MEGAscript T7 Kit (Life Technologies, Darmstadt, Germany). To obtain modified TE mRNA, we performed the IVT reaction at 37°C

for 4 hr, and the mixture contained 7.5 mM ATP, 1.875 mM GTP (both from MEGAscript T7 Kit), 7.5 mM 5-methylcytidine (5mCTP), 7.5 mM pseudouridine ( $\Psi$ ) (both from TriLink BioTechnologies, San Diego, CA, USA), 2.5 mM 3'-O-Me-m<sup>7</sup>G(5')ppp(5') G RNA cap structure analog (New England Biolabs, Frankfurt am Main, Germany), 40 U RiboLock RNase inhibitor (Thermo Scientific, Waltham, MA, USA), and 1.5 µg PCR product. In contrast with modified TE mRNA, unmodified TE mRNA was synthesized by using 7.5 mM ATP, 1.875 mM GTP, 7.5 mM CTP, and 7.5 mM UTP instead of modified nucleotides  $\Psi$  and 5mCTP. Afterward, template DNA was removed by adding 1 µL TURBO DNase (from MEGAscript T7 Kit) and incubation at 37°C for 15 min. Then the reaction mixture was purified using the RNeasy Mini Kit (QIAGEN, Hilden, Germany) according to manufacturer's instructions. The mRNA was treated for 30 min at 37°C with 15 U Antarctic phosphatase (New England Biolabs, Frankfurt am Main, Germany) and purified again using the RNeasy Mini Kit. The concentration of mRNA was adjusted to 100 ng/µL by adding nuclease-free water. The quality and purity of the PCR product and the synthetic mRNA were assessed by 1% agarose gel electrophoresis and staining with 1× GelRed (Biotium, Fremont, CA, USA) in 1× Tris-borate-EDTA (TBE) for 30 min at room temperature (RT). The obtained TE mRNA was stored at  $-80^\circ\text{C}$ .

#### Cy3 Labeling of Synthetic TE mRNA

Cu(I)-free azide-(dibenzocyclooctyne [DBCO]) click chemistry was used to perform Cy3 labeling of synthetic TE mRNA. Therefore, 1.9 mM 5-azido-C<sub>3</sub>-UTP (Jena Bioscience, Jena, Germany) and 5.6 mM  $\Psi$  were used during the IVT instead of 7.5 mM  $\Psi$ . After the IVT, the reaction mixture was purified using the RNeasy MinElute Cleanup Kit (QIAGEN) according to the manufacturer's instructions. Afterward, Cy3 was conjugated to 5-azido-C<sub>3</sub>-UTP-modified mRNA by incubation with 5-fold molar excess of DBCO-sulfo-Cy3 (Jena Bioscience) in a total amount of 40 µL nuclease-free water for 1 hr at 37°C. The molar amount of DBCO-sulfo-Cy3 in relation to the amount of azide-labeled mRNA was calculated using the manufacturer's data sheet. Then the reaction mixture was purified using the RNeasy MinElute Cleanup Kit, and the quality and purity of the

**Table 1. Comparison of the Detected Elastin Amount Using Fastin Assay and Dot Blot Assay**

Cell Type	Fastin Elastin Assay			Dot Blot Assay		
	Elastin Amount ( $\mu\text{g}$ )			Intensity of Antibody Staining		
	w/o mRNA	w TE mRNA	Fold-Change	w/o mRNA	w TE mRNA	Fold-Change
EA.hy926 cells	3.17 $\pm$ 1.18	50.66 $\pm$ 9.40	16	27.52 $\pm$ 7.29	36.45 $\pm$ 7.24	1.3
Fibroblasts	5.02 $\pm$ 0.88	52.46 $\pm$ 8.67	10.5	83.72 $\pm$ 1.07	91.39 $\pm$ 0.71	1.1
WBS_MSCs	2.66 $\pm$ 1.15	35.32 $\pm$ 4.9	13.3	19.6 $\pm$ 0.74	26.09 $\pm$ 2.31	1.3

w, with; w/o, without.

mRNA were verified by 1% agarose gel electrophoresis and staining with 1 $\times$  GelRed in 1 $\times$  TBE for 30 min at RT. The concentration was determined using a photometer, and the mRNA was stored at  $-80^\circ\text{C}$ .

#### Isolation of MSCs from a Patient with WBS

To isolate MSCs from a patient with WBS (WBS\_MSCs), we rinsed the thymus tissue of a WBS patient with Dulbecco's PBS (DPBS; Life Technologies, Darmstadt, Germany) and cut it into small pieces. The tissue pieces were then incubated with 5 mL of 0.1% collagenase II solution (Merck Millipore, Darmstadt, Germany) in DMEM low glucose (Thermo Fisher Scientific, Waltham, MA, USA) at  $37^\circ\text{C}$ . The solution containing tissue pieces was briefly mixed every 30–60 min. After 3 hr, the tissue suspension was filtered through a 100- $\mu\text{m}$  tissue strainer and the filtrate was centrifuged for 5 min at  $300 \times g$ . After centrifugation, the supernatant was aspirated and the cell pellet was resuspended in 30 mL of MSC medium and seeded in three T175 cell culture flasks. The cells were then incubated in 25 mL medium at  $37^\circ\text{C}$  and 5%  $\text{CO}_2$ . The obtained cells are called WBS\_MSCs.

#### Characterization of Isolated WBS\_MSCs Using Flow Cytometry

Flow cytometry analysis was performed to determine the expression of surface markers on the cultured WBS\_MSCs. To perform the antibody staining, we first washed  $5 \times 10^5$  cells with 500  $\mu\text{L}$  DPBS and then with 1 mL DPBS containing 0.5% BSA (Sigma-Aldrich). Subsequently, the cells were incubated for 30 min at  $37^\circ\text{C}$  in 500  $\mu\text{L}$  DPBS/0.5% BSA containing antibodies (1:100 dilution). PE-conjugated mouse anti-human CD90/Thy-1 or CD45 antibodies (both from R&D Systems, Minneapolis, MN, USA), Alexa 488-conjugated mouse anti-human CD105 (Bio-Rad, Hercules, CA, USA), and PE-conjugated mouse anti-human CD31 (BD Biosciences, Heidelberg, Germany) were used for the staining. Afterward, cells were centrifuged at  $300 \times g$  for 5 min, washed, fixed in 1 $\times$  CellFIX solution (10 $\times$ ; Becton Dickinson, Heidelberg, Germany), and measured immediately using a FACScan flow cytometer (Becton Dickinson). A total of 10,000 cells was counted and analyzed using CellQuest Pro Software (Becton Dickinson).

#### Cultivation of Cells

EA.hy926 cells were cultivated in DMEM with high glucose (Lonza, Basel, Switzerland) supplemented with 10% heat-inactivated FBS (Life Technologies), 1% L-glutamine (Life Technologies), and 1%


penicillin/streptomycin (Life Technologies). Human BJ fibroblasts were cultivated in DMEM with high glucose containing 10% heat-inactivated FBS, 3% HEPES (Life Technologies), 1% L-glutamine, and 1% penicillin/streptomycin. WBS\_MSCs were cultivated in DMEM/Ham's F12 (1:1) medium (Biochrom, Berlin, Germany) containing 10% heat-inactivated FBS, 1% L-glutamine, and 1% penicillin/streptomycin. Cells were cultivated at  $37^\circ\text{C}$  and 5%  $\text{CO}_2$ . Medium was changed every 3–4 days. After reaching 80% confluency, cells were washed once with DPBS and detached using 0.05% Trypsin-EDTA (Life Technologies). Subsequently, trypsin was inactivated by adding trypsin neutralization solution (TNS) 0.05% in 0.1% BSA (PromoCell, Heidelberg, Germany). Cell amount was measured using CASY Cell Counter (Schärfe System, Reutlingen, Germany).

#### Transfection of Cells with Synthetic TE mRNA

For the transfection,  $3 \times 10^5$  EA.hy926 cells, BJ fibroblasts, or WB\_MSCs were seeded per well of a six-well plate, and the cells were cultivated overnight at  $37^\circ\text{C}$  in a 5%  $\text{CO}_2$  atmosphere. To generate transfection complexes, called lipoplexes, we incubated 2.5  $\mu\text{g}$  TE mRNA with 4  $\mu\text{L}$  Lipofectamine 2000 in 500  $\mu\text{L}$  OptiMEM I reduced serum-free media (Invitrogen, Carlsbad, CA, USA) for 20 min at RT. Cell medium was discarded and cells were washed once with 1 mL DPBS. Afterward, 500  $\mu\text{L}$  OptiMEM was added to each well of the six-well plate. Subsequently, 500  $\mu\text{L}$  OptiMEM containing lipoplexes was added to the wells. Additionally, 500  $\mu\text{L}$  OptiMEM containing only 4  $\mu\text{L}$  Lipofectamine 2000 was added as a control. After 4 hr of incubation at  $37^\circ\text{C}$ , transfection mixtures were removed and 1 mL cell culture medium without phenol red was added to each well. Cells were cultivated at  $37^\circ\text{C}$  and 5%  $\text{CO}_2$  overnight. The supernatant was collected and stored at  $-80^\circ\text{C}$ . Cells were detached using 0.05% Trypsin-EDTA (Life Technologies) and 0.05% TNS in 0.1% BSA (PromoCell, Heidelberg, Germany).

#### Analysis of Transfection Efficiency Using Cy3-Labeled TE mRNA

To assess the uptake of the Cy3 TE mRNA into EA.hy926 cells and human fibroblasts, we transfected  $3 \times 10^5$  cells as described above and 4 hr post-transfection, cells were washed twice with 1 mL DPBS. The Cy3 TE mRNA uptake into the cells was examined using fluorescence microscopy (Axiovert135; Carl Zeiss, Oberkochen, Germany). Additionally, the cells were detached, washed with 1 mL DPBS, and resuspended in 500  $\mu\text{L}$  1 $\times$  CellFIX solution to perform flow cytometry analysis. Flow cytometry measurements



**Figure 7. Detection of Elastin Amount after the Microinjection of Synthetic Modified TE mRNA into Pig Skin**

2.5 µg synthetic modified TE mRNA was complexed with Lipofectamine 2000 and injected using hollow microneedles into the pig skin. After 72 hr, elastin amount in the skin was detected using Fastin assay. The elastin amount is presented relative to the skin samples treated only with the transfection reagent without mRNA. Data are presented as box plot, median with 25–75 percentiles (box) and 5–95 percentiles (whiskers) ( $n = 7$ ). Statistical differences were determined using paired t test.  $**p < 0.01$ .

were performed using a FACScan flow cytometer. A total of 10,000 cells was counted and analyzed using CellQuest Pro Software.

#### **Analysis of Cell Viability after the Transfection of EA.hy926 Cells with Modified and Unmodified TE mRNA**

To analyze the cell viability after the transfection of EA.hy926 cells with modified and unmodified TE mRNA, we seeded  $4 \times 10^5$  cells per well of a six-well plate, cultivated them overnight, and transfected for 4 hr at 37°C with lipoplexes containing 2.5 µg modified or unmodified TE mRNA. As negative controls, cells were treated only with OptiMEM or OptiMEM containing transfection reagent. After 24 hr, the cell viability was measured using PrestoBlue assay (Invitrogen, Carlsbad, CA, USA). Therefore, cells were washed with 1 mL DPBS, and 50 µL PrestoBlue cell viability reagent was added to 450 µL cell culture medium per well and incubated for 1.5 hr at 37°C. The fluorescence intensity of 100 µL supernatant was measured in triplicates at an excitation wavelength of 530 nm and emission wavelength of 600 nm using a multimode microplate reader (Mithras LB 940; Berthold Technologies).

#### **Analysis of Elastin Synthesis in Cells**

##### **Fastin Assay**

Fastin Elastin Kit (Biocolor, Carrickfergus, UK) was used to detect soluble TE and  $\alpha$ -elastin polypeptides by first binding a dye label (5,10,15,20-tetraphenyl-21H,23H-porphine tetra-sulfonate [TPPS]) to the basic and non-polar amino acid sequences in mammalian elastin. By adding the Dye Dissociation Reagent, the elastin-bound dye is released into solution and measured at 513 nm using a microplate reader (EON Microplate Spectrophotometer; BioTek, Bad

Friedrichshall, Germany). Fastin assay was performed according to the manufacturer's instructions. An  $\alpha$ -elastin standard from 1.25 to 80 µg was used to detect the elastin quantity in the samples.

##### **Dot Blot Analysis**

To detect the produced elastin by dot blot analysis, we blotted 500 µL supernatant using the Bio-Dot SF device (Bio-Rad, Hercules, CA, USA) on nitrocellulose membrane according to the manufacturer's instructions. The membrane was incubated on a tilting shaker overnight at 4°C in TBST (20 mM Tris [pH 7.5], 500 mM NaCl, 0.05% Tween 20) containing 5% milk powder (Carl Roth, Karlsruhe, Germany). The following day, the membrane was incubated for 1 hr at RT with polyclonal antibody to human aortic elastin (EPC, Owensville, MO, USA) in TBST (1:200 dilution). Then the membrane was washed 3× for 5 min with TBST. Afterward, the membrane was incubated for 45 min at RT with alkaline-phosphatase-conjugated anti-rabbit IgG (Sigma-Aldrich, Munich, Germany) 1:10,000 diluted in TBST. Subsequently, the membranes were washed 3× for 5 min with TBST and incubated with the BCIP/NBT (5-bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium) solution (Sigma-Aldrich, Munich, Germany) for colorimetric detection of alkaline-phosphatase-conjugated antibodies. The intensities of dot blots were analyzed using ImageJ software.

#### **Delivery of TE mRNA into Pig Skin**

##### **Preparation of Porcine Skin**

The skin was separated postmortem from the pig's ears, washed with 0.9% NaCl solution (Fresenius Kabi, Bad Homburg, Germany), and trimmed to 1-mm thickness using a dermatome (Acculan 3Ti dermatome; B. Braun, Melsungen, Germany). Afterward, round pieces with a diameter of 15 mm were punched out. Using an iodine solution (0.5 M I<sub>2</sub>; Fluca Analytical, Seelze, Germany), we disinfected the surface of the punched skin pieces and then rinsed one to two times with DPBS. Furthermore, the skin was incubated for 30 min in an antibiotic solution composed of 250 µg/mL gentamicin (Sigma-Aldrich, St. Louis, MO, USA) and 1,250 µg/mL amphotericin B (PromoCell, Heidelberg, Germany) in DMEM (Thermo Fisher Scientific, Waltham, MA, USA) and then washed one to two times with DPBS. The excessive DPBS was removed using a sterile swab from the surface before performing intradermal injection.

##### **Intradermal Delivery of Synthetic TE mRNA into Ex Vivo Porcine Skin Model**

To perform the microinjections, 2.5 µg of synthetic TE mRNA was complexed with 4 µL Lipofectamine 2000 in OptiMEM in a total volume of 35 µL. The appropriate volume of OptiMEM containing Lipofectamine 2000 served as control. The intradermal injection of synthetic mRNA was performed using the hollow microinjection needles MicronJet600 from NanoPass Technologies (Nes Ziona, Israel). MicronJet600 is composed of three hollow microneedles of 600-µm length that are cleared for intradermal injections. After the injection of synthetic mRNA, each skin piece was washed 1× with 1 mL DPBS and transferred into ThinCert cell culture inserts with a pore size of 1 µm (Greiner Bio-One, Frickenhausen, Germany). The inserts



containing the skin were placed in one well of a 12-well plate, which contained 1.5 mL human endothelial cell culture medium (VascuLife EnGS-Mv Microvascular Endothelial Kit; without hydrocortisone hemisuccinate; CellSystems, Troisdorf, Germany). The skin was collected after 72 hr to determine the elastin amount by using Fastin assay.

### Detection of TE mRNA-Mediated Elastin Synthesis in Pig Skin Using Fastin Assay

#### Preparation of Skin

The medium on skin samples was swabbed, and the transfected region of the skin was cut from the skin piece using a scalpel. Samples were dried for 15 min at RT and then transferred into a reaction tube and immediately closed to prevent weight fluctuations due to evaporation. The weight of the individual skin samples was on average at  $17 \pm 3$  mg.

#### Fastin Assay

Skin samples were treated for 1 hr at  $100^\circ\text{C}$  with  $750\ \mu\text{L}$   $0.25\ \text{M}$  oxalic acid and then centrifuged at  $10,000 \times g$  for 10 min. The supernatant was collected and transferred into new reaction tubes. An  $\alpha$ -elastin standard from 1.25 to  $80\ \mu\text{g}$  was used to detect the elastin quantity in the samples. To each sample,  $500\ \mu\text{L}$  elastin precipitating reagent was added, incubated for 15 min, and then centrifuged at  $16,000 \times g$  for 10 min. The supernatant was discarded,  $1\ \text{mL}$  of dye reagent was added to each tube, mixed, and incubated for 90 min at RT and 550 rpm. Then the samples were centrifuged at  $12,000 \times g$  for 10 min. The supernatant was discarded and  $400\ \mu\text{L}$  of dye dissociation reagent was added, mixed, and incubated 10 min at RT until the pellet is dissolved. Afterward,  $100\ \mu\text{L}$  of this solution was transferred into one well of a 96-well plate in triplicates, and the absorption was measured at 513 nm using a microplate reader (EON Microplate Spectrophotometer).

#### Statistical Analysis

Data are shown as mean  $\pm$  SEM or median with 25–75 percentiles (box) and 5–95 percentiles (whiskers). One-way ANOVA followed by Bonferroni's multiple comparison test was performed to compare the means of more than two groups. Student's two-sample t test was performed for comparison of means between two groups. All statistical analyses were performed double-tailed using GraphPad Prism version 6.01. Differences of  $p < 0.05$  were considered significant.

#### SUPPLEMENTAL INFORMATION

Supplemental Information includes three figures and can be found with this article online at <https://doi.org/10.1016/j.omtn.2018.03.013>.

#### AUTHOR CONTRIBUTIONS

M.L., R.M.P., and M.A.-A. conceived and designed the experiments. R.M.P., S.G., M.P., L.H., M.Y., and F.G. performed the experiments with support from M.L., A.B., T.K., A.N., M.A.-A., and H.P.W. and analyzed the data. E.K., Y.L., H.P.W., and C.S. contributed reagents/materials/analysis tools. M.A.-A. wrote the paper and supervised the project.

#### ACKNOWLEDGMENTS

This study was funded by the German Heart Foundation/German Foundation of Heart Research through Grant F/38/13 and the European Social Funds in Baden-Wuerttemberg, Germany, and the Ministry of Science, Research, and the Arts of the State of Baden-Wuerttemberg (MWK-BW). Furthermore, we acknowledge support by Deutsche Forschungsgemeinschaft and Open Access Publishing Fund of University of Tübingen.

#### REFERENCES

- Baldwin, A.K., Simpson, A., Steer, R., Cain, S.A., and Kielty, C.M. (2013). Elastic fibres in health and disease. *Expert Rev. Mol. Med.* *15*, e8.
- Uitto, J. (1979). Biochemistry of the elastic fibers in normal connective tissues and its alterations in diseases. *J. Invest. Dermatol.* *72*, 1–10.
- Vrhovski, B., and Weiss, A.S. (1998). Biochemistry of tropoelastin. *Eur. J. Biochem.* *258*, 1–18.
- Annabi, N., Mithieux, S.M., Camci-Unal, G., Dokmeci, M.R., Weiss, A.S., and Khademhosseini, A. (2013). Elastomeric recombinant protein-based biomaterials. *Biochem. Eng. J.* *77*, 110–118.
- Swee, M.H., Parks, W.C., and Pierce, R.A. (1995). Developmental regulation of elastin production. Expression of tropoelastin pre-mRNA persists after down-regulation of steady-state mRNA levels. *J. Biol. Chem.* *270*, 14899–14906.
- Chin, K., Wieslander, C., Shi, H., Balgobin, S., Montoya, T.I., Yanagisawa, H., and Word, R.A. (2016). Pelvic organ support in animals with partial loss of fibulin-5 in the vaginal wall. *PLoS ONE* *11*, e0152793.
- Moore, J., and Thibeault, S. (2012). Insights into the role of elastin in vocal fold health and disease. *J. Voice* *26*, 269–275.
- Shapiro, S.D., Endicott, S.K., Province, M.A., Pierce, J.A., and Campbell, E.J. (1991). Marked longevity of human lung parenchymal elastic fibers deduced from prevalence of D-aspartate and nuclear weapons-related radiocarbon. *J. Clin. Invest.* *87*, 1828–1834.
- Dayal, D., Giri, D., and Senniappan, S. (2017). A rare association of central hypothyroidism and adrenal insufficiency in a boy with Williams-Beuren syndrome. *Ann. Pediatr. Endocrinol. Metab.* *22*, 65–67.
- Strømme, P., Bjørnstad, P.G., and Ramstad, K. (2002). Prevalence estimation of Williams syndrome. *J. Child Neurol.* *17*, 269–271.
- Urbán, Z., Riaz, S., Seidl, T.L., Katahira, J., Smoot, L.B., Chitayat, D., Boyd, C.D., and Hinek, A. (2002). Connection between elastin haploinsufficiency and increased cell proliferation in patients with supravalvular aortic stenosis and Williams-Beuren syndrome. *Am. J. Hum. Genet.* *71*, 30–44.
- Zhang, P., Huang, A., Ferruzzi, J., Mecham, R.P., Starcher, B.C., Tellides, G., Humphrey, J.D., Giordano, F.J., Niklason, L.E., and Sessa, W.C. (2012). Inhibition of microRNA-29 enhances elastin levels in cells haploinsufficient for elastin and in bioengineered vessels—brief report. *Arterioscler. Thromb. Vasc. Biol.* *32*, 756–759.
- Pober, B.R. (2010). Williams-Beuren syndrome. *N. Engl. J. Med.* *362*, 239–252.
- Li, D.Y., Brooke, B., Davis, E.C., Mecham, R.P., Sorensen, L.K., Boak, B.B., Eichwald, E., and Keating, M.T. (1998). Elastin is an essential determinant of arterial morphogenesis. *Nature* *393*, 276–280.
- Milewicz, D.M., Urbán, Z., and Boyd, C. (2000). Genetic disorders of the elastic fiber system. *Matrix Biol.* *19*, 471–480.
- Wise, S.G., and Weiss, A.S. (2009). Tropoelastin. *Int. J. Biochem. Cell Biol.* *41*, 494–497.
- Warren, L., Manos, P.D., Ahfeldt, T., Loh, Y.H., Li, H., Lau, F., Ebina, W., Mandal, P.K., Smith, Z.D., Meissner, A., et al. (2010). Highly efficient reprogramming to pluripotency and directed differentiation of human cells with synthetic modified mRNA. *Cell Stem Cell* *7*, 618–630.
- Zangi, L., Lui, K.O., von Gise, A., Ma, Q., Ebina, W., Ptaszek, L.M., Später, D., Xu, H., Tabebordbar, M., Gorbатов, R., et al. (2013). Modified mRNA directs the fate of heart progenitor cells and induces vascular regeneration after myocardial infarction. *Nat. Biotechnol.* *31*, 898–907.

19. Kormann, M.S., Hasenpusch, G., Aneja, M.K., Nica, G., Flemmer, A.W., Herber-Jonat, S., Huppmann, M., Mays, L.E., Illenyi, M., Schams, A., et al. (2011). Expression of therapeutic proteins after delivery of chemically modified mRNA in mice. *Nat. Biotechnol.* 29, 154–157.
20. Stadler, C.R., Bähr-Mahmud, H., Celik, L., Hebich, B., Roth, A.S., Roth, R.P., Karikó, K., Türeci, Ö., and Sahin, U. (2017). Elimination of large tumors in mice by mRNA-encoded bispecific antibodies. *Nat. Med.* 23, 815–817.
21. Steinle, H., Behring, A., Schlensak, C., Wendel, H.P., and Avci-Adali, M. (2017). Concise review: application of in vitro transcribed messenger RNA for cellular engineering and reprogramming: progress and challenges. *Stem Cells* 35, 68–79.
22. Tsamis, A., Krawiec, J.T., and Vorp, D.A. (2013). Elastin and collagen fibre microstructure of the human aorta in ageing and disease: a review. *J. R. Soc. Interface* 10, 20121004.
23. Choi, W.S., Mitsumoto, A., and Kochevar, I.E. (2009). Involvement of reactive oxygen species in TGF-beta1-induced tropoelastin expression by human dermal fibroblasts. *Photochem. Photobiol.* 85, 1425–1433.
24. Kucich, U., Rosenbloom, J.C., Abrams, W.R., and Rosenbloom, J. (2002). Transforming growth factor-beta stabilizes elastin mRNA by a pathway requiring active Smads, protein kinase C-delta, and p38. *Am. J. Respir. Cell Mol. Biol.* 26, 183–188.
25. Kähäri, V.M., Olsen, D.R., Rhudy, R.W., Carrillo, P., Chen, Y.Q., and Uitto, J. (1992). Transforming growth factor-beta up-regulates elastin gene expression in human skin fibroblasts. Evidence for post-transcriptional modulation. *Lab. Invest.* 66, 580–588.
26. Kothapalli, C.R., and Ramamurthi, A. (2010). Induced elastin regeneration by chronically activated smooth muscle cells for targeted aneurysm repair. *Acta Biomater.* 6, 170–178.
27. Uitto, J., Hsu-Wong, S., Katchman, S.D., Bashir, M.M., and Rosenbloom, J. (1995). Skin elastic fibres: regulation of human elastin promoter activity in transgenic mice. *Ciba Found. Symp.* 192, 237–253, discussion 253–258.
28. Katchman, S.D., Hsu-Wong, S., Ledo, I., Wu, M., and Uitto, J. (1994). Transforming growth factor-beta up-regulates human elastin promoter activity in transgenic mice. *Biochem. Biophys. Res. Commun.* 203, 485–490.
29. Mizuno, T., Yau, T.M., Weisel, R.D., Kiani, C.G., and Li, R.K. (2005). Elastin stabilizes an infarct and preserves ventricular function. *Circulation* 112 (Suppl 9), 181–188.
30. Xiong, J., Wang, S.M., Chen, L.H., Lin, Y., Zhu, Y.F., and Ye, C.S. (2008). Elastic fibers reconstructed using adenovirus-mediated expression of tropoelastin and tested in the elastase model of abdominal aortic aneurysm in rats. *J. Vasc. Surg.* 48, 965–973.
31. Xie, H., Lucchesi, L., Zheng, B., Ladich, E., Pineda, T., Merten, R., Gregory, C., Rutten, M., and Gregory, K. (2017). Treatment of burn and surgical wounds with recombinant human tropoelastin produces new elastin fibers in scars. *J. Burn Care Res.* 38, e859–e867.
32. Akhurst, R.J., and Derynck, R. (2001). TGF-beta signaling in cancer—a double-edged sword. *Trends Cell Biol.* 11, S44–S51.
33. Welch, D.R., Fabra, A., and Nakajima, M. (1990). Transforming growth factor beta stimulates mammary adenocarcinoma cell invasion and metastatic potential. *Proc. Natl. Acad. Sci. USA* 87, 7678–7682.
34. Saxena, M., Van, T.T., Baird, F.J., Coloe, P.J., and Smooker, P.M. (2013). Pre-existing immunity against vaccine vectors—friend or foe? *Microbiology* 159, 1–11.
35. Vannucci, L., Lai, M., Chiuppesi, F., Ceccherini-Nelli, L., and Pistello, M. (2013). Viral vectors: a look back and ahead on gene transfer technology. *New Microbiol.* 36, 1–22.
36. Blanchevoe, C., Floquet, N., Scandolera, A., Baud, S., Maurice, P., Bocquet, O., Blaise, S., Ghoneim, C., Cantarelli, B., Delacoux, F., et al. (2013). Interaction between the elastin peptide VGVAPG and human elastin binding protein. *J. Biol. Chem.* 288, 1317–1328.
37. Hinek, A. (1994). Nature and the multiple functions of the 67-kD elastin-/laminin binding protein. *Cell Adhes. Commun.* 2, 185–193.
38. Xu, J., and Shi, G.-P. (2014). Vascular wall extracellular matrix proteins and vascular diseases. *Biochim. Biophys. Acta* 1842, 2106–2119.
39. Ito, S., Ishimaru, S., and Wilson, S.E. (1997). Inhibitory effect of type 1 collagen gel containing alpha-elastin on proliferation and migration of vascular smooth muscle and endothelial cells. *Cardiovasc. Surg.* 5, 176–183.
40. Karikó, K., Muramatsu, H., Ludwig, J., and Weissman, D. (2011). Generating the optimal mRNA for therapy: HPLC purification eliminates immune activation and improves translation of nucleoside-modified, protein-encoding mRNA. *Nucleic Acids Res.* 39, e142.
41. Weissman, D., Pardi, N., Muramatsu, H., and Karikó, K. (2013). HPLC purification of in vitro transcribed long RNA. In *Synthetic Messenger RNA and Cell Metabolism Modulation*, P.M. Rabinovich, ed. (Springer), pp. 43–54.
42. Thess, A., Grund, S., Mui, B.L., Hope, M.J., Baumhof, P., Fotin-Mleczek, M., and Schlake, T. (2015). Sequence-engineered mRNA without chemical nucleoside modifications enables an effective protein therapy in large animals. *Mol. Ther.* 23, 1456–1464.
43. Flynn, P.M., Shenep, J.L., Mao, L., Crawford, R., Williams, B.F., and Williams, B.G. (1994). Influence of needle gauge in Mantoux skin testing. *Chest* 106, 1463–1465.
44. Fritze, O., Schleicher, M., König, K., Schenke-Layland, K., Stock, U., and Harasztsi, C. (2010). Facilitated noninvasive visualization of collagen and elastin in blood vessels. *Tissue Eng. Part C Methods* 16, 705–710.
45. Avci-Adali, M., Behring, A., Steinle, H., Keller, T., Krajewski, S., Schlensak, C., and Wendel, H.P. (2014). In vitro synthesis of modified mRNA for induction of protein expression in human cells. *J. Vis. Exp.* (93), e51943.


**Supplemental Information**

***De Novo* Synthesis of Elastin by Exogenous  
Delivery of Synthetic Modified mRNA into Skin  
and Elastin-Deficient Cells**

**Mario Lescan, Regine Mariette Perl, Sonia Golombek, Martin Pilz, Ludmilla Hann, Mahua Yasmin, Andreas Behring, Timea Keller, Andrea Nolte, Franziska Gruhn, Efrat Kochba, Yotam Levin, Christian Schlensak, Hans Peter Wendel, and Meltem Avcı-Adalı**

## Supplementary Data:

### Detection of elastin using dot blot assay




**Supplementary Figure 1: Dot blot images with detected elastin after the transfection of EA.hy296 cells, human fibroblasts, or MSCs derived from WBS patient (WBS\_MSCs) with synthetic modified TE mRNA.**  $3 \times 10^5$  EA.hy926 cells, human fibroblasts, or WBS\_MSCs were transfected with 2.5  $\mu\text{g}$  synthetic modified TE mRNA. As a control, cells were incubated with OptiMEM containing Lipofectamine 2000 without TE mRNA. After 24 h, 500  $\mu\text{l}$  supernatant was blotted on nitrocellulose membrane and elastin was detected using polyclonal antibody to human aortic elastin and the subsequent detection of alkaline phosphatase conjugated secondary antibody.

### Elastin ELISA

For the detection of translated elastin in the supernatant of mRNA transfected cells,  $4 \times 10^5$  EAHy.926 cells were seeded, cultivated overnight, and transfected with 2.5  $\mu\text{g}$   $\Psi/5\text{mCTP}$ ,  $\text{m1}\Psi$  or  $\text{m1}\Psi/5\text{mCTP}$  modified TE mRNA using Lipofectamine 2000. After 4 hr, transfection complexes were removed and 1 ml medium was added. The


supernatant was collected after 24 hr and centrifuged 10 min at 3,000 x g to remove cell debris. Elastin concentration was measured using ELISA Kit for human elastin (Antibodies-online, Cloud-Clone Corp., Aachen, Germany) according to manufacturer's protocol. Cells incubated with OptiMEM containing Lipofectamine 2000 was used as control.



**Supplementary Figure 2: Detection of elastin concentration after the transfection of EA.hy293 cells with Ψ/5mCTP, m1Ψ or m1Ψ/5mCTP modified TE mRNA using elastin ELISA.**  $4 \times 10^5$  EA.hy926 cells were transfected with 2.5  $\mu$ g differently modified synthetic TE mRNA. After 24 hr, the elastin concentration was detected using elastin ELISA. As a control, cells were incubated with OptiMEM containing Lipofectamine 2000 without TE mRNA. Results are shown as mean  $\pm$  SEM (n=4). Statistical differences were determined using one-way ANOVA with Bonferroni's multiple comparisons test. (\*p<0.05, \*\*p<0.01)

### **Detection of elastin by two-photon laser scanning microscopy (LSM)**

To analyze the elastin fiber formation,  $3 \times 10^5$  fibroblasts were transfected with 2.5  $\mu\text{g}$   $\Psi$ /5mCTP modified TE mRNA and 30,000 cells were seeded on each 1  $\text{cm}^3$  gelatin scaffold (Gelfoam<sup>®</sup>, Pfizer, New York, NY, USA). After a cultivation for 21 days at 37°C and 5%  $\text{CO}_2$ , the elastin fiber formation was analyzed using two-photon LSM as described in the study by Fritze et al. [1]. Therefore, Mira<sup>TM</sup> 900 Ti:Sapphire Laser pumped by a Verdi V5 Diode-Pumped Laser (Coherent, Santa Clara, CA, USA) coupled to an upright Zeiss Axioskop 2 FS MOT motor-based confocal laser-scanning microscope was used. After excitation with 760 nm light, elastin autofluorescence was detected as a broad emission signal from 403 to 531 nm with a maximum at approximately 457 nm. At an excitation wavelength of 760 nm, the scaffolds seeded with TE mRNA transfected cells revealed an increased autofluorescence signal compared to the scaffolds seeded with cells without TE mRNA transfection.



**Supplementary Figure 3: Analysis of elastin synthesis by using two-photon laser scanning microscopy (LSM).** 30,000 human fibroblasts without or with 2.5  $\mu\text{g}$  TE mRNA transfection were seeded on gelatin scaffolds, and cultivated for 21 days. **A)** Two-photon LSM images of scaffolds seeded with or without TE mRNA transfected fibroblasts at an excitation wavelength of 760 nm. **B)** Spectral distribution measured by setting the excitation wavelength to 760 nm. The crosses (+) show the analyzed region of interests (ROIs).

## Literature

[1] O. Fritze, M. Schleicher, K. König, K. Schenke-Layland, U. Stock, C. Harasztosi, Facilitated noninvasive visualization of collagen and elastin in blood vessels, *Tissue Eng Part C Methods* 16(4) (2010) 705-10.