

Human trophoblast stem cell self-renewal and differentiation: Role of decorin

Pinki Nandi¹, Hyobin Lim¹, Eloy Jose Torres-Garcia¹, Peeyush K Lala^{1,2,3,*}

¹Department of Anatomy and Cell Biology, Schulich School of Medicine and Dentistry, University of Western Ontario, London, Ontario, Canada.

² Department of Oncology, Schulich School of Medicine and Dentistry, University of Western Ontario, London, Ontario, Canada.

³ Associate Scientist, Children's Health Research Institute, Schulich School of Medicine and Dentistry, University of Western Ontario, London, Ontario, Canada.

*Corresponding Author (Correspondence to pklala@uwo.ca)

Supplementary Figure 1




Figure legend for supplementary figure 1: S1a shows the negative controls for SOX2 (left panel) and NANOG (right panel) in HTR-monolayer. Figure S1b shows the negative controls for SOX2 (left panel) and NANOG (right panel) in HTR-spheroids. Figure S1c shows staining for OCT4 in HTR monolayer and spheroids.

Supplementary Figure 2




Figure legend for supplementary figure 2: qRT-PCR analysis of ES (SOX2, NANOG), TS (CDX2, EOMES) and differentiation markers (CGB and HLA-G) in a population of pooled HTR-spheroids.