

Web appendix 3: Results of sensitivity analyses for the PSM-COPD study

	Baseline		12 months			
	Telephone PSM mean (sd)	Usual Care mean (sd)	Telephone PSM mean (sd)	Usual Care mean (sd)	Mean difference ¹ (95% CI)	p-value
Per-protocol analysis						
SGRQ-C total	26.5 (13.5)	29.5 (14.5)	27.6 (14.6)	30.8 (17.0)	-1.4 (-3.6, 0.8)	0.2
SGRQ-C symptom	48.1 (21.1)	47.8 (20.7)	49.2 (21.2)	50.0 (22.7)	-1.9 (-4.9, 1.2)	0.2
SGRQ-C activity	34.1 (19.8)	38.7 (21.3)	33.9 (20.1)	39.3 (24.4)	-2.4 (-5.6, 0.8)	0.1
SGRQ-C impact	14.4 (12.3)	17.5 (13.9)	15.8 (13.8)	19.3 (15.6)	-1.6 (-3.8, 0.5)	0.1
Analysis to assess impact of assessment window						
SGRQ-C total	27.1 (13.9)	29.1 (14.2)	28.1 (15.5)	30.7 (16.8)	-1.1 (-3.3, 1.1)	0.3
SGRQ-C symptom	48.6 (21.2)	47.7 (20.7)	49.3 (21.4)	50.0 (22.6)	-1.6 (-4.7, 1.4)	0.3
SGRQ-C activity	35.0 (19.9)	38.5 (21.1)	34.4 (20.7)	39.1 (24.2)	-2.5 (-5.6, 0.7)	0.1
SGRQ-C impact	14.8 (12.8)	17.0 (13.4)	16.4 (14.8)	19.2 (15.3)	-1.1 (-3.3, 1.0)	0.3
Imputation to assess impact of missing data						
SGRQ-C total	27.9 (14.5)	29.5 (14.5)	28.5 (15.9)	31.1 (17.3)	-1.4 (-3.4, 0.6)	0.2
SGRQ-C symptom	48.3 (21.8)	47.7 (20.5)	48.6 (21.1)	50.2 (22.4)	-2.0 (-4.7, 0.7)	0.1
SGRQ-C activity	36.3 (21.0)	38.5 (21.2)	34.7 (21.4)	39.2 (23.8)	-3.0 (-5.9, -0.2)	0.04
SGRQ-C impact	15.5 (13.4)	17.6 (13.8)	17.5 (15.3)	19.9 (15.5)	-0.9 (-2.9, 1.1)	0.4

¹Telephone PSM compared to usual care (negative values favour Telephone PSM).