

S1 Table. Baseline parameters of patients with liver cirrhosis compared with *Candida* DNA-positive and -negative duodenal samples.

Parameter	DNA-positive duodenal samples (n=44), median (range)	DNA-negative duodenal samples (n=10), median (range)	Level of significance (p)
MELD score, median (range)	14 (6-40)	12 (5-20)	0.242
Age (years), median (range)	61 (29-84)	57 (32-65)	0.063
Albumin (g/l), median (range)	35.4 (21-46)	31.3 (29-47)	0.675
Bilirubin ($\mu\text{mol/l}$), median (range)	36 (7-286)	35.6 (10-82)	0.672
INR, median (range)	1.3 (1-2)	1.35 (1-2)	0.927
Hemoglobin (mmol/l), median (range)	6.8 (5-9)	7 (5-8)	0.674
ALAT ($\mu\text{kat/l}$), median (range)	0.5 (0-3)	0.6 (0-2)	0.983
GGT ($\mu\text{kat/l}$), median (range)	1.6 (0-22)	1.6 (1-25)	0.749
Serum creatinine, ($\mu\text{mol/l}$), median (range)	87 (29-591)	67 (48-114)	0.243
GFR (ml/min), median (range)	80.3 (8-150)	91.5 (44-123)	0.364
White blood cell count ($\text{exp}9/\text{l}$), median (range)	5.75 (2-13)	5.4 (2-11)	0.779
C-reactive protein (mg/dl), median (range)	17.8 (1-104)	15.6 (1-44)	0.723

GFR = glomerular filtration rate; INR = international normalized ratio; ALAT = Aspartat-Amino-Transferase; GGT = Gamma-glutamyltransferase.