

S3 Table. Association between drug therapy and the presence and quantification of Candida DNA in duodenal fluid.

| Drug | Presence of fungal DNA (OR (IQR)) | Level of significance (p) | Median DNA quantification (range) | Level of significance (p) |
|----------------------------------|--|---------------------------|--|---------------------------|
| Antibiotic treatment (yes/no) | 89.5% vs. 77.1% 2.5 (0.477–13.300) | 0.277 | 2.8x10⁶ (2.2x10³–4.4x10⁸) vs. 3.2x10⁴ (2.3x10²–6.3x10⁹) | 0.001 |
| Betablocker (yes/no) | 80.0% vs. 84.2% 0.704 (0.302–5.889) | 0.737 | 2.5x10 ⁵ (2.3x10 ² –6.3x10 ⁹) vs. 1.3x10 ⁶ (8.8x10 ² –4.1x10 ⁹) | 0.390 |
| Protone pump inhibitors (yes/no) | 89.3% vs. 73.1% 0.137 (0.74–1.428) | 0.169 | 5.0x10 ⁵ (2.3x10 ² –4.1x10 ⁹) vs. 1.2x10 ⁵ (7.8x10 ² –6.3x10 ⁹) | 0.818 |

OR = odds ratio; IQR = interquartile range