

Tobacco-smoking-related prevalence of methanogens in the oral fluid microbiota

Ghiles Grine¹, Elodie Terror^{1,2}, Mahmoud Abdelwadoud Boualam¹, Gérard Aboudharam^{1,2},

Hervé Chaudet¹, Raymond Ruimy³, Michel Drancourt^{1*}

SUPPLEMENTARY MATERIAL:

Supplementary text: English translation of the anonymized questionnaire filled by all the participants to this study.

Patient questionnaire

PATIENT INFORMATION

General data

Last name and first name:

Birth date:

Sex:

Age:

Profession:

Medical background:

Followed treatments:

Dental status:

Periodontal data

	NO	Yes
Previous periodontal treatment:	<input type="checkbox"/>	<input type="checkbox"/>
Presence of inflammation:	<input type="checkbox"/>	<input type="checkbox"/>
Presence of dental plaque:	<input type="checkbox"/>	<input type="checkbox"/>
Presence of tartar:	<input type="checkbox"/>	<input type="checkbox"/>
Presence of recession:	<input type="checkbox"/>	<input type="checkbox"/>
Presence of mobility:	<input type="checkbox"/>	<input type="checkbox"/>
Presence of periodontal abscess:	<input type="checkbox"/>	<input type="checkbox"/>
Loss of attachment:	<input type="checkbox"/>	<input type="checkbox"/>
Presence of implants:	<input type="checkbox"/>	<input type="checkbox"/>