

Supplementary Information

Liver X receptors regulate hepatic F4/80⁺CD11b⁺ Kupffer cells/macrophages and innate immune responses in mice

Kaori Endo-Umeda¹, Hiroyuki Nakashima², Shihoko Komine-Aizawa³, Naoki Umeda¹, Shuhji Seki², and Makoto Makishima^{1,*}

¹Division of Biochemistry, Department of Biomedical Sciences, Nihon University School of Medicine, 30-1 Oyaguchi-kamicho, Itabashi-ku, Tokyo 173-8610, Japan

²Department of Immunology and Microbiology, National Defense Medical College, 3-2 Namiki, Tokorozawa, Saitama 359-8513, Japan

³Division of Microbiology, Department of Pathology and Microbiology, Nihon University School of Medicine, 30-1 Oyaguchi-kamicho, Itabashi-ku, Tokyo 173-8610, Japan


*Corresponding author. e-mail: makishima.makoto@nihon-u.ac.jp

F4/80 (green)

CD11b (red)

WT

LXR α / β -KO



Supplementary Figure 1. Double immunostaining of F4/80 (green) and CD11b (red) in the liver of WT and LXR α / β -KO mice. The open triangles indicate F4/80⁺CD11b⁺ cells. Scale bar = 50 μ m.