

Table S5. Relationship between best response and changes in serum level of CEA and CA 19-9 before start of treatment to cycle 4 (8 weeks of treatment) in patients with elevated CEA ($\geq 5\mu\text{g/L}$) or CA 19-9 ($\geq 35\text{ kU/L}$) at baseline

Patients with baseline CEA $\geq 5\ \mu\text{g/L}$ (<i>n</i> =384)		Change in serum level of CEA		Patients with baseline CA 19-9 $\geq 35\ \text{kU/L}$ (<i>n</i> =260)		Change in serum level of CA 19-9	
		Drop	Rise			Drop	Rise
Clinical response	<i>n</i> (%)	<i>n</i> (%)	<i>n</i> (%)	Clinical response	<i>n</i> (%)	<i>n</i> (%)	<i>n</i> (%)
CR+PR	187 (49)	177 (95)	10 (5)	CR+PR	130 (50)	124 (95)	6 (5)
SD	170 (44)	128 (75)	42 (25)	SD	110 (42)	87 (79)	23 (21)
PD	20 (5)	6 (30)	14 (70)	PD	16 (6)	8 (50)	8 (50)
NE	7 (2)			NE	4 (2)		

Abbreviations: CR, Complete response; NE, Not evaluable; PD, Progressive disease; PR, Partial response; SD, Stable disease.