

## **Supporting Information**

### **Extracellular Vesicles as Markers and Mediators in Sepsis**

Pierre Raeven, MD, PhD<sup>1,2\*</sup>

Johannes Zipperle, MSc.<sup>1</sup>

Susanne Drechsler, DVM<sup>1</sup>

1. Ludwig Boltzmann Institute for Experimental and Clinical Traumatology, Vienna, Austria.
2. Department of Anesthesiology, General Intensive Care Medicine and Pain Management, Medical University of Vienna, Vienna, Austria.

**\*Author to whom correspondence should be addressed:**

Pierre Raeven, Email: [pierre.raeven@meduniwien.ac.at](mailto:pierre.raeven@meduniwien.ac.at)

**Table S1. Important markers of extracellular vesicles derived from various cell types.**

| <b>Cell type</b>     | <b>Markers</b>  |
|----------------------|---|
| Total cells          | PhtdSer   |
| Platelets            | Fibrinogen, CD61, CD42a, CD9, CD63, CD81, CD41, CD41a, PAC-1, CD62P |
| ECs                  | CD62e, CD144, CD146, CD105, CD106, TM, TF, EPCR, CD31, CD142        |
| Myeloid cell lineage | CD13  |
| Leukocytes           | CD45, CD11a   |
| Granulocytes         | CD65b, CD66b, PKH-26, CD16b, CD11b                                  |
| Monocytes            | CD14, CD15 CD11b, CD142   |
| Erythrocytes         | CD235a (GlycophorinA)   |
| Lymphocytes          | CD3, CD19, CD62L  |

APC: activated protein C; CD: cluster of differentiation; EC: endothelial cell; EPCR: endothelial protein C receptor; PhtdSer: phosphatidylserine; TF: tissue factor; TM: thrombomodulin.