## **Supplementary Data**

## Supplementary Table S1. Baseline Characteristics According to Annualized Continuous Glucose Monitoring Use

	Annualized CGM use			
	0–2 days per week (n=24)	3–5 days per week (n=54)	6–7 days per week (n=42)	P*
Age (years)	12.2±2.9	13.0±2.5	12.7 ± 2.7	0.53
Diabetes duration (years)	$6.0 \pm 3.5$	$5.9 \pm 3.4$	$6.4 \pm 3.8$	0.81
Sex, male	63%	39%	60%	0.83
Race, white	96%	91%	100%	0.28
BMI z-score (SDS)	$0.45 \pm 0.99$	$0.63 \pm 1.01$	$0.38 \pm 0.92$	0.44
Insulin regimen, pump	83%	85%	83%	0.96
Blood glucose monitoring frequency (times per day)	$6.4 \pm 2.7$	$6.3 \pm 2.5$	$7.3 \pm 2.1$	0.09
$HbA_{1c}(\%)$	$8.1 \pm 0.6$	$8.0 \pm 0.8$	$7.9 \pm 0.9$	0.74
Family structure, two-parent	96%	89%	95%	0.87
Household income, >\$100K	39%	45%	68%	0.02
Parental education, college or above	67%	72%	76%	0.41

\*P-value for analysis of variance or Mantel-Haenszel Chi-square tests. BMI, body mass index; HbA $_{\rm Ic}$ , glycated hemoglobin; SDS, standard deviation score.