

Supporting Information

Novel Topologically Complex Scaffold Derived from Alkaloid Haemanthamine

Karthik Govindaraju ¹, Marco Masi ², Margaux Colin³, Veronique Mathieu³, Antonio Evidente², Todd W. Hudnall ^{1,*} and Alexander Kornienko ^{1,*}


¹ Department of Chemistry and Biochemistry, Texas State University, San Marcos, TX 78666

² Dipartimento di Scienze Chimiche, Università di Napoli Federico II, Complesso Universitario Monte Sant'Angelo, Via Cintia 4, 80126, Napoli, Italy


³ Department of Pharmacotherapy and Pharmaceutics, Faculté de Pharmacie, Université Libre de Bruxelles (ULB), Brussels, Belgium

NMR Spectra


1: ^1H NMR (400 MHz, CDCl_3)




1: ^{13}C NMR (100 MHz, CDCl_3)




2: ^1H NMR (400 MHz, CDCl_3)




3: ^1H NMR (400 MHz, CDCl_3)




4: ^1H NMR (400 MHz, CDCl_3)




4: ^{13}C NMR (100 MHz, CDCl_3)




5: ^1H NMR (400 MHz, CDCl_3)




5: ^{13}C NMR (100 MHz, CDCl_3)




6: ¹H NMR (400 MHz, CDCl₃)




6: ¹³C NMR (100 MHz, CDCl₃)



7: ^1H NMR (400 MHz, CDCl_3)



7: ^{13}C NMR (100 MHz, CDCl_3)



9: ^1H NMR (400 MHz, CDCl_3)

