

Co-occurrence network reveals a higher fragmentation of the
bacterial community in Kaidu River than its tributaries in
Northwestern China

Yang Hu^{1,2}, Chengrong Bai^{1,2}, Jian Cai^{1,2}, Jiangyu Dai³, Keqiang Shao¹, Xiangming
Tang¹, Guang Gao^{1,*}

¹ State Key Laboratory of Lake Science and Environment, Nanjing Institute of
Geography and Limnology, Chinese Academy of Sciences, Nanjing 210008, China

² University of Chinese Academy of Sciences, Beijing, 100000, China

³ State Key Laboratory of Hydrology-Water Resources and Hydraulic Engineering,
Nanjing Hydraulic Research Institute, Nanjing, 210029, P.R. China

Running headline: Bacterial community in lotic ecosystems

* Corresponding author: Guang Gao, Nanjing Institute of Geography and Limnology, Chinese Academy
of Sciences, 73 East Beijing Road, Nanjing 210008, China

E-mail: guanggao@niglas.ac.cn

Phone: (+86) 25 86882187

Fax: (86) 25 57714759




Fig. S1 The rarefaction curve of bacterial community in the Kaidu River and its tributaries. KD represents Kaidu River, TR represents its tributaries




Fig. S2 The dendrogram of the bacterial communities in the Kaidu River and its tributaries based on UPGMA algorithm





Fig. S3 The relationships between Bray-Curtis dissimilarity and geographic distance in the Kaidu River and its tributaries