

Supplementary Information

Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust

Harry M.T. Choi,¹ Maayan Schwarzkopf,¹ Mark E. Fornace,² Aneesh Acharya,¹
Georgios Artavanis,¹ Johannes Stegmaier,^{3,4,5} Alexandre Cunha,^{3,6} and Niles A. Pierce^{1,7,8,*}

Contents

S1 Additional materials and methods	6
S1.1 Probe set and amplifier details	6
S1.2 Confocal microscope settings	8
S1.3 Flow cytometer settings	9
S1.4 Image analysis	11
S1.4.1 Raw pixel intensities	11
S1.4.2 Measurement of signal, background, and signal-to-background	11
S1.4.3 Measurement of background components	12
S1.4.4 Measurement of split-initiator HCR suppression	12
S1.4.5 Normalized voxel intensities for qHCR imaging: analog mRNA relative quantitation with subcellular resolution in an anatomical context	14
S1.4.6 Dot detection and colocalization for dHCR imaging: digital mRNA absolute quantitation in an anatomical context	14
S1.5 Flow cytometry data analysis	17
S1.5.1 Raw cell intensities	17
S1.5.2 Measurement of signal, background, signal-to-background, background components, and split-initiator HCR suppression for transgenic targets	17
S1.5.3 Measurement of signal, background, signal-to-background, background components, and split-initiator HCR suppression for endogenous targets	17
S1.5.4 Normalized single-cell intensities for qHCR flow cytometry: analog mRNA relative quantitation for high-throughput expression profiling of human and bacterial cells	18
S2 Protocols for in situ HCR v3.0	19
S2.1 Protocols for whole-mount chicken embryos	19
S2.1.1 Preparation of fixed whole-mount chicken embryos	19
S2.1.2 Buffer recipes for sample preparation	20
S2.1.3 Multiplexed in situ HCR v3.0 using split-initiator probes	21
S2.1.4 Buffer recipes for in situ HCR v3.0	22
S2.1.5 Sample mounting for microscopy	23
S2.1.6 Reagents and supplies	24

¹Division of Biology & Biological Engineering, California Institute of Technology, Pasadena, CA 91125, USA. ²Division of Chemistry & Chemical Engineering, California Institute of Technology, Pasadena, CA 91125, USA. ³Center for Advanced Methods in Biological Image Analysis, Beckman Institute, California Institute of Technology, Pasadena, CA 91125, USA. ⁴Institute for Automation & Applied Informatics, Karlsruhe Institute of Technology, Karlsruhe, Germany. ⁵Institute of Imaging & Computer Vision, RWTH Aachen University, Aachen, Germany. ⁶Center for Data-Driven Discovery, California Institute of Technology, Pasadena, CA 91125, USA. ⁷Division of Engineering & Applied Science, California Institute of Technology, Pasadena, CA 91125, USA. ⁸Weatherall Institute of Molecular Medicine, University of Oxford, Oxford OX3 9DS, UK. *Email: niles@caltech.edu

S2.2	Protocols for mammalian cells on a chambered slide	25
S2.2.1	Preparation of fixed mammalian cells on a chambered slide	25
S2.2.2	Buffer recipes for sample preparation	26
S2.2.3	Multiplexed in situ HCR v3.0 using split-initiator probes	27
S2.2.4	Buffer recipes for in situ HCR v3.0	28
S2.2.5	Reagents and supplies	29
S2.3	Protocols for mammalian cells in suspension	30
S2.3.1	Preparation of fixed mammalian cells in suspension	30
S2.3.2	Buffer recipes for sample preparation	31
S2.3.3	Multiplexed in situ HCR v3.0 using split-initiator probes	32
S2.3.4	Buffer recipes for in situ HCR v3.0	34
S2.3.5	Reagents and supplies	35
S2.4	Protocols for bacteria in suspension	36
S2.4.1	Preparation of fixed bacteria in suspension	36
S2.4.2	Buffer recipes for sample preparation	37
S2.4.3	Multiplexed in situ HCR v3.0 using split-initiator probes	38
S2.4.4	Buffer recipes for in situ HCR v3.0	40
S2.4.5	Reagents and supplies	41
S3	Additional studies	42
S3.1	Validation of split-initiator HCR suppression in vitro and in situ (cf. Figure 2)	42
S3.2	In situ validation of automatic background suppression with split-initiator probes in whole-mount chicken embryos (cf. Figure 3)	44
S3.2.1	Measurement of background and signal-to-background for unoptimized standard and split-initiator probe sets as a function of probe set size	45
S3.2.2	Measurement of background and signal-to-background for standard and split-initiator probes with identical target-binding domains	49
S3.2.3	Measurement of signal, background, signal-to-background, background components, and split-initiator HCR suppression	51
S3.3	Multiplexed 4-channel mRNA imaging with high signal-to-background in whole-mount chicken embryos (cf. Figure 4)	53
S3.4	qHCR imaging: analog mRNA relative quantitation with subcellular resolution in whole-mount chicken embryos (cf. Figure 5)	55
S3.4.1	Testing for a crowding effect	55
S3.4.2	Redundant 2-channel detection of <i>Dmbx1</i>	60
S3.4.3	Redundant 2-channel detection of <i>Epha4</i>	61
S3.5	In situ validation of automatic background suppression with split-initiator probes for mRNA flow cytometry with cultured human and bacterial cells	63
S3.5.1	Measurement of signal, background, signal-to-background, background components, and split-initiator HCR suppression for <i>d2eGFP</i> transgenic target in HEK cells	63
S3.5.2	Measurement of signal, background, signal-to-background, background components, and split-initiator HCR suppression for <i>GAPDH</i> endogenous target in HEK cells	65
S3.5.3	Measurement of signal, background, signal-to-background, background components, and split-initiator HCR suppression for <i>eGFP</i> transgenic target in <i>E. coli</i>	67
S3.6	qHCR flow cytometry: analog mRNA relative quantitation for high-throughput expression profiling of human and bacterial cells (cf. Figure 6)	69
S3.6.1	Testing for a crowding effect	69
S3.6.2	Redundant 2-channel detection of <i>GAPDH</i> endogenous target in HEK cells	70
S3.6.3	Redundant 2-channel detection of <i>PGK1</i> endogenous target in HEK cells	72
S3.6.4	Redundant 2-channel detection of <i>fusA</i> endogenous target in <i>E. coli</i>	74

S3.6.5	Multiplexed 2-channel detection of <i>GAPDH</i> and <i>PGK1</i> endogenous targets in HEK cells . . .	76
S3.6.6	Multiplexed 2-channel detection of <i>fusA</i> and <i>icd</i> endogenous targets in <i>E. coli</i>	78
S3.7	dHCR imaging: digital mRNA absolute quantitation in an anatomical context (cf. Figure 7)	80
S3.7.1	Redundant 2-channel detection of single <i>BRAF</i> mRNAs in HEK cells using in situ HCR v3.0	80
S3.7.2	Redundant 2-channel detection of single <i>Dmbx1</i> mRNAs in whole-mount chicken embryos using in situ HCR v3.0	81
S3.7.3	Redundant 2-channel detection of single <i>kdrl</i> mRNAs in whole-mount zebrafish embryos using in situ HCR v2.0 (Shah <i>et al.</i> , 2016)	82
S4	Probe sequences	83
S4.1	Standard probes for Figures 3, S5, and S7	84
S4.2	Split-initiator probes for Figures 3, S6, S8, S9, and S10	85
S4.3	Standard probes for Figures S9 and S10	86
S4.4	Split-initiator probes for Figure S11	88
S4.5	Split-initiator probes for Figures 4 and S12	89
S4.6	Split-initiator probes for Figures S13–S17	93
S4.7	Split-initiator probes Figures 5, S18, and S19	95
S4.8	Split-initiator probes for Figure 6, S20–S28	99
S4.9	Split-initiator probes for Figures 7, S29, and S30	108

List of Figures

S1	Illustration of gates used for flow cytometry analysis of HEK cells	10
S2	Illustration of gates used for flow cytometry analysis of <i>E. coli</i>	10
S3	Test tube validation of split-initiator HCR suppression for amplifiers B1, B2, B4, and B5 (cf. Figure 2)	42
S4	Test tube validation of split-initiator HCR suppression for amplifier B3 (cf. Figure 2)	43
S5	Measurement of background for unoptimized standard probe sets as a function of probe set size (cf. Figure 3A)	45
S6	Measurement of background for unoptimized split-initiator probe sets as a function of probe set size (cf. Figure 3A)	46
S7	Measurement of signal and background for unoptimized standard probe sets as a function of probe set size (cf. Figure 3B)	47
S8	Measurement of signal and background for unoptimized split-initiator probe sets as a function of probe set size (cf. Figure 3B)	48
S9	Measurement of background for standard and split-initiator probes with identical target-binding domains	49
S10	Measurement of signal and background for standard and split-initiator probes with identical target-binding domains	50
S11	Measurement of signal and background, background components, and split-initiator HCR suppression	51
S12	Measurement of signal and background for multiplexed 4-channel mRNA imaging (cf. Figure 4) . .	53
S13	Comparison of signal intensity distributions for individual and simultaneous imaging of <i>EphA4</i> and <i>Egr2</i>	55
S14	Characterizing signal plus background for <i>EphA4</i> and <i>Egr2</i> in a 2-target experiment.	56
S15	Characterizing signal plus background for <i>EphA4</i> in a 1-target experiment	57
S16	Characterizing signal plus background for <i>Egr2</i> in a 1-target experiment	58
S17	Characterizing background for <i>EphA4</i> and <i>Egr2</i>	59
S18	Redundant 2-channel detection of <i>Dmbx1</i>	60
S19	Redundant 2-channel detection of <i>EphA4</i> (cf. Figure 5)	61
S20	Measurement of signal and background, background components, and split-initiator HCR suppression for <i>d2eGFP</i> transgenic target in HEK cells (cf. Figure 6A)	63

S21	Measurement of signal and background, background components, and split-initiator HCR suppression for <i>GAPDH</i> endogenous target in HEK cells	65
S22	Measurement of signal and background, background components, and split-initiator HCR suppression for <i>eGFP</i> transgenic target in <i>E. coli</i> (cf. Figure 6A)	67
S23	Comparison of signal intensity distributions for individual and multiplexed floHCR of <i>GAPDH</i> and <i>ACTB</i>	69
S24	Measurement of signal and background, background components, and split-initiator HCR suppression for redundant 2-channel detection of <i>GAPDH</i> endogenous target (cf. Figure 6B)	70
S25	Measurement of signal and background, background components, and split-initiator HCR suppression for redundant 2-channel detection of <i>PGK1</i> endogenous target	72
S26	Measurement of signal and background, background components, and split-initiator HCR suppression for redundant 2-channel detection of <i>fusA</i> endogenous target (cf. Figure 6B)	74
S27	Measurement of signal and background, background components, and split-initiator HCR suppression for multiplexed 2-channel detection of <i>GAPDH</i> and <i>PGK1</i> endogenous targets	76
S28	Measurement of signal and background, background components, and split-initiator HCR suppression for multiplexed 2-channel detection of <i>fusA</i> and <i>icd</i> endogenous targets	78
S29	Redundant 2-channel detection of single <i>BRAF</i> mRNAs in HEK cells using in situ HCR v3.0 (cf. Figure 7A)	80
S30	Redundant 2-channel detection of single <i>Dmbx1</i> mRNAs in whole-mount chicken embryos using in situ HCR v3.0 (cf. Figure 7B)	81
S31	Analysis of single-molecule mRNA imaging in whole-mount zebrafish embryos using in situ HCR v2.0	82

List of Tables

S1	Sequences for gel studies (cf. Figure 2)	6
S2	Organisms, target mRNAs, probe sets, amplifiers, and figure numbers for in situ HCR experiments	7
S3	Confocal microscope settings	8
S4	Flow cytometer settings	9
S5	Experiment types for qHCR imaging using in situ HCR v3.0	12
S6	Parameters used for dot detection in dHCR images.	16
S7	Experiment types for flow cytometry using in situ HCR v3.0 with a transgenic target mRNA	17
S8	Experiment types for flow cytometry using in situ HCR v3.0 with an endogenous target mRNA	18
S9	In situ validation of split-initiator HCR suppression	42
S10	Estimated background for standard and split-initiator probes sets as a function of probe set size (cf. Figure 3A)	46
S11	Estimated signal-to-background for standard and split-initiator probes sets as a function of probe set size (cf. Figure 3B)	48
S12	Estimated background for standard and split-initiator probes with identical target-binding domains	49
S13	Estimated signal-to-background for standard and split-initiator probes with identical target-binding domains	50
S14	Estimated signal-to-background, background components, and split-initiator HCR suppression	52
S15	Estimated signal-to-background for multiplexed 4-channel mRNA imaging	54
S16	Estimated signal-to-background for redundant 2-channel detection of <i>Dmbx1</i> and <i>EphA4</i>	62
S17	Estimated signal-to-background, background components, and split-initiator HCR suppression for <i>d2eGFP</i> transgenic target in HEK cells (cf. Figure 6A)	64
S18	Estimated signal-to-background, background components, and split-initiator HCR suppression for <i>GAPDH</i> endogenous target in HEK cells	66
S19	Estimated signal-to-background, background components, and split-initiator HCR suppression for <i>eGFP</i> transgenic target in <i>E. coli</i> (cf. Figure 6A)	68

S20	Estimated signal-to-background, background components, and split-initiator HCR suppression for redundant 2-channel detection of <i>GAPDH</i> endogenous target (cf. Figure 6B)	71
S21	Estimated signal-to-background, background components, and split-initiator HCR suppression for redundant 2-channel detection of <i>PGK1</i> endogenous target	73
S22	Estimated signal-to-background, background components, and split-initiator HCR suppression for redundant 2-channel detection of <i>fusA</i> endogenous target (cf. Figure 6B)	75
S23	Estimated signal-to-background, background components, and split-initiator HCR suppression for multiplexed 2-channel detection of <i>GAPDH</i> and <i>PGK1</i> endogenous targets	77
S24	Estimated signal-to-background, background components, and split-initiator HCR suppression for multiplexed 2-channel detection of <i>fusA</i> and <i>icd</i> endogenous targets	79
S25	Dot colocalization fractions for redundant 2-channel detection of single <i>BRAF</i> mRNAs in HEK cells using in situ HCR v3.0 (cf. Figure 7A)	80
S26	Dot colocalization fractions for redundant 2-channel detection of single <i>Dmbx1</i> mRNAs in whole-mount chicken embryos using in situ HCR v3.0 (cf. Figure 7B)	81
S27	Dot colocalization fractions for redundant 2-channel detection of single <i>kdrl</i> mRNAs in whole-mount zebrafish embryos using in situ HCR v3.0	82

S1 Additional materials and methods

S1.1 Probe set and amplifier details

HCR Amplifier	Oligo	Length (nt)	Sequence (5' to 3')	Figures
B1-Alexa647	I1	36	gAggAgggCAgCAAACgggAAgAgTCTTCCTTTACg	S3
	P1	45	gAggAgggCAgCAAACggAAAgACgTTgTggCTgTTgTAgTTgTA	S3
	P2	45	CgTTCTTCTgCTTgTCggCCATgATTAgAAgAgTCTTCCTTTACg	S3
B2-Alexa647	I1	36	CCTCgTAAATCCTCATCAATCATCCAgtAAACCgCC	S3
	P1	45	CCTCgTAAATCCTCATCAAAAgACgTTgTggCTgTTgTAgTTgTA	S3
	P2	45	CgTTCTTCTgCTTgTCggCCATgATAAATCATCCAgtAAACCgCC	S3
B3-Alexa647	I1	36	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	2, S4
	P1	45	gTCCCTgCCTCTATATCTTTAgACgTTgTggCTgTTgTAgTTgTA	2, S4
	P2	45	CgTTCTTCTgCTTgTCggCCATgATTTCCACTCAACTTTAACCCg	2, S4
B4-Alexa647	I1	36	CCTCAACCTACCTCCAACTCTCACCATATTCgCTTC	S3
	P1	45	CCTCAACCTACCTCCAACAAAgACgTTgTggCTgTTgTAgTTgTA	S3
	P2	45	CgTTCTTCTgCTTgTCggCCATgATATCTCACCATATTCgCTTC	S3
B5-Alexa647	I1	36	CTCACTCCCAATCTCTATCTACCCTACAAATCCAAT	S3
	P1	45	CTCACTCCCAATCTCTATAAAgACgTTgTggCTgTTgTAgTTgTA	S3
	P2	45	CgTTCTTCTgCTTgTCggCCATgATAACTACCCTACAAATCCAAT	S3

Table S1. Sequences for gel studies (cf. Figure 2). Initiator sequence in green, spacer sequence in blue, target-binding sequence in black. In all cases, the Target is: 5'-TACAACTACAACAAGCCACAACgTCTATATCATggCCgACAAgCagAAgAACg-3'.

Organism	Target	Standard probes (v2.0)	Split-initiator probe pairs (v3.0)	HCR amplifier	Amplification time	Figures
<i>G. gallus domesticus</i>	<i>Sox10</i>	5, 10, 20	5, 10, 20	B3-Alexa647	overnight	3, S5, S7
	<i>Sox10</i>			B3-Alexa647	overnight	3, S6, S8
	<i>Sox10</i>	20	20	B3-Alexa647	overnight	S9, S10
	<i>Sox10</i>			B3-Alexa647	overnight	S9, S10
	<i>EphA4</i>			B2-Alexa647	overnight	S11
	<i>FoxD3</i>		12	B4-Alexa488	overnight	4, S12
	<i>Dmbx1</i>		20	B1-Alexa514	overnight	4, S12
	<i>Sox10</i>		20	B3-Alexa546	overnight	4, S12
	<i>EphA4</i>		20	B2-Alexa647	overnight	4, S12
	<i>EphA4</i>		20	B1-Alexa546	overnight	S13, S14, S15
	<i>Egr2</i>		20	B3-Alexa647	overnight	S13, S14, S16
	<i>Dmbx1</i>		20	B1-Alexa546	overnight	S18
	<i>Dmbx1</i>		20	B2-Alexa647	overnight	S18
	<i>EphA4</i>		20	B1-Alexa546	overnight	5, S19
<i>EphA4</i>		20	B2-Alexa647	overnight	5, S19	
<i>H. sapiens sapiens</i>	<i>d2eGFP</i>		12	B3-Alexa594	overnight	6A, S20
	<i>GAPDH</i>		10	B5-Alexa488	overnight	6B, S23, S24
	<i>GAPDH</i>		10	B4-Alexa594	overnight	6B, S1, S21, S24
	<i>ACTB</i>		10	B2-Alexa594	overnight	S23
	<i>PGK1</i>		18	B1-Alexa488	overnight	S25
	<i>PGK1</i>		18	B2-Alexa594	overnight	S25, S27
	<i>GAPDH</i>		10	B4-Alexa488	overnight	S27
<i>E. coli</i>	<i>eGFP</i>		12	B3-Alexa594	overnight	6A, S22
	<i>fusA</i>		18	B3-Alexa488	overnight	6B, S26, S28
	<i>fusA</i>		18	B2-Alexa594	overnight	6B, S2, S26
	<i>icd</i>		20	B1-Alexa594	overnight	S28
<i>H. sapiens sapiens</i>	<i>BRAF</i>		23	B3-Alexa647	45 min	7A, S29
	<i>BRAF</i>		23	B4-Alexa546	45 min	7A, S29
<i>G. gallus domesticus</i>	<i>Dmbx1</i>		25	B1-Alexa594	90 min	7B, S30
	<i>Dmbx1</i>		25	B2-Alexa647	90 min	7B, S30

Table S2. Organisms, target mRNAs, probe sets, amplifiers, and figure numbers for in situ HCR experiments.

S1.2 Confocal microscope settings

Target	Fluorophore	Laser (nm)	Beam Splitter	Filter (nm)	Pixel size ($x \times y \times z \mu\text{m}$)	Voxel size ($x \times y \times z \mu\text{m}$)	Focal planes	Figures
<i>EphA4</i>	Alexa647	633	MBS 488/561/633	650–689	$0.208 \times 0.208 \times 2.7$		1	S11
<i>Sox10</i>	Alexa647	633	MBS 488/561/633	650–699	$0.415 \times 0.415 \times 2.7$		1	3, S5–S10
<i>FoxD3</i>	Alexa488	488	MBS 488/561/633	491–525	$0.664 \times 0.664 \times 4$		1	4, S12
<i>Dmbx1</i>	Alexa514	514	MBS 458/514	546–564	$0.664 \times 0.664 \times 4$		1	4, S12
<i>Sox10</i>	Alexa546	561	MBS 488/561/633	573–612	$0.664 \times 0.664 \times 4$		1	4, S12
<i>EphA4</i>	Alexa647	633	MBS 488/561/633	654–687	$0.664 \times 0.664 \times 4$		1	4, S12
<i>Dmbx1</i>	Alexa546	561	MBS 488/561/633	563–592	$0.208 \times 0.208 \times 2.7$	$2.1 \times 2.1 \times 2.7$	1	5, S18
<i>Dmbx1</i>	Alexa647	633	MBS 488/561/633	650–689	$0.208 \times 0.208 \times 2.7$	$2.1 \times 2.1 \times 2.7$	1	5, S18
<i>EphA4</i>	Alexa546	561	MBS 488/561/633	563–592	$0.208 \times 0.208 \times 2.7$	$2.1 \times 2.1 \times 2.7$	1	5, S19
<i>EphA4</i>	Alexa647	633	MBS 488/561/633	650–689	$0.208 \times 0.208 \times 2.7$	$2.1 \times 2.1 \times 2.7$	1	5, S19
<i>BRAF</i>	Alexa546	561	MBS 405/488/561/640 (T10/R90)	566–623	$0.0624 \times 0.0624 \times 0.42$		17	7A, S29
<i>BRAF</i>	Alexa647	640	MBS 405/488/561/640 (T10/R90)	656–700	$0.0624 \times 0.0624 \times 0.42$		17	7A, S29
—	DAPI	405	MBS 405/488/561/640 (T10/R90)	410–470	$0.0624 \times 0.0624 \times 0.42$		17	S29
<i>Dmbx1</i>	Alexa594	561	MBS 405/488/561/640 (T10/R90)	580–647	$0.099 \times 0.099 \times 0.420$		22	7B, S30
<i>Dmbx1</i>	Alexa647	640	MBS 405/488/561/640 (T10/R90)	645–700	$0.099 \times 0.099 \times 0.420$		22	7B, S30
<i>EphA4</i>	Alexa546	561	MBS 488/561/633	563–592	$0.415 \times 0.415 \times 2.7$	$2.1 \times 2.1 \times 2.7$	1	S14–S17
<i>Egr2</i>	Alexa647	633	MBS 488/561/633	650–689	$0.415 \times 0.415 \times 2.7$	$2.1 \times 2.1 \times 2.7$	1	S14–S17

Table S3. Confocal microscope settings.

S1.3 Flow cytometer settings

Target	Fluorophore	Laser (nm)	Filter	Emission (nm)	Figures
<i>d2eGFP</i>	Alexa594	561	Y2	615/20	6A, S20
<i>GAPDH</i>	Alexa488	488	B1	525/50	6B, S23, S24, S27
<i>GAPDH</i>	Alexa594	561	Y2	615/20	6B, S1, S21, S24
<i>ACTB</i>	Alexa594	561	Y2	615/20	S23
<i>PGK1</i>	Alexa488	488	B1	525/50	S25
<i>PGK1</i>	Alexa594	561	Y2	615/20	S25, S27
<i>eGFP</i>	Alexa594	561	Y2	615/20	6A, S22
<i>fusA</i>	Alexa488	488	B1	525/50	6B, S26, S28
<i>fusA</i>	Alexa594	561	Y2	615/20	6B, S2, S26
<i>icd</i>	Alexa594	561	Y2	615/20	S28

Table S4. Flow cytometer settings.




Figure S1. Illustration of gates used for flow cytometry analysis of HEK cells. (A) Scatter plots for ungated data. Top: side scatter area (SSC-A) vs forward scatter area (FSC-A). Bottom: forward scatter height (FSC-H) vs forward scatter area (FSC-A). (B) Scatter plots after applying one gate. Top: gate on FSC-A vs SSC-A to remove debris and select cells. Bottom: gate on FSC-A vs FSC-H to remove clumps of cells and select single cells. (C) Scatter plots after applying both gates. (D) Signal plus background for ungated (top) and gated (bottom) samples. Target: *GAPDH*. Probe set: 10 split-initiator probe pairs. Amplifier: B4-Alexa594. The depicted gates were used for the SIG+BACK data in Figure S21.




Figure S2. Illustration of gates used for flow cytometry analysis of *E. coli*. (A) Scatter plots for ungated sample (top) and gated sample (bottom): side scatter area (SSC-A) vs forward scatter area (FSC-A). (B) Signal plus background for ungated sample (top) and gated sample (bottom). Target: *fusA*. Probe set: 18 split-initiator probe pairs. Amplifier: B2-Alexa594. The depicted gate was used for the SIG+BACK data in Figure S26.

S1.4 Image analysis

We build on an image analysis framework developed over a series of publications (Choi *et al.*, 2010, 2014, 2016; Trivedi *et al.*, 2018). For convenience, here we provide a self-contained description of the details relevant to the present work.

S1.4.1 Raw pixel intensities

The total fluorescence within a pixel is a combination of signal and background. Fluorescent background (BACK) arises from three sources in each channel:

- autofluorescence (AF): fluorescence inherent to the sample,
- non-specific detection (NSD): probes that bind non-specifically in the sample and subsequently trigger HCR amplification,
- non-specific amplification (NSA): HCR hairpins that bind non-specifically in the sample.

Fluorescent signal (SIG) in each channel corresponds to:

- signal (SIG): probes that hybridize specifically to the target mRNA and subsequently trigger HCR amplification.

For pixel i of replicate embryo n , we denote the background

$$X_{n,i}^{\text{BACK}} = X_{n,i}^{\text{NSD}} + X_{n,i}^{\text{NSA}} + X_{n,i}^{\text{AF}}, \quad (\text{S1})$$

the signal:

$$X_{n,i}^{\text{SIG}}, \quad (\text{S2})$$

and the total fluorescence (SIG+BACK):

$$X_{n,i}^{\text{SIG+BACK}} = X_{n,i}^{\text{SIG}} + X_{n,i}^{\text{BACK}}. \quad (\text{S3})$$

S1.4.2 Measurement of signal, background, and signal-to-background

For each target mRNA, background (BACK) is characterized for pixels in a representative rectangular region of no- or low-expression and the combination of signal plus background (SIG+BACK) is characterized for pixels in a representative rectangular region of high expression (e.g., Figures 3C, S7A, S8A, S10A, and S12A). For the pixels in these regions, we characterize the distribution by plotting an intensity histogram (e.g., Figures 3D, S7B, S8B, S10B, and S12B) and characterize average performance by calculating the mean pixel intensity (\bar{X}_n^{BACK} or $\bar{X}_n^{\text{SIG+BACK}}$ for replicate embryo n). Performance across replicate embryos is characterized by calculating the sample means (\bar{X}^{BACK} and $\bar{X}^{\text{SIG+BACK}}$) and standard errors ($s_{\bar{X}^{\text{BACK}}}$ and $s_{\bar{X}^{\text{SIG+BACK}}}$). The mean signal is then estimated as

$$\bar{X}^{\text{SIG}} = \bar{X}^{\text{SIG+BACK}} - \bar{X}^{\text{BACK}} \quad (\text{S4})$$

with the standard error estimated via uncertainty propagation as

$$s_{\bar{X}^{\text{SIG}}} \leq \sqrt{(s_{\bar{X}^{\text{SIG+BACK}}})^2 + (s_{\bar{X}^{\text{BACK}}})^2}. \quad (\text{S5})$$

The signal-to-background ratio is estimated as:

$$\bar{X}^{\text{SIG/BACK}} = \bar{X}^{\text{SIG}} / \bar{X}^{\text{BACK}} \quad (\text{S6})$$

with standard error estimated via uncertainty propagation as

$$s^{\text{SIG/BACK}} \leq \bar{X}^{\text{SIG/BACK}} \sqrt{\left(\frac{s_{\bar{X}^{\text{SIG}}}}{\bar{X}^{\text{SIG}}}\right)^2 + \left(\frac{s_{\bar{X}^{\text{BACK}}}}{\bar{X}^{\text{BACK}}}\right)^2}. \quad (\text{S7})$$

These upper bounds on estimated standard errors hold under the assumption that the correlation between SIG and BACK is non-negative. Tables S10–S16 display signal, background, and/or signal-to-background values to characterize the performance of in situ HCR v3.0 within whole-mount chicken embryos.

Experiment type	Quantity	Reagents			
		Probes	Hairpins	Expression region	
A	1	SIG+NSD+NSA+AF = SIG+BACK	odd + even	✓	high
	1	NSD+NSA+AF = BACK	odd + even	✓	no/low
B	2	NSA+AF		✓	high
	3	AF			high
C	4	SIG ^{odd} +NSD ^{odd} +NSA+AF = SIG ^{odd} +BACK ^{odd}	odd	✓	high
	4	NSD ^{odd} +NSA+AF = BACK ^{odd}	odd	✓	no/low
	5	SIG ^{even} +NSD ^{even} +NSA+AF = SIG ^{even} +BACK ^{even}	even	✓	high
	5	NSD ^{even} +NSA+AF = BACK ^{even}	even	✓	no/low

Table S5. Experiment types for qHCR imaging using in situ HCR v3.0. (A) Characterize signal, background, and signal-to-background. (B) Characterize components of background (AF, NSA, NSD). (C) Characterize split-initiator HCR suppression.

S1.4.3 Measurement of background components

Calculation of the signal-to-background ratio (Section S1.4.2) requires only a Type 1 experiment (using the terminology of Table S5A), yielding the values $\bar{X}^{\text{SIG+BACK}}$ and \bar{X}^{BACK} that are needed to calculate SIG/BACK. If desired, additional control experiments that omit certain reagents can be used to characterize the individual components of background (AF, NSA, NSD). A Type 2 experiment (no probes, hairpins only) yields $\bar{X}^{\text{NSA+AF}}$ and a Type 3 experiment (no probes, no hairpins) yields \bar{X}^{AF} .^{*} The background components can then be estimated via calculations analogous to (S4) and (S5). The estimated means are:

$$\bar{X}^{\text{NSD}} = \bar{X}^{\text{BACK}} - \bar{X}^{\text{NSA+AF}} \quad (\text{S8})$$

$$\bar{X}^{\text{NSA}} = \bar{X}^{\text{NSA+AF}} - \bar{X}^{\text{AF}} \quad (\text{S9})$$

with estimated standard errors:

$$s_{\bar{X}^{\text{NSD}}} \leq \sqrt{(s_{\bar{X}^{\text{BACK}}})^2 + (s_{\bar{X}^{\text{NSA+AF}}})^2} \quad (\text{S10})$$

$$s_{\bar{X}^{\text{NSA}}} \leq \sqrt{(s_{\bar{X}^{\text{NSA+AF}}})^2 + (s_{\bar{X}^{\text{AF}}})^2}. \quad (\text{S11})$$

These upper bounds on estimated standard errors hold under the assumption that the correlations are non-negative for the different components of background. For a given quantity, if the estimated mean is less than the estimated standard error, we report the standard error as an upper bound, and use this bound for uncertainty propagation.

Table S14B provides estimates for AF, NSA, and NSD when imaging *EphA4* in whole-mount chicken embryos using in situ HCR v3.0, all of which are small compared to SIG. Kinetically trapped HCR hairpins automatically suppress NSA and the combination of split-initiator probes and HCR hairpins automatically suppress NSD. Furthermore, HCR generates amplified SIG. All of these factors contribute to achieving a high signal-to-background ratio. If a Type 1 experiment demonstrates $\text{SIG} \gg \text{BACK}$, as is typically the case using in situ HCR v3.0, then there is little motivation to perform Type 2 and Type 3 experiments to characterize the individual background components (AF, NSA, NSD) as these are all bounded above by BACK.

S1.4.4 Measurement of split-initiator HCR suppression

To characterize performance of in situ HCR v3.0 in suppressing triggering of HCR by individual split-initiator probes, we augment experiments of Type 1 with additional control experiments that omit certain reagents. First, let us define the NSD and SIG observed using odd probes only or even probes only:

^{*}If a microscope generates non-negligible fluorescence intensities in the absence of sample, this so-called instrument noise (NOISE) should be taken into consideration when calculating background and signal contributions, leading to four Experiment Types (1. SIG+BACK+NOISE, 2. BACK+NOISE, 3. NSA+AF+NOISE, 4. AF+NOISE, 5. NOISE; cf. Table S5AB).

- odd non-specific detection (NSD^{odd}): odd probes that bind non-specifically in the sample and subsequently trigger HCR amplification.
- even non-specific detection (NSD^{even}): even probes that bind non-specifically in the sample and subsequently trigger HCR amplification.
- odd signal (SIG^{odd}): odd probes that hybridize specifically to the target mRNA and subsequently trigger HCR amplification.
- even signal (SIG^{even}): even probes that hybridize specifically to the target mRNA in the sample and subsequently trigger HCR amplification.

A Type 4 experiment (odd probes only, with hairpins) yields $\bar{X}^{\text{BACK}^{\text{odd}}+\text{SIG}^{\text{odd}}}$ and $\bar{X}^{\text{BACK}^{\text{odd}}}$ and a Type 5 experiment (even probes only, with hairpins) yields $\bar{X}^{\text{BACK}^{\text{even}}+\text{SIG}^{\text{even}}}$ and $\bar{X}^{\text{BACK}^{\text{even}}}$. These quantities in turn can be used to calculate SIG^{odd} and SIG^{even} via calculations analogous to (S4) and (S5). The estimated means are:

$$\bar{X}^{\text{SIG}^{\text{odd}}} = \bar{X}^{\text{SIG}^{\text{odd}}+\text{BACK}^{\text{odd}}} - \bar{X}^{\text{BACK}^{\text{odd}}} \quad (\text{S12})$$

$$\bar{X}^{\text{SIG}^{\text{even}}} = \bar{X}^{\text{SIG}^{\text{even}}+\text{BACK}^{\text{even}}} - \bar{X}^{\text{BACK}^{\text{even}}} \quad (\text{S13})$$

with estimated standard errors:

$$s_{\bar{X}^{\text{SIG}^{\text{odd}}}} \leq \sqrt{(s_{\bar{X}^{\text{SIG}^{\text{odd}}+\text{BACK}^{\text{odd}}}})^2 + (s_{\bar{X}^{\text{BACK}^{\text{odd}}}})^2} \quad (\text{S14})$$

$$s_{\bar{X}^{\text{SIG}^{\text{even}}}} \leq \sqrt{(s_{\bar{X}^{\text{SIG}^{\text{even}}+\text{BACK}^{\text{even}}}})^2 + (s_{\bar{X}^{\text{BACK}^{\text{even}}}})^2} \quad (\text{S15})$$

These upper bounds on estimated standard errors hold under the assumption that the correlations are non-negative for SIG^{odd} and BACK^{odd} and for SIG^{even} and BACK^{even}. For a given quantity, if the estimated mean is less than the estimated standard error, we report the standard error as an upper bound, and use this bound for uncertainty propagation.

Split-initiator HCR suppression can then be characterized by calculating SIG/SIG^{odd} and SIG/SIG^{even}, with higher values corresponding to more effective suppression. This in situ characterization is akin to the gel studies (Figures 2, S3, and S4) that compare test tube triggering of HCR by odd/even probe pairs colocalized by the target (lane 3) or by either probe alone (lanes 4 or 5). For the in situ data, signal-to-signal ratios are obtained via calculations analogous to (S6) and (S7). The estimated means are:

$$\bar{X}^{\text{SIG}/\text{SIG}^{\text{odd}}} = \bar{X}^{\text{SIG}} / \bar{X}^{\text{SIG}^{\text{odd}}} \quad (\text{S16})$$

$$\bar{X}^{\text{SIG}/\text{SIG}^{\text{even}}} = \bar{X}^{\text{SIG}} / \bar{X}^{\text{SIG}^{\text{even}}} \quad (\text{S17})$$

with estimated standard errors:

$$s_{\bar{X}^{\text{SIG}/\text{SIG}^{\text{odd}}}} \leq \bar{X}^{\text{SIG}/\text{SIG}^{\text{odd}}} \sqrt{\left(\frac{s_{\bar{X}^{\text{SIG}}}}{\bar{X}^{\text{SIG}}}\right)^2 + \left(\frac{s_{\bar{X}^{\text{SIG}^{\text{odd}}}}}{\bar{X}^{\text{SIG}^{\text{odd}}}}\right)^2} \quad (\text{S18})$$

$$s_{\bar{X}^{\text{SIG}/\text{SIG}^{\text{even}}}} \leq \bar{X}^{\text{SIG}/\text{SIG}^{\text{even}}} \sqrt{\left(\frac{s_{\bar{X}^{\text{SIG}}}}{\bar{X}^{\text{SIG}}}\right)^2 + \left(\frac{s_{\bar{X}^{\text{SIG}^{\text{even}}}}}{\bar{X}^{\text{SIG}^{\text{even}}}}\right)^2}. \quad (\text{S19})$$

These upper bounds on estimated standard errors hold under the assumption that the correlations are non-negative for SIG and SIG^{odd} and for SIG and SIG^{even}.

Table S14C displays the signal-to-signal ratios SIG/SIG^{odd} and SIG/SIG^{even} when imaging *EphA4* in whole-mount chicken embryos using in situ HCR v3.0. Split-initiator probes colocalized by the target are more than an order of magnitude more effective at triggering HCR than odd or even probes alone. Interestingly, with this assay, we are quantifying the automatic *background* suppression capabilities of the split-initiator probes by measuring automatic *signal* suppression, taking advantage of the fact that the target molecules in the embryo will colocalize odd/even probe pairs for a Type 1 experiment and will localize odd or even probes to the same expression region for a Type 4 or 5 experiment.

S1.4.5 Normalized voxel intensities for qHCR imaging: analog mRNA relative quantitation with subcellular resolution in an anatomical context

For quantitative mRNA imaging using in situ HCR, precision increases with voxel size as long as the imaging voxels remain smaller than the features in the expression pattern (see Section S2.2 of Trivedi *et al.* (2018)). To increase precision, we calculate raw voxel intensities by averaging neighboring pixel intensities while still maintaining a subcellular voxel size. To facilitate relative quantitation between voxels, we estimate the normalized HCR signal of voxel j in replicate n as:

$$x_{n,j} \equiv \frac{X_{n,j}^{\text{SIG+BACK}} - X^{\text{BOT}}}{X^{\text{TOP}} - X^{\text{BOT}}}, \quad (\text{S20})$$

which translates and rescales the data so that the voxel intensities in each channel fall in the interval $[0,1]$. Here,

$$X^{\text{BOT}} \equiv \bar{X}^{\text{BACK}} \quad (\text{S21})$$

is the mean background across replicates (see Section S1.4.2) and

$$X^{\text{TOP}} \equiv \max_{n,j} X_{n,j}^{\text{SIG+BACK}} \quad (\text{S22})$$

is the maximum total fluorescence for a voxel across replicates.

Pairwise expression scatter plots that each display normalized voxel intensities for two channels (e.g., Figures 4 and 5 of Trivedi *et al.* (2018)) provide a powerful quantitative framework for performing multidimensional read-out/read-in analyses (Figure 6 of Trivedi *et al.* (2018)). Read-out from anatomical space to expression space enables discovery of expression clusters of voxels with quantitatively related expression levels and ratios (amplitudes and slopes in the expression scatter plots), while read-in from expression space to anatomical space enables discovery of the corresponding anatomical locations of these expression clusters within the embryo. The simple and practical normalization approach of (S20)–(S22) translates and rescales all voxels identically within a given channel (enabling comparison of amplitudes and slopes in scatter plots between replicates), and does not attempt to remove scatter in the normalized signal estimate that is caused by scatter in the background.

To validate relative mRNA quantitation with subcellular resolution ($2 \times 2 \times 2.7 \mu\text{m}$ voxels) in whole-mount chicken embryos, Figures 5C, S18C, and S19C display highly correlated normalized voxel intensities for 2-channel redundant detection of *Dmbx1* and *EphA4*. In this setting, accuracy corresponds to linearity with zero intercept, and precision corresponds to scatter around the line (Trivedi *et al.*, 2018).

S1.4.6 Dot detection and colocalization for dHCR imaging: digital mRNA absolute quantitation in an anatomical context

To validate the performance of in situ HCR (v3.0) for single-molecule imaging, we perform a 2-channel redundant detection experiment in which a target mRNA is detected using two independent probe sets and HCR amplifiers. Let N_1 denote the number of dots detected in channel 1, N_2 the number of dots detected in channel 2, and N_{12} the number of colocalized dots appearing in both channels. We define the colocalization fraction for each channel:

$$C_1 = N_{12}/N_1, \quad (\text{S23})$$

$$C_2 = N_{12}/N_2. \quad (\text{S24})$$

As the false-positive and false-negative rates for single-molecule detection go to zero, C_1 and C_2 will both approach 1 from below, providing a quantitative basis for evaluating performance. Colocalization results using in situ HCR v3.0 with split-initiator probes (23–25 probe pairs per channel) in cultured human cells (Table S25) and whole-mount chicken embryos (Table S26) are $\approx 84\%$, compared with $\approx 50\%$ using in situ HCR v2.0 (39 standard probe pairs per channel) in a previous study in whole-mount zebrafish embryos (Table S27)(Shah *et al.*, 2016).

Single molecules were identified in each channel using the following dot detection algorithm applied to a three-dimensional confocal image stack:

- **Step 1: Blur noise.** To remove noise smaller than the dots of interest, the image was convolved with an isotropic Gaussian blur (standard deviation σ_{blur}).
- **Step 2: Local background subtraction.** To eliminate variations in pixel intensity arising from background variations that occur on a length scale larger than the dots of interest, local background subtraction was performed by subtracting the mean pixel intensity of a cube (edge length d_{back}) from the intensity of the pixel at the center of the cube.
- **Step 3: Global threshold on pixel intensity.** To eliminate dim features, the resulting pixel intensities were subjected to a global threshold (t_{pixel}), and the range $[t_{\text{pixel}}, 1]$ was renormalized to a $[0, 1]$ scale.
- **Step 4: Watershed dot detection.** To identify single mRNA molecules as dots within the image, regional image maxima were segmented using the minimum saliency watershed method (Coupric & Bertrand, 1997; Yoo *et al.*, 2002). In this method, two maxima are labeled as the same dot if the minimum boundary height between the maxima is less than a given threshold ($t_{\text{watershed}}$). Dot coordinates were estimated as the intensity-weighted centroid of each watershed basin, and dot intensities were estimated as the integrated pixel intensity within each basin.
- **Step 5: Global threshold on dot intensity.** To eliminate dim dots, the resulting dot intensities were normalized on a $[0, 1]$ scale and a global threshold (t_{dot}) was applied. The resulting number of dots N_i was recorded for channel i .

After identifying the dots in each channel of a 2-channel redundant detection image, a dot i in Channel 1 and a dot j in Channel 2 were considered colocalized if all of the following four statements were true:

- **Test 1:** The xy centroids differed by less than a lateral distance threshold ($r_{xy} = 0.22 \mu\text{m}$).
- **Test 2:** Dot i is the closest dot in Channel 1 to dot j in Channel 2.
- **Test 3:** Dot j is the closest dot in Channel 2 to dot i in Channel 1.
- **Test 4:** The z centroids differed by less than the axial distance threshold ($r_z = 0.42 \mu\text{m}$).

Note that due to the lower axial resolution, dot colocalization was tested separately for xy and z . The same distance thresholds were used across all sample types and replicates.[†]

[†]For chicken embryo replicate 3, Channels 1 and 2 were manually aligned by applying a constant offset of $+0.35 \mu\text{m}$ to the z -coordinates of Channel 1 after the detected dots showed a clear bias in z coordinates between the two channels. No other images were manually aligned.

		HEK cells		Chicken embryos		Zebrafish embryos	
	lateral resolution d_{xy} (μm)	0.0624		0.099		0.2167	
	axial resolution d_z (μm)	0.42		0.42		0.3369	
	Step 1: Gaussian blur radius σ_{blur} (μm)	0.2		0.2		0.1	
	Step 2: Mean subtraction cube length d_{back} (μm)	2.5		3.0		5.5	
	Replicate	Ch1	Ch2	Ch1	Ch2	Ch1	Ch2
Step 3: Global pixel threshold (t_{pixel})	1	0.003	0.012	0.04	0.02	0.02	0.02
	2	0.005	0.012	0.04	0.02	0.01	0.003
	3	0.004	0.01	0.04	0.02	0.01	0.005
Step 4: Watershed saliency minimum ($t_{\text{watershed}}$)	1	0.15	0.15	0.15	0.15	0.15	0.15
	2	0.15	0.15	0.15	0.15	0.30	0.20
	3	0.15	0.15	0.15	0.15	0.35	0.30
Step 5: Global dot intensity threshold (t_{dot})	1	0.01	0.005	0.00	0.01	0.01	0.01
	2	0.02	0.02	0.006	0.013	0.025	0.025
	3	0.03	0.012	0.01	0.025	0.04	0.04

Table S6. Parameters used for dot detection in dHCR images.

S1.5 Flow cytometry data analysis

Data analysis for flow cytometry experiments on cultured cells closely follows the image analysis of Section S1.4 as detailed below.

S1.5.1 Raw cell intensities

The components of background (AF, NSA, NSD) and signal (SIG) are defined as before (Section S1.4.1), where n is treated as an index over cells, and $i = 1$ for each cell since the flow cytometer returns one value per cell.

S1.5.2 Measurement of signal, background, signal-to-background, background components, and split-initiator HCR suppression for transgenic targets

For a transgenic target mRNA, signal and background are characterized based on flow cytometry experiments of Types 1a and 1b (Table S7A) with SIG+BACK measured in transgenic cells containing the target and BACK measured in wildtype (WT) cells lacking the target. This approach parallels that for characterizing signal and background in images (Section S1.4.2) with transgenic cells taking the place of a region of high expression and WT cells taking the place of a region of no/low expression. If desired, additional control experiments of Types 2 and 3 (Table S7B) can be performed to characterize the components of background (AF, NSA, NSD) using the calculations of Section S1.4.3. Likewise, additional control experiments of Types 4a, 4b, 5a, and 5b (Table S7C) can be performed to characterize split-initiator HCR suppression using the calculations of Section S1.4.4. For transgenic target mRNAs in human and bacterial cells, Figures S20 and S22 display distributions of cell intensities characterizing signal and background (panel A) and split-initiator HCR suppression (panel B). Tables S17 and S19 display corresponding values for SIG, BACK, and SIG/BACK (panel A), background components AF, NSA, and NSD (panel B), and SIG/SIG^{odd} and SIG/SIG^{even} (panel C).

Experiment type	Quantity	Reagents			
		Probes	Hairpins	Cell type	
A	1a	$SIG+NSD+NSA+AF = SIG+BACK$	odd + even	✓	transgenic
	1b	$NSD+NSA+AF = BACK$	odd + even	✓	WT
B	2	NSA+AF		✓	transgenic
	3	AF			transgenic
C	4a	$SIG^{odd}+NSD^{odd}+NSA+AF = SIG^{odd}+BACK^{odd}$	odd	✓	transgenic
	4b	$NSD^{odd}+NSA+AF = BACK^{odd}$	odd	✓	WT
	5a	$SIG^{even}+NSD^{even}+NSA+AF = SIG^{even}+BACK^{even}$	even	✓	transgenic
	5b	$NSD^{even}+NSA+AF = BACK^{even}$	even	✓	WT

Table S7. Experiment types for flow cytometry using in situ HCR v3.0 with a transgenic target mRNA. (A) Characterize signal, background, and signal-to-background. (B) Characterize components of background (AF, NSA, NSD). (C) Characterize split-initiator HCR suppression.

S1.5.3 Measurement of signal, background, signal-to-background, background components, and split-initiator HCR suppression for endogenous targets

For an endogenous target mRNA, signal and background are characterized based on flow cytometry experiments of Types 1a and 1b (Table S8A) with SIG+BACK measured using a probe set (odd + even) that address the target in WT cells and BACK measured using a probe set (Tg(odd) and Tg(even)) that addresses a different transgenic target absent from WT cells. Use of a previously validated transgenic probe set to measure background in WT cells ensures that a low measured fluorescence value does not simply indicate a dysfunctional probe set, but indeed represents low background generated by a probe set that is known to be functional if the target is present in the sample. If desired,

Experiment type	Quantity	Reagents		Cell type	
		Probes	Hairpins		
A	1a	$SIG+NSD+NSA+AF = SIG+BACK$	odd + even	✓	WT
	1b	$NSD+NSA+AF = BACK$	Tg(odd) + Tg(even)	✓	WT
B	2	NSA+AF		✓	WT
	3	AF			WT
C	4a	$SIG^{odd}+NSD^{odd}+NSA+AF = SIG^{odd}+BACK^{odd}$	odd	✓	WT
	4b	$NSD^{odd}+NSA+AF = BACK^{odd}$	Tg(odd)	✓	WT
	5a	$SIG^{even}+NSD^{even}+NSA+AF = SIG^{even}+BACK^{even}$	even	✓	WT
	5b	$NSD^{even}+NSA+AF = BACK^{even}$	Tg(even)	✓	WT

Table S8. Experiment types for flow cytometry using in situ HCR v3.0 with an endogenous target mRNA. (A) Characterize signal, background, and signal-to-background. (B) Characterize components of background (AF, NSA, NSD). (C) Characterize split-initiator HCR suppression. Here, Tg(odd) and Tg(even) denote odd and even probes from a probe set targeting a transgenic mRNA that is absent from WT cells.

additional control experiments of Types 2 and 3 (Table S8B) can be performed to characterize the components of background (AF, NSA, NSD) using the calculations of Section S1.4.3. Likewise, additional control experiments of Types 4a, 4b, 5a, and 5b (Table S8C) can be performed to characterize split-initiator HCR suppression using the calculations of Section S1.4.4. For endogenous target mRNA *GAPDH*, Figure S21 displays distributions of cell intensities characterizing signal and background (panel A) and split-initiator HCR suppression (panel B). Table S18 displays corresponding values for SIG, BACK, and SIG/BACK (panel A), background components AF, NSA, and NSD (panel B), and SIG/SIG^{odd} and SIG/SIG^{even} (panel C).

S1.5.4 Normalized single-cell intensities for qHCR flow cytometry: analog mRNA relative quantitation for high-throughput expression profiling of human and bacterial cells

For relative mRNA quantitation between cells, the single-cell intensities within a channel are normalized using equation (S20) with BOT (mean BACK intensity across cells) and TOP (maximum SIG+BACK intensity for a single cell) defined by equations (S21) and (S22). Redundant detection experiments validating mRNA single-cell relative quantitation are displayed for endogenous targets *GAPDH* (Figure S24 and Table S20) and *PGK1* (Figure S25 and Table S21) in HEK cells, and for endogenous target *fusA* in *E. coli* (Figure S26 and Table S22).

S2 Protocols for in situ HCR v3.0

S2.1 Protocols for whole-mount chicken embryos

S2.1.1 Preparation of fixed whole-mount chicken embryos

1. Collect chicken embryos on 3M filter paper and place in a petri dish containing Ringer's solution.
2. Transfer embryos into a new petri dish with fresh Ringer's solution.
NOTE: This is to rinse away egg yolk before fixation.
3. Transfer into a petri dish containing 4% paraformaldehyde (PFA).
CAUTION: Use PFA with extreme care as it is a hazardous material.
NOTE: Use fresh PFA to avoid increased autofluorescence.
4. Fix the samples at room temperature for 1 h.
5. Transfer embryos into a petri dish containing PBST.
6. Dissect the embryos off the filter paper. Remove the fixed vitelline membrane (opaque) and cut the squares around area pellucida, without leaving excess of extra-embryonic tissue.
7. Transfer embryos into a 2 mL Eppendorf tube containing PBST.
8. Nutate for 5 mins with the tube positioned horizontally in a small ice bucket.
9. Wash embryos two additional times with 2 mL of PBST, each time nutating for 5 min on ice.
10. Dehydrate embryos with 2 × 5 min washes of 2 mL methanol (MeOH) on ice.
11. Store embryos at -20 °C overnight before use.
NOTE: Embryos can be stored for six months at -20 °C.
12. Transfer the required number of embryos for an experiment to a 2 mL Eppendorf tube.
NOTE: Do not place more than 4 embryos in each 2 mL Eppendorf tube.
13. Rehydrate with a series of graded 2 mL MeOH/PBST washes for 5 min each on ice:
 - (a) 75% MeOH / 25% PBST
 - (b) 50% MeOH / 50% PBST
 - (c) 25% MeOH / 75% PBST
 - (d) 100% PBST
 - (e) 100% PBST.
14. Treat embryos with 2 mL of 10 µg/mL proteinase K solution for 2 min (stage HH 8) or 2.5 min (stages HH 10–11) at room temperature.
NOTE: Proteinase K concentration and treatment time should be reoptimized for each batch of proteinase K, or for samples at a different developmental stage.
15. Postfix with 2 mL of 4% PFA for 20 min at room temperature.
16. Wash embryos 2 × 5 min with 2 mL of PBST on ice.
17. Wash embryos with 2 mL of 50% PBST / 50% 5× SSCT for 5 min on ice.
18. Wash embryos with 2 mL of 5× SSCT for 5 min on ice.

S2.1.2 Buffer recipes for sample preparation

Ringer's solution

123 mM NaCl
1.53 mM CaCl₂
4.96 mM KCl₂
0.81 mM Na₂HPO₄
0.15 mM KH₂PO₄

For 2 L of solution

14.4 g of NaCl
340 mg of CaCl₂
740 mg of KCl
230 mg of Na₂HPO₄
40 mg of KH₂PO₄
Bring volume up to 1.5 L with ultrapure H₂O
Adjust pH to 7.4 and fill up to 2 L with ultrapure H₂O
Filter sterilize with 0.22 μm bottle top filter

4% Paraformaldehyde (PFA)

4% PFA
1× PBS

For 25 mL of solution

1 g of PFA powder
25 mL of 1× PBS
Heat solution at 50–60 °C to dissolve powder

PBST

1× PBS
0.1% Tween 20

For 50 mL of solution

5 mL of 10× PBS
500 μL of 10% Tween 20
Fill up to 50 mL with ultrapure H₂O

Proteinase K solution

10 μg/mL proteinase K

For 2 mL of solution

1 μL of 20 mg/mL proteinase K
Fill up to 2 mL with PBST

NOTE: Avoid using calcium chloride and magnesium chloride in PBS as this leads to increased autofluorescence in the embryos.

S2.1.3 Multiplexed in situ HCR v3.0 using split-initiator probes

Detection stage

1. For each sample, transfer 1-4 embryos to a 2 mL Eppendorf tube.
NOTE: Do not place more than 4 embryos in each 2 mL Eppendorf tube.
2. Incubate embryos in 1 mL of 30% probe hybridization buffer on ice for 5 min.
CAUTION: probe hybridization buffer contains formamide, a hazardous material.
NOTE: Pre-heat probe hybridization buffer to 37 °C before use.
3. Remove the buffer and pre-hybridize with 1 mL of 30% probe hybridization buffer for 30 min at 37 °C.
4. Prepare probe solution by adding 4 pmol of each probe mixture (odd & even: 2 μ L of 2 μ M stock per probe mixture) to 1 mL of 30% probe hybridization buffer at 37 °C.
NOTE: For Figures 7B and S30 (dHCR), 10 pmol of each probe was used to improve probe hybridization efficiency.
5. Remove the pre-hybridization solution and add the probe solution.
6. Incubate embryos overnight (12–16 h) at 37 °C.
7. Remove excess probes by washing embryos 4 \times 15 min with 1 mL of 30% probe wash buffer at 37 °C:
CAUTION: probe wash buffer contains formamide, a hazardous material.
NOTE: Pre-heat probe wash buffer to 37 °C before use.
8. Wash samples 2 \times 5 min with 5 \times SSCT at room temperature.

Amplification stage

1. Pre-amplify embryos with 500 μ L of amplification buffer for 5 min at room temperature.
NOTE: Equilibrate amplification buffer to room temperature before use.
2. Prepare 30 pmol of each fluorescently labeled hairpin by snap cooling 10 μ L of 3 μ M stock in hairpin storage buffer (heat at 95 °C for 90 seconds and cool to room temperature in a dark drawer for 30 min).
3. Prepare hairpin solution by adding all snap-cooled hairpins to 500 μ L of amplification buffer at room temperature.
4. Remove the pre-amplification solution and add the hairpin solution.
5. Incubate the embryos overnight (12–16 h) in the dark at room temperature.
NOTE: For Figures 7B and S30 (dHCR), a 90 min amplification time was used to ensure single-molecule dots are diffraction-limited.
6. Remove excess hairpins by washing with 1 mL of 5 \times SSCT at room temperature:
 - (a) 2 \times 5 min
 - (b) 2 \times 30 min
 - (c) 1 \times 5 min

S2.1.4 Buffer recipes for in situ HCR v3.0

Probes, amplifiers, probe hybridization buffer, and probe wash buffer should be stored at -20 °C. Amplification buffer should be stored at 4 °C. Keep these reagents on ice at all times during probe and amplifier preparation. Make sure all solutions are well mixed before use.

30% probe hybridization buffer

30% formamide
 5× sodium chloride sodium citrate (SSC)
 9 mM citric acid (pH 6.0)
 0.1% Tween 20
 50 µg/mL heparin
 1× Denhardt's solution
 10% dextran sulfate

For 40 mL of solution

12 mL formamide
 10 mL of 20× SSC
 360 µL 1 M citric acid, pH 6.0
 400 µL of 10% Tween 20
 200 µL of 10 mg/mL heparin
 800 µL of 50× Denhardt's solution
 8 mL of 50% dextran sulfate
 Fill up to 40 mL with ultrapure H₂O

30% probe wash buffer

30% formamide
 5× sodium chloride sodium citrate (SSC)
 9 mM citric acid (pH 6.0)
 0.1% Tween 20
 50 µg/mL heparin

For 40 mL of solution

12 mL formamide
 10 mL of 20× SSC
 360 µL 1 M citric acid, pH 6.0
 400 µL of 10% Tween 20
 200 µL of 10 mg/mL heparin
 Fill up to 40 mL with ultrapure H₂O

Amplification buffer

5× sodium chloride sodium citrate (SSC)
 0.1% Tween 20
 10% dextran sulfate

For 40 mL of solution

10 mL of 20× SSC
 400 µL of 10% Tween 20
 8 mL of 50% dextran sulfate
 Fill up to 40 mL with ultrapure H₂O

5× SSCT

5× sodium chloride sodium citrate (SSC)
 0.1% Tween 20

For 40 mL of solution

10 mL of 20× SSC
 400 µL of 10% Tween 20
 Fill up to 40 mL with ultrapure H₂O

50% dextran sulfate

50% dextran sulfate

For 40 mL of solution

20 g of dextran sulfate powder
 Fill up to 40 mL with ultrapure H₂O

S2.1.5 Sample mounting for microscopy

1. Make a chamber for mounting each embryo by aligning two stacks of double-sided tape (2 pieces per stack) 1 cm apart on a 25 mm × 75 mm glass slide.
2. Place an embryo between the tape stacks on the slide and remove as much solution as possible.
3. Align the embryo for dorsal imaging and carefully touch the slide with a kimwipe to further dry the area around the embryo.
4. Add two drops of SlowFade Gold antifade mountant on top of the embryo.
5. Place a 22 mm × 30 mm No. 1 coverslip on top of the stacks to close the chamber.

NOTE: See Section S1.2 for details of confocal microscopes used to image whole-mount chicken embryos.

S2.1.6 Reagents and supplies

Paraformaldehyde (PFA) (Sigma Cat. # P6148)
Methanol (Mallinckrodt Chemicals Cat. # 3016-16)
Proteinase K, molecular biology grade (NEB Cat. # P8107S)
Formamide (Deionized) (Ambion Cat. # AM9342)
20× sodium chloride sodium citrate (SSC) (Life Technologies Cat. # 15557-044)
Heparin (Sigma Cat. # H3393)
50% Tween 20 (Life Technologies Cat. # 00-3005)
50× Denhardt's solution (Life Technologies Cat. # 750018)
Dextran sulfate, mol. wt. > 500,000 (Sigma Cat. # D6001)
25 mm × 75 mm glass slide (VWR Cat. # 48300-025)
22 mm × 30 mm No. 1 coverslip (VWR Cat. # 48393-026)
SlowFade Gold antifade mountant (Life Technologies Cat. # S36937)

S2.2 Protocols for mammalian cells on a chambered slide

S2.2.1 Preparation of fixed mammalian cells on a chambered slide

1. Coat bottom of each chamber by applying 300 μL of 0.01% poly-D-lysine prepared in cell culture grade H_2O .
NOTE: A volume of 300 μL is sufficient per chamber on an 8-chamber slide.
2. Incubate for at least 30 min at room temperature.
3. Aspirate the coating solution and wash each chamber twice with molecular biology grade H_2O .
4. Plate desired number of cells in each chamber.
5. Grow cells to desired confluency for 24–48 h.
6. Aspirate growth media and wash each chamber with 300 μL of DPBS.
NOTE: Avoid using calcium chloride and magnesium chloride in DPBS as this leads to increased autofluorescence.
7. Add 300 μL of 4% formaldehyde to each chamber.
CAUTION: Use formaldehyde with extreme care as it is a hazardous material.
8. Incubate for 10 min at room temperature.
9. Remove fixative and wash each chamber $2 \times 300 \mu\text{L}$ of DPBS.
10. Aspirate DPBS and add 300 μL of ice-cold 70% ethanol.
11. Permeabilize cells overnight at -20°C .
12. Cells can be stored at -20°C or 4°C until use.

S2.2.2 Buffer recipes for sample preparation

4% formaldehyde in PBS

4% formaldehyde

1× PBS

For 10 mL of solution

2.5 mL of 16% formaldehyde

1 mL of 10× PBS

Fill up to 10 mL with molecular biology grade H₂O

NOTE: Avoid using calcium chloride and magnesium chloride in PBS as this leads to increased autofluorescence in the samples.

S2.2.3 Multiplexed in situ HCR v3.0 using split-initiator probes

Detection stage

1. Aspirate EtOH and air dry samples at room temperature.
NOTE: Drying of sample is optional.
2. Wash samples two times with 300 μL of $2\times$ SSC.
3. Pre-hybridize samples in 300 μL of 30% probe hybridization buffer for 30 min at 37 $^{\circ}\text{C}$.
CAUTION: probe hybridization buffer contains formamide, a hazardous material.
NOTE: Pre-heat probe hybridization buffer to 37 $^{\circ}\text{C}$ before use.
4. Prepare probe solution by adding 1.2 pmol of each probe mixture (odd & even: 0.6 μL of 2 μM stock per probe mixture) to 300 μL of 30% probe hybridization buffer at 37 $^{\circ}\text{C}$.
NOTE: For Figures 7A and S29 (dHCR), 3 pmol of each probe was used to improve probe hybridization efficiency.
5. Remove the pre-hybridization solution and add the probe solution.
6. Incubate samples overnight (12–16 h) at 37 $^{\circ}\text{C}$.
7. Remove excess probes by washing 4×5 min with 300 μL of 30% probe wash buffer at 37 $^{\circ}\text{C}$.
CAUTION: probe wash buffer contains formamide, a hazardous material.
NOTE: Pre-heat probe wash buffer to 37 $^{\circ}\text{C}$ before use.
8. Wash samples 2×5 min with $5\times$ SSCT at room temperature.

Amplification stage

1. Pre-amplify samples in 300 μL of amplification buffer for 30 min at room temperature.
NOTE: Equilibrate amplification buffer to room temperature before use.
2. Prepare 18 pmol of each fluorescently labeled hairpin by snap cooling 6 μL of 3 μM stock in hairpin storage buffer (heat at 95 $^{\circ}\text{C}$ for 90 seconds and cool to room temperature in a dark drawer for 30 min).
3. Prepare hairpin solution by adding all snap-cooled hairpins to 300 μL of amplification buffer at room temperature.
4. Remove the pre-amplification solution and add the hairpin solution.
5. Incubate samples overnight (12–16 h) in the dark at room temperature.
NOTE: For Figures 7A and S29 (dHCR), a 45 min amplification time was used to ensure single-molecule dots are diffraction-limited.
6. Remove excess hairpins by washing 5×5 min with 300 μL of $5\times$ SSCT at room temperature.
7. Aspirate $5\times$ SSCT and add ≈ 100 μL of SlowFade Gold antifade mountant with DAPI.
8. Samples can be stored at 4 $^{\circ}\text{C}$ protected from light prior to imaging.

S2.2.4 Buffer recipes for in situ HCR v3.0

Probes, amplifiers, probe hybridization buffer, and probe wash buffer should be stored at -20 °C. Amplification buffer should be stored at 4 °C. Keep these reagents on ice at all times during probe and amplifier preparation. Make sure all solutions are well mixed before use.

30% probe hybridization buffer

30% formamide
 5× sodium chloride sodium citrate (SSC)
 9 mM citric acid (pH 6.0)
 0.1% Tween 20
 50 µg/mL heparin
 1× Denhardt's solution
 10% dextran sulfate

For 40 mL of solution

12 mL formamide
 10 mL of 20× SSC
 360 µL 1 M citric acid, pH 6.0
 400 µL of 10% Tween 20
 200 µL of 10 mg/mL heparin
 800 µL of 50× Denhardt's solution
 8 mL of 50% dextran sulfate
 Fill up to 40 mL with ultrapure H₂O

30% probe wash buffer

30% formamide
 5× sodium chloride sodium citrate (SSC)
 9 mM citric acid (pH 6.0)
 0.1% Tween 20
 50 µg/mL heparin

For 40 mL of solution

12 mL formamide
 10 mL of 20× SSC
 360 µL 1 M citric acid, pH 6.0
 400 µL of 10% Tween 20
 200 µL of 10 mg/mL heparin
 Fill up to 40 mL with ultrapure H₂O

Amplification buffer

5× sodium chloride sodium citrate (SSC)
 0.1% Tween 20
 10% dextran sulfate

For 40 mL of solution

10 mL of 20× SSC
 400 µL of 10% Tween 20
 8 mL of 50% dextran sulfate
 Fill up to 40 mL with ultrapure H₂O

5× SSCT

5× sodium chloride sodium citrate (SSC)
 0.1% Tween 20

For 40 mL of solution

10 mL of 20× SSC
 400 µL of 10% Tween 20
 Fill up to 40 mL with ultrapure H₂O

50% dextran sulfate

50% dextran sulfate

For 40 mL of solution

20 g of dextran sulfate powder
 Fill up to 40 mL with ultrapure H₂O

S2.2.5 Reagents and supplies

Molecular biology grade H₂O (Corning Cat. # 46-000-CV)
16% Formaldehyde (w/v), Methanol-free (Life Technologies Cat. # 28906)
DPBS, no calcium, no magnesium (Life Technologies Cat. # 14190144)
10× PBS (Ambion Cat. # AM9624)
Formamide (Deionized) (Ambion Cat. # AM9342)
20× sodium chloride sodium citrate (SSC) (Life Technologies Cat. # 15557-044)
Heparin (Sigma Cat. # H3393)
10% Tween 20 (BioRad Cat. # 161-0781)
50× Denhardt's solution (Life Technologies Cat. # 750018)
Dextran sulfate, mol. wt. > 500,000 (Sigma Cat. # D6001)
ibidi μ -slide ibitreat (ibidi Cat. # 80826)
SlowFade Gold antifade mountant with DAPI (Life Technologies Cat. # S36938)

S2.3 Protocols for mammalian cells in suspension

S2.3.1 Preparation of fixed mammalian cells in suspension

1. Aspirate growth media from culture plate and wash cells with DPBS.

NOTE: Avoid using calcium chloride and magnesium chloride in DPBS as this leads to increased autofluorescence.

2. Add 3 mL of trypsin per 10 cm plate and incubate in a 5% CO₂ incubator for 5 min at 37 °C.

3. Quench trypsin by adding 3 mL of growth media.

4. Transfer cells to a conical tube and centrifuge for 5 min at 180 × g.

5. Aspirate supernatant and re-suspend cells in 4% formaldehyde to reach ≈ 10⁶ cells/mL.

CAUTION: Use formaldehyde with extreme care as it is a hazardous material.

6. Fix cells for 1 hr at room temperature.

7. Centrifuge for 5 min at 180 × g and remove supernatant.

8. Wash cells 4 times with PBST (use the same volume as formaldehyde solution).

NOTE: Centrifuge for 5 min at 180 × g and aspirate supernatant between washes.

9. Re-suspend cells in ice-cold 70% ethanol (use the same volume as formaldehyde and PBST solutions).

10. Permeabilize cells overnight at 4 °C.

11. Cells can be stored at 4 °C until use.

S2.3.2 Buffer recipes for sample preparation

4% formaldehyde in PBST

4% formaldehyde

1× PBS, 0.1% Tween 20

For 36 mL of solution

9 mL of 16% formaldehyde

3.6 mL of 10× PBST

180 μ L of 10% Tween 20

Fill up to 36 mL with ultrapure H₂O

NOTE: Avoid using calcium chloride and magnesium chloride in PBS as this leads to increased autofluorescence in the samples.

S2.3.3 Multiplexed in situ HCR v3.0 using split-initiator probes

Detection stage

1. Transfer desired amount ($0.5-1 \times 10^6$) of fixed cells into a 1.5 mL Eppendorf tube.
2. Centrifuge for 5 min to remove EtOH.
NOTE: All centrifugation steps are done at $180 \times g$.
3. Wash cells twice with 500 μL of PBST. Centrifuge for 5 min to remove supernatant.
4. Re-suspend the pellet with 400 μL of 30% LMW probe hybridization buffer and pre-hybridize for 30 min at 37 °C.
CAUTION: probe hybridization buffer contains formamide, a hazardous material.
NOTE: Pre-heat probe hybridization buffer to 37 °C before use.
5. In the meantime, prepare probe solution by adding 2 pmol of each probe mixture (odd & even: 1 μL of 2 μM stock per probe mixture) to 100 μL of 30% LMW probe hybridization buffer at 37 °C.
6. Add the probe solution directly to the sample to reach a final probe concentration of 4 nM.
7. Incubate the sample overnight at 37 °C.
8. Centrifuge for 5 min to remove probe solution.
9. Re-suspend the cell pellet with 500 μL of 30% probe wash buffer.
CAUTION: probe wash buffer contains formamide, a hazardous material.
NOTE: Pre-heat probe wash buffer to 37 °C before use.
10. Incubate for 10 min at 37 °C and remove the wash solution by centrifugation for 5 min.
11. Repeat steps 9 and 10 for three additional times.
12. Re-suspend the cell pellet with 500 μL of 5 \times SSCT.
13. Incubate for 5 min at room temperature.
14. Proceed to hairpin amplification.

Amplification stage

1. Centrifuge for 5 min to pellet the cells.
2. Re-suspend the cell pellet with 150 μL of LMW amplification buffer and pre-amplify for 30 min at room temperature.
NOTE: Equilibrate amplification buffer to room temperature before use.
3. Prepare 15 pmol of each fluorescently labeled hairpin by snap cooling 5 μL of 3 μM stock in hairpin storage buffer (heat at 95 °C for 90 seconds and cool to room temperature in a dark drawer for 30 min).
4. Prepare hairpin mixture by adding all snap-cooled hairpins to 100 μL of LMW amplification buffer at room temperature.
5. Add the hairpin mixture directly to the sample to reach a final hairpin concentration of 60 nM.
6. Incubate the sample overnight (>12 h) in the dark at room temperature.
7. Centrifuge for 5 min and remove the hairpin solution.

8. Re-suspend the cell pellet with 500 μ L of 5 \times SSCT.
9. Without incubation, remove the wash solution by centrifugation for 5 min.
10. Repeat steps 8 and 9 for five additional times.
11. Re-suspend the cell pellet in desired buffer and volume.
NOTE: Samples can be stored at 4 °C protected from light before flow cytometry or imaging.
12. Filter cells before flow cytometry.

S2.3.4 Buffer recipes for in situ HCR v3.0

Probes, amplifiers, probe hybridization buffer, and probe wash buffer should be stored at -20 °C. Amplification buffer should be stored at 4 °C. Keep these reagents on ice at all times during probe and amplifier preparation. Make sure all solutions are well mixed before use.

30% probe hybridization buffer (LMW d.s.)

30% formamide
 5× sodium chloride sodium citrate (SSC)
 9 mM citric acid (pH 6.0)
 0.1% Tween 20
 50 µg/mL heparin
 1× Denhardt's solution
 10% low MW dextran sulfate

For 40 mL of solution

12 mL formamide
 10 mL of 20× SSC
 360 µL 1 M citric acid, pH 6.0
 400 µL of 10% Tween 20
 200 µL of 10 mg/mL heparin
 800 µL of 50× Denhardt's solution
 8 mL of 50% low MW dextran sulfate
 Fill up to 40 mL with ultrapure H₂O

30% probe wash buffer

30% formamide
 5× sodium chloride sodium citrate (SSC)
 9 mM citric acid (pH 6.0)
 0.1% Tween 20
 50 µg/mL heparin

For 40 mL of solution

12 mL formamide
 10 mL of 20× SSC
 360 µL 1 M citric acid, pH 6.0
 400 µL of 10% Tween 20
 200 µL of 10 mg/mL heparin
 Fill up to 40 mL with ultrapure H₂O

Amplification buffer (LMW d.s.)

5× sodium chloride sodium citrate (SSC)
 0.1% Tween 20
 10% low MW dextran sulfate

For 40 mL of solution

10 mL of 20× SSC
 400 µL of 10% Tween 20
 8 mL of 50% low MW dextran sulfate
 Fill up to 40 mL with ultrapure H₂O

5× SSCT

5× sodium chloride sodium citrate (SSC)
 0.1% Tween 20

For 40 mL of solution

10 mL of 20× SSC
 400 µL of 10% Tween 20
 fill up to 40 mL with ultrapure H₂O

1× PBST

1× PBS
 0.1% Tween 20

For 40 mL of solution

10 mL of 10× PBST (0.5% Tween 20)
 200 µL of 10% Tween 20
 Fill up to 40 mL with ultrapure H₂O

50% dextran sulfate

50% dextran sulfate

For 40 mL of solution

20 g of low MW dextran sulfate powder
 Fill up to 40 mL with ultrapure H₂O

S2.3.5 Reagents and supplies

DPBS, no calcium, no magnesium (Life Technologies Cat. # 14190144)
Trypsin-EDTA (0.25%), phenol red (Life Technologies Cat. # 25200072)
16% Formaldehyde (w/v), methanol-free (Life Technologies Cat. # 28908)
10× PBST (Rockland Cat. # MB-075-1000)
10% Tween 20 solution (Bio-Rad Cat. # 161-0781)
Formamide (Deionized) (Ambion Cat. # AM9342)
20× sodium chloride sodium citrate (SSC) (Life Technologies Cat. # 15557-044)
Heparin (Sigma Cat. # H3393)
50% Tween 20 (Life Technologies Cat. # 00-3005)
50× Denhardt's solution (Life Technologies Cat. # 750018)
Dextran sulfate, mol. wt. 6,500-10,000 (Sigma Cat. # D4911)

S2.4 Protocols for bacteria in suspension

S2.4.1 Preparation of fixed bacteria in suspension

1. Grow *E. coli* from streaked plate or frozen glycerol stocks in 2–3 mL of LB media overnight in a 37 °C shaker.
2. Dilute to make a 5 mL liquid culture with $OD_{600} = 0.05$.
3. Incubate in a 37 °C shaker until $OD_{600} \approx 0.5$ (exponential phase).
4. Aliquot 1 mL of cells and centrifuge for 10 min.
NOTE: Centrifugation should be as gentle as possible to pellet cells. For E. coli all centrifugation steps are done at $4000 \times g$.
5. Remove supernatant and re-suspend cells in 750 μL 1 \times phosphate-buffered saline (PBS).
NOTE: Remove all solutions via pipetting throughout the protocol.
6. Add 250 μL of 4% formaldehyde to and incubate overnight at 4 °C.
CAUTION: Use formaldehyde with extreme care as it is a hazardous material.
7. Centrifuge for 10 min and remove supernatant.
8. Re-suspend cells in 150 μL 1 \times PBS.
9. Add 850 μL of 100% MeOH and store at -20 °C before use.

S2.4.2 Buffer recipes for sample preparation

LB media

5 g of Novagen LB Broth Miller powder
Fill up to 200 mL with ultrapure H₂O
Autoclave at 121 °C for 20 min

4% formaldehyde in PBS

4% formaldehyde
1× PBS

For 4 mL of solution

1 mL of 16% formaldehyde
0.4 mL of 10× PBS
Fill up to 4 mL with ultrapure H₂O

NOTE: Avoid using calcium chloride and magnesium chloride in PBS as this leads to increased autofluorescence in the samples.

S2.4.3 Multiplexed in situ HCR v3.0 using split-initiator probes

Detection stage

1. Transfer 150 μL of fixed cells into a 1.5 mL Eppendorf tube.
2. Centrifuge for 5 min and remove supernatant.
3. Wash cells with 500 μL of 1 \times PBST. Centrifuge for 5 min to remove supernatant.
4. Re-suspend the pellet with 400 μL of 30% LMW probe hybridization buffer and pre-hybridize for 1 hr at 37 $^{\circ}\text{C}$.
CAUTION: probe hybridization buffer contains formamide, a hazardous material.
NOTE: Pre-heat probe hybridization buffer to 37 $^{\circ}\text{C}$ before use.
5. In the meantime, prepare probe solution by adding 2 pmol of each probe mixture (odd & even: 1 μL of 2 μM stock per probe mixture) to 100 μL of 30% LMW probe hybridization buffer at 37 $^{\circ}\text{C}$.
6. Add the probe solution directly to the sample to reach a final probe concentration of 4 nM.
7. Incubate the sample overnight at 37 $^{\circ}\text{C}$.
8. Add 1 mL of probe wash buffer to the sample.
CAUTION: probe wash buffer contains formamide, a hazardous material.
NOTE: Pre-heat probe wash buffer to 37 $^{\circ}\text{C}$ before use.
9. Centrifuge for 5 min and remove the wash solution.
10. Re-suspend the cell pellet with 500 μL wash solution.
11. Incubate for 5 min at 37 $^{\circ}\text{C}$ and remove the wash solution by centrifugation for 5 min.
12. Repeat steps 10 and 11 for two additional times but with 10 min incubation.
13. Proceed to hairpin amplification.

Amplification stage

1. Re-suspend the cell pellet with 150 μL of LMW amplification buffer and pre-amplify for 30 min at room temperature.
NOTE: Equilibrate amplification buffer to room temperature before use.
2. Prepare 15 pmol of each fluorescently labeled hairpin by snap cooling 5 μL of 3 μM stock in hairpin storage buffer (heat at 95 $^{\circ}\text{C}$ for 90 seconds and cool to room temperature in a dark drawer for 30 min).
3. Prepare hairpin mixture by adding all snap-cooled hairpins to 100 μL of LMW amplification buffer at room temperature.
4. Add the hairpin mixture directly to the sample to reach a final hairpin concentration of 60 nM.
5. Incubate the sample overnight (>12 h) in the dark at room temperature.
6. Add 1 mL of 5 \times SSCT at room temperature to the sample to dilute the solution.
7. Centrifuge for 5 min and remove the hairpin solution.
8. Re-suspend the cell pellet with 500 μL of 5 \times SSCT and incubate 5 min at room temperature.
9. Centrifuge for 5 min and remove the wash solution.

10. Repeat steps 8 and 9 for two additional times but with a 10 min incubation.
11. Re-suspend the cell pellet in desired buffer and volume.
NOTE: Samples can be stored at 4 °C protected from light before flow cytometry.
12. Filter cells before flow cytometry.

S2.4.4 Buffer recipes for in situ HCR v3.0

Probes, amplifiers, probe hybridization buffer, and probe wash buffer should be stored at -20 °C. Amplification buffer should be stored at 4 °C. Make sure all solutions are well mixed before use.

30% probe hybridization buffer (low MW D. S.)

30% formamide
5× sodium chloride sodium citrate (SSC)
9 mM citric acid (pH 6.0)
0.1% Tween 20
50 µg/mL heparin
1× Denhardt's solution
10% low MW dextran sulfate

For 40 mL of solution

12 mL formamide
10 mL of 20× SSC
360 µL 1 M citric acid, pH 6.0
400 µL of 10% Tween 20
200 µL of 10 mg/mL heparin
800 µL of 50× Denhardt's solution
8 mL of 50% low MW dextran sulfate
Fill up to 40 mL with ultrapure H₂O

30% probe wash buffer

30% formamide
5× sodium chloride sodium citrate (SSC)
9 mM citric acid (pH 6.0)
0.1% Tween 20
50 µg/mL heparin

For 40 mL of solution

12 mL formamide
10 mL of 20× SSC
360 µL 1 M citric acid, pH 6.0
400 µL of 10% Tween 20
200 µL of 10 mg/mL heparin
Fill up to 40 mL with ultrapure H₂O

Amplification buffer (low MW D. S.)

5× sodium chloride sodium citrate (SSC)
0.1% Tween 20
10% low MW dextran sulfate

For 40 mL of solution

10 mL of 20× SSC
400 µL of 10% Tween 20
8 mL of 50% low MW dextran sulfate
Fill up to 40 mL with ultrapure H₂O

5× SSCT

5× sodium chloride sodium citrate (SSC)
0.1% Tween 20

For 40 mL of solution

10 mL of 20× SSC
400 µL of 10% Tween 20
fill up to 40 mL with ultrapure H₂O

1× PBST

1× PBS
0.1% Tween 20

For 40 mL of solution

10 mL of 10× PBST (0.5% Tween 20)
200 µL of 10% Tween 20
Fill up to 40 mL with ultrapure H₂O

50% dextran sulfate

50% dextran sulfate

For 40 mL of solution

20 g of low MW dextran sulfate powder
Fill up to 40 mL with ultrapure H₂O

S2.4.5 Reagents and supplies

LB Broth Miller (Novagen Cat. # 71753-5)
16% Formaldehyde (w/v), methanol-free (Life Technologies Cat. # 28908)
10× PBS (Ambion Cat. # AM9624)
10× PBST (Rockland Cat. # MB-075-1000)
10% Tween 20 solution (Bio-Rad Cat. # 161-0781)
Formamide (Deionized) (Ambion Cat. # AM9342)
20× sodium chloride sodium citrate (SSC) (Life Technologies Cat. # 15557-044)
Heparin (Sigma Cat. # H3393)
50% Tween 20 (Life Technologies Cat. # 00-3005)
50× Denhardt's solution (Life Technologies Cat. # 750018)
Dextran sulfate, mol. wt. 6,500-10,000 (Sigma Cat. # D4911)

S3 Additional studies

S3.1 Validation of split-initiator HCR suppression in vitro and in situ (cf. Figure 2)

Figures S3 and S4 display gel studies measuring split-initiator HCR suppression for amplifiers B1–B5, revealing typical ≈ 60 -fold suppression. Table S9 displays signal-to-signal ratios for the same amplifiers used in situ within whole-mount chicken embryos (imaging) and/or within cultured human or bacterial cells (flow cytometry), revealing typical ≈ 50 -fold suppression.




Figure S3. Test tube validation of split-initiator HCR suppression for amplifiers B1, B2, B4, and B5 (cf. Figure 2). (A) Agarose gel electrophoresis. Reaction conditions: hairpins H1 and H2 at 0.5 μ M each (Lanes 1-7); DNA oligos I1, P1, P2, and/or Target at 5 nM each (lanes noted on the gel); 5 \times SSCT buffer; overnight reaction at room temperature. Hairpins H1 and H2 labeled with Alexa 647 fluorophore (green channel). dsDNA 1 kb ladder pre-stained with SYBR Gold (red channel). (B) Quantification of the polymer band in panel (A).

Organism	Target	Samples	Channel	SIG/SIG ^{odd}	SIG/SIG ^{even}	Table
<i>G. gallus domesticus</i>	<i>EphA4</i>	3 embryos	B2-Alexa647	>800	57 \pm 5	S14
<i>H. sapiens sapiens</i>	<i>Tg(d2eGFP)</i>	55,000 cells	B3-Alexa594	461 \pm 9	22.2 \pm 0.1	S17
<i>H. sapiens sapiens</i>	<i>GAPDH</i>	30,000 cells	B4-Alexa594	55 \pm 5	42.9 \pm 0.6	S18
<i>H. sapiens sapiens</i>	<i>GAPDH</i>	20,000 cells	B5-Alexa488	40.4 \pm 0.8	>3000	S20
<i>H. sapiens sapiens</i>	<i>GAPDH</i>	20,000 cells	B4-Alexa594	67 \pm 2	52 \pm 1	S20
<i>H. sapiens sapiens</i>	<i>PGK1</i>	54,000 cells	B1-Alexa488	>5000	49.0 \pm 0.5	S21
<i>H. sapiens sapiens</i>	<i>PGK1</i>	54,000 cells	B2-Alexa594	41 \pm 1	13 \pm 1	S21
<i>H. sapiens sapiens</i>	<i>GAPDH</i>	18,000 cells	B4-Alexa488	93 \pm 6	91 \pm 14	S23
<i>H. sapiens sapiens</i>	<i>PGK1</i>	18,000 cells	B2-Alexa594	21 \pm 2	18.8 \pm 0.4	S23
<i>E. coli</i>	<i>Tg(eGFP)</i>	18,000 cells	B3-Alexa594	>3000	9.9 \pm 0.5	S19
<i>E. coli</i>	<i>fusA</i>	3,400 cells	B3-Alexa488	40 \pm 20	14 \pm 3	S22
<i>E. coli</i>	<i>fusA</i>	3,400 cells	B2-Alexa594	50 \pm 20	6.2 \pm 0.6	S22
<i>E. coli</i>	<i>fusA</i>	35,000 cells	B3-Alexa488	600 \pm 300	17.2 \pm 1.3	S24
<i>E. coli</i>	<i>icd</i>	35,000 cells	B1-Alexa594	800 \pm 300	85 \pm 4	S24

Table S9. In situ validation of split-initiator HCR suppression.




Figure S4. Test tube validation of split-initiator HCR suppression for amplifier B3 (cf. Figure 2). (A) Agarose gel electrophoresis (Replicate 1 is displayed in Figure 2). Reaction conditions: hairpins H1 and H2 at $0.5 \mu\text{M}$ each (Lanes 1-7); DNA oligos I1, P1, P2, and/or Target at 5 nM each (lanes noted on the gel); $5\times$ SSCT buffer; overnight reaction at room temperature. Hairpins H1 and H2 labeled with Alexa 647 fluorophore (green channel). dsDNA 1 kb ladder pre-stained with SYBR Gold (red channel). (B) Quantification of the polymer band in panel (A).

S3.2 In situ validation of automatic background suppression with split-initiator probes in whole-mount chicken embryos (cf. Figure 3)

The following studies are included:

- **Measurement of background and signal-to-background for unoptimized standard and split-initiator probe sets as a function of probe set size.** Probe sets with 5, 10, or 20 standard probes or split-initiator probe pairs. For standard probes, the probe set size is increased from 5 to 10 to 20 by adding untested probes to a previously validated set of 5 probes Choi *et al.* (2016); each standard probe has a 50-nt target-binding site and carries two HCR initiators (one at each end). Each split-initiator probe pair addresses the nearly identical target subsequence as the corresponding standard probe; each split-initiator probe has a 25-nt target-binding site and carries half an HCR initiator; the two probes within a pair are separated by 2 nt along the target so the overall target binding site for a probe pair is 52 nt. For background comparisons (Figures S5 and S6 and Table S10), the PMT gain is held constant for all experiments to enable comparison of background intensities between experiments. For signal-to-background comparisons (Figures S7 and S8 and Table S11), the PMT gain is adjusted to use the full dynamic range for each probe set.
- **Measurement of background and signal-to-background for standard and split-initiator probes with identical target-binding domains.** For these studies, standard probes are constructed from split-initiator probe pairs as follows. For each split-initiator probe pair in a probe set, the full initiator is shifted onto either the odd probe (full-initiator odd probe + initiator-free even probe) or onto the even probe (initiator-free odd probe + full-initiator even probe). Within each pair, one probe is then a full-initiator standard probe and one probe is a helper probe that contains no initiator. The helper probes are employed to ensure that each standard probe pair (full-initiator probe + helper probe) has the same target-binding capabilities as its analogous split-initiator probe pair. Probe sets with 20 standard probe pairs or split-initiator probe pairs. For background comparisons (Figure S9 and Table S12), the PMT gain is held constant for all experiments to enable comparison of background intensities between experiments. For signal-to-background comparisons (Figure S10 and Table S13), the PMT gain is adjusted to use the full dynamic range for each probe set.
- **Measurement of signal, background, and signal-to-background, background components, and split-initiator HCR suppression.** Measurement of signal, background, and signal-to-background (Figure S11A and Table S14A) using the methods of Section S1.4.2. Measurement of background components (AF, NSA, NSD; Figure S11B and Table S14B) using the methods of Section S1.4.3. Measurement of split-initiator HCR suppression (Figure S11C and Table S14C) using the methods of Section S1.4.4.

S3.2.1 Measurement of background and signal-to-background for unoptimized standard and split-initiator probe sets as a function of probe set size




Figure S5. Measurement of background for unoptimized standard probe sets as a function of probe set size (cf. Figure 3A). Confocal images collected with the microscope PMT gain adjusted to avoid saturating pixels using 20 standard probes (row 3). Protocol: in situ HCR v2.0 (Section S8 of Choi *et al.* (2016)). Target mRNA: *Sox10*. Probe set: 5, 10, or 20 standard probes. Amplifier: B3-Alexa647. Whole-mount chicken embryos fixed stage HH 11.




Figure S6. Measurement of background for unoptimized split-initiator probe sets as a function of probe set size (cf. Figure 3A). Confocal images collected with the microscope PMT gain adjusted to avoid saturating pixels using 20 standard probes (Figure S5, row 3). Protocol: in situ HCR v3.0 (Section S2.1). Target mRNA: *Sox10*. Probe set: 5, 10, or 20 split-initiator probe pairs. Amplifier: B3-Alexa647. Whole-mount chicken embryos fixed stage HH 11.

Probe type	Probe set size	BACK
Standard (v2.0)	5	210 ± 30
	10	1160 ± 60
	20	1500 ± 100
Split-initiator (v3.0)	5	29 ± 1
	10	26 ± 3
	20	28 ± 1

Table S10. Estimated background for standard and split-initiator probes sets as a function of probe set size (cf. Figure 3A). Mean ± standard error, $N = 3$ replicate embryos. Analysis based on rectangular regions depicted in Figures S5 and S6 using methods of Section S1.4.2.




Figure S7. Measurement of signal and background for unoptimized standard probe sets as a function of probe set size (cf. Figure 3B). (A) Confocal images collected with the microscope PMT gain optimized for each probe set. (B) Pixel intensity histograms for Signal + Background (pixels within solid boundary) and Background (pixels within dashed boundary) per embryo in panel A. Protocol: in situ HCR v2.0 (Section S8 of Choi *et al.* (2016)). Target mRNA: *Sox10*. Probe set: 5, 10, or 20 standard probes. Amplifier: B3-Alexa647. Whole-mount chicken embryos fixed stage HH 11.




Figure S8. Measurement of signal and background for unoptimized split-initiator probe sets as a function of probe set size (cf. Figure 3B). (A) Confocal images collected with the microscope PMT gain optimized for each probe set. (B) Pixel intensity histograms for Signal + Background (pixels within solid boundary) and Background (pixels within dashed boundary) per embryo in panel A. Protocol: in situ HCR v3.0 (Section S2.1). Target mRNA: *Sox10*. Probe set: 5, 10, or 20 split-initiator probe pairs. Amplifier: B3-Alexa647. Whole-mount chicken embryos fixed stage HH 11.

Probe type	Probe set size	BACK	SIG+BACK	SIG	SIG/BACK
Standard (v2.0)	5	200.3 ± 1.1	1770 ± 70	1570 ± 70	7.8 ± 0.3
	10	540 ± 40	2170 ± 40	1630 ± 60	3.0 ± 0.2
	20	720 ± 10	2200 ± 200	1400 ± 200	2.0 ± 0.3
Split-initiator (v3.0)	5	43.7 ± 0.6	1270 ± 70	1230 ± 70	28.2 ± 1.7
	10	34.0 ± 1.2	1480 ± 30	1450 ± 30	42.5 ± 1.8
	20	37 ± 3	2460 ± 40	2420 ± 40	65 ± 5

Table S11. Estimated signal-to-background for standard and split-initiator probes sets as a function of probe set size (cf. Figure 3B). For this signal-to-background study, the microscope PMT gain was optimized for each probe set so raw background and signal values should only be compared within row. Mean ± standard error, $N = 3$ replicate embryos. Analysis based on rectangular regions depicted in Figures S7 and S8 using methods of Section S1.4.2.

S3.2.2 Measurement of background and signal-to-background for standard and split-initiator probes with identical target-binding domains




Figure S9. Measurement of background for standard and split-initiator probes with identical target-binding domains. (A) Confocal images collected with the microscope PMT gain adjusted to avoid saturating pixels using full-initiator odd probe sets (top row). Protocol: in situ HCR v3.0 (Section S2.1). Target mRNA: *Sox10*. Probe sets (20 probe pairs per probe set): full-initiator odd probes + initiator-free even probes (top row), initiator-free odd probes + full-initiator even probes (middle row), split-initiator odd and even probes (bottom row). Amplifier: B3-Alexa647. Whole-mount chicken embryos fixed stage HH 10.

Sample	BACK
Full-initiator probes (odd)	910 ± 80
Full-initiator probes (even)	330 ± 40
Split-initiator probes	19.9 ± 0.9

Table S12. Estimated background for standard and split-initiator probes with identical target-binding domains. Mean ± standard error, $N = 3$ replicate embryos. Analysis based on rectangular regions depicted in Figure S9 using methods of Section S1.4.2.




Figure S10. Measurement of signal and background for standard and split-initiator probes with identical target-binding domains. (A) Confocal images collected with the microscope PMT gain optimized for each probe set. (B) Pixel intensity histograms for Signal + Background (pixels within solid boundary) and Background (pixels within dashed boundary) per embryo in panel A. Protocol: in situ HCR v3.0 (Section S2.1). Target mRNA: *Sox10*. Probe sets (20 probe pairs per probe set): full-initiator odd probes + initiator-free even probes (top row), initiator-free odd probes + full-initiator even probes (middle row), split-initiator odd and even probes (bottom row). Amplifier: B3-Alexa647. Whole-mount chicken embryos fixed stage HH 10.

Sample	BACK	SIG+BACK	SIG	SIG/BACK
Full-initiator probes (odd)	910 ± 80	1230 ± 110	320 ± 140	0.4 ± 0.2
Full-initiator probes (even)	330 ± 40	1610 ± 150	1280 ± 160	3.9 ± 0.6
Split-initiator probes	26.4 ± 0.7	1870 ± 110	1840 ± 110	70 ± 5

Table S13. Estimated signal-to-background for standard and split-initiator probes with identical target-binding domains. Mean ± standard error, $N = 3$ replicate embryos. Analysis based on rectangular regions depicted in Figure S10 using methods of Section S1.4.2.

S3.2.3 Measurement of signal, background, signal-to-background, background components, and split-initiator HCR suppression




Figure S11. Measurement of signal and background, background components, and split-initiator HCR suppression. Confocal image: fluorescence merged with bright field. (A) Signal and background: use experiment of Type 1 in Table S5A (odd probes + even probes + hairpins) to measure SIG+BACK (region of high expression) and BACK (region of no/low expression). (B) Background components: use experiment of Type 2 in Table S5B (no probes, hairpins only) to measure NSA+AF (region of high expression); use experiment of Type 3 (no probes, no hairpins) to measure AF (region of high expression). (C) Split-initiator HCR suppression: use experiment of Type 4 in Table S5C (odd probes, hairpins) to measure SIG^{odd}+BACK^{odd} (region of high expression) and BACK^{odd} (region of no/low expression); use experiment of Type 5 in Table S5C (even probes, hairpins) to measure SIG^{even}+BACK^{even} (region of high expression) and BACK^{even} (region of no/low expression). Left: confocal images collected with the microscope PMT gain optimized to avoid saturating pixels using the full method (top row). Right: pixel intensity histograms for a region of high expression (pixels within solid boundary) and/or low/no expression (pixels within dashed boundary) per embryo. Protocol: in situ HCR v3.0 (Section S2.1). Target mRNA: *EphA4*. Probe set: 20 split-initiator probe pairs. Amplifier: B2-Alexa647. Whole-mount chicken embryos fixed stage HH 11.

	Channel		Reagents		Expression			
	B2-Alexa647		Probes	Hairpins	region	Figure		
A	SIG+NSD+NSA+AF = SIG+BACK		1260	± 30	odd + even	✓	high	S11C
	NSD+NSA+AF = BACK		19	± 1	odd + even	✓	low/no	S11C
	SIG		1240	± 30				
	SIG/BACK		67	± 4				
B	NSA+AF		8.1	± 0.9		✓	high	S11B
	AF		6.97	± 0.08			high	S11A
	NSA		1.2	± 0.9				
	NSD		10.5	± 1.3				
C	NSD ^{odd} +NSA+AF = BACK ^{odd}		11.0	± 1.0	odd	✓	low/no	S11D
	NSD ^{even} +NSA+AF = BACK ^{even}		10.0	± 0.6	even	✓	low/no	S11E
	SIG ^{odd} +BACK ^{odd}		11.5	± 1.1	odd	✓	high	S11D
	SIG ^{even} +BACK ^{even}		31.8	± 1.7	even	✓	high	S11E
	NSD ^{odd}		2.8	± 1.4				
	NSD ^{even}		1.9	± 1.1				
	SIG ^{odd}		< 1.5					
	SIG ^{even}		21.8	± 1.8				
	SIG/SIG ^{odd}		> 800					
	SIG/SIG ^{even}		57	± 5				

Table S14. Estimated signal-to-background, background components, and split-initiator HCR suppression. (A) Signal-to-background (SIG/BACK) based on methods of Section S1.4.2. (B) Background components (AF, NSA, NSD) based on methods of Section S1.4.3. (C) Split-initiator HCR suppression (SIG/SIG^{odd}, SIG/SIG^{even}) based on methods of Section S1.4.4. Mean \pm standard error, $N = 3$ replicate embryos. Analysis based on rectangular regions depicted in Figure S11.

S3.3 Multiplexed 4-channel mRNA imaging with high signal-to-background in whole-mount chicken embryos (cf. Figure 4)




Figure S12. Measurement of signal and background for multiplexed 4-channel mRNA imaging (cf. Figure 4). (A) Individual channels of 4-channel confocal images. For each of three replicate embryos, a representative optical section was selected for each channel based on the expression depth of the corresponding target mRNA. (B) Pixel intensity histograms for Signal + Background (pixels within solid boundary) and Background (pixels within dashed boundary). Ch1: target mRNA *FoxD3*, probe set with 12 split-initiator probe pairs, amplifier B4-Alexa488. Ch2: target mRNA *Dmbx1*, probe set with 20 split-initiator probe pairs, amplifier B1-Alexa514. Ch 3: target mRNA *Sox10*, probe set with 20 split-initiator probe pairs, amplifier B3-Alexa546. Ch4: *EphA4*, probe set with 20 split-initiator probe pairs, amplifier B2-Alexa647. Whole-mount chicken embryos fixed stage HH 10.

Target mRNA	BACK	SIG+BACK	SIG	SIG/BACK
<i>FoxD3</i>	36 ± 3	1270 ± 50	1230 ± 50	34 ± 3
<i>Dmbx1</i>	16.4 ± 0.2	730 ± 20	710 ± 20	43.4 ± 1.0
<i>Sox10</i>	34.0 ± 0.2	2030 ± 80	2000 ± 80	59 ± 3
<i>EphA4</i>	35 ± 3	980 ± 70	950 ± 70	27 ± 3

Table S15. Estimated signal-to-background for multiplexed 4-channel mRNA imaging. Mean ± standard error, $N = 3$ replicate embryos. Analysis based on rectangular regions depicted in Figure S12 using methods of Section S1.4.2.

S3.4 qHCR imaging: analog mRNA relative quantitation with subcellular resolution in whole-mount chicken embryos (cf. Figure 5)

S3.4.1 Testing for a crowding effect

In order to perform multiplexed quantitative imaging using HCR, it is important that there is not a crowding effect in which amplification polymers tethered to one target molecule affect the signal intensity for a different target molecule. To test for a possible crowding effect, we imaged two target mRNAs that are highly expressed in the same cells (*EphA4* and *Egr2*) individually (1-target studies) and also simultaneously (2-target studies) within whole-mount chicken embryos. Figure S13 compares the signal intensity distributions for 1-target and 2-target studies, revealing similar intensity distributions whether targets were detected alone or together, suggesting that there is not a significant crowding effect (either antagonistic or synergistic).




Figure S13. Comparison of signal intensity distributions for individual and simultaneous imaging of *EphA4* and *Egr2*. (A) Raw voxel intensity scatter plot: *Egr2* channel vs *EphA4* channel. (B) Raw voxel intensity histogram for *EphA4* channel. (C) Raw voxel intensity histogram for *Egr2* channel. In panels B and C, solid lines denote average histograms over 3 replicate embryos while symbols denote individual histograms (1 histogram per replicate). Orange data: signal plus background for *EphA4* and *Egr2* (Figure S14). Cyan data: signal plus background for *EphA4* and background for *Egr2* (Figure S15). Blue data: background for *EphA4* and signal plus background for *Egr2* (Figure S16). Black data (near origin): background for *EphA4* and *Egr2* (Figure S17). Voxel size: $2.1 \times 2.1 \times 2.7 \mu\text{m}$. Whole-mount wildtype chicken embryos fixed stage HH 10.




Figure S14. Characterizing signal plus background for *EphA4* and *Egr2* in a 2-target experiment. (A) Individual channels from 2-channel confocal images depicting regions used to estimate signal plus background. *EphA4* channel: 20 probe pairs and amplifier B1-Alexa546. *Egr2* channel: 20 probe pairs and amplifier B3-Alexa647. For each of 3 replicate embryos, a representative optical section was selected based on the expression depth of the target mRNAs. Pixel size: $0.4 \times 0.4 \mu\text{m}$. (B) Raw voxel intensity scatter plots for the selected regions of panel A representing signal plus background for *EphA4* and *Egr2*. (C) Raw voxel intensity histograms for the selected regions of panel A representing signal plus background for *EphA4* and *Egr2*. Same microscope settings used for all replicates in Figures S14–S17. Voxel size: $2.1 \times 2.1 \times 2.7 \mu\text{m}$. Whole-mount wildtype chicken embryos fixed stage HH 10.




Figure S15. Characterizing signal plus background for *EphA4* in a 1-target experiment. (A) Individual channels from 2-channel confocal images depicting regions used to estimate signal plus background for *EphA4* and background for *Egr2*. *EphA4* channel: 20 probe pairs and amplifier B1-Alexa546. *Egr2* channel: no probes, no amplifier. For each of 3 replicate embryos, a representative optical section was selected based on the expression depth of the target mRNA. Pixel size: $0.4 \times 0.4 \mu\text{m}$. (B) Raw voxel intensity scatter plots for the selected regions of panel A representing signal plus background for *EphA4* and background for *Egr2*. (C) Raw voxel intensity histograms for the selected regions of panel A representing signal plus background for *EphA4* and background for *Egr2*. Same microscope settings used for all replicates in Figures S14–S17. Voxel size: $2.1 \times 2.1 \times 2.7 \mu\text{m}$. Whole-mount wildtype chicken embryos fixed stage HH 10.




Figure S16. Characterizing signal plus background for *Egr2* in a 1-target experiment. (A) Individual channels from 2-channel confocal images depicting regions used to estimate background for *EphA4* and signal plus background for *Egr2*. *EphA4* channel: no probes, no amplifier. *Egr2* channel: 20 probe pairs and amplifier B3-Alexa647. For each of 3 replicate embryos, a representative optical section was selected based on the expression depth of the target mRNA. Pixel size: $0.4 \times 0.4 \mu\text{m}$. (B) Raw voxel intensity scatter plots for the selected regions of panel A representing background for *EphA4* and signal plus background for *Egr2*. (C) Raw voxel intensity histograms for the selected regions of panel A representing background for *EphA4* and signal plus background for *Egr2*. Same microscope settings used for all replicates in Figures S14–S17. Voxel size: $2.1 \times 2.1 \times 2.7 \mu\text{m}$. Whole-mount wildtype chicken embryos fixed stage HH 10.




Figure S17. Characterizing background for *EphA4* and *Egr2*. Individual channels from 2-channel confocal images depicting regions used to estimate background using the standard HCR v3.0 in situ protocol (Section S2.1) omitting probes (BACK \approx AF + NSA; see Section S1.4 for definitions). For each of 3 replicate embryos, a representative optical section was selected at approximately the depth where *EphA4* and *Egr2* are expressed. Same microscope settings used for all replicates in Figures S14–S17. Pixel size: $0.4 \times 0.4 \mu\text{m}$. (B) Raw voxel intensity scatter plots for the selected region of panel A. (C) Raw voxel intensity histograms for the scatter plots of panel B. Voxel size: $2.1 \times 2.1 \times 2.7 \mu\text{m}$. Whole-mount wildtype chicken embryos fixed stage HH 10.


S3.4.2 Redundant 2-channel detection of *Dmbx1*

Figure S18. Redundant 2-channel detection of *Dmbx1*. (A) Confocal images: individual channels and merge. Solid boundaries denote regions of high expression; dashed boundaries denote regions of no/low expression. Pixel size: $0.2 \times 0.2 \mu\text{m}$. Probe sets: 20 split-initiator probe pairs per channel. Amplifiers: B1-Alexa546 (Ch1) and B2-Alexa647 (Ch2). Whole-mount chicken embryos fixed stage HH 10. (B) Raw voxel intensity scatter plots representing signal plus background for voxels within solid boundaries of panel A. Voxel size: $2.1 \times 2.1 \times 2.7 \mu\text{m}$. Dashed lines represent BOT and TOP values (Table S16) used to normalize data for panel C using methods of Section S1.4.5. (C) Normalized voxel intensity scatter plots representing estimated normalized signal (Pearson correlation coefficient, r).


S3.4.3 Redundant 2-channel detection of *EphA4*

Figure S19. Redundant 2-channel detection of *EphA4* (cf. Figure 5). (A) Confocal images: individual channels and merge. Solid boundaries denote regions of high expression; dashed boundaries denote regions of no/low expression. Pixel size: $0.2 \times 0.2 \mu\text{m}$. Probe sets: 20 split-initiator probe pairs per channel. Amplifiers: B1-Alexa546 (Ch1) and B2-Alexa647 (Ch2). Whole-mount chicken embryos fixed stage HH 10. (B) Raw voxel intensity scatter plots representing signal plus background for voxels within solid boundaries of panel A. Voxel size: $2.1 \times 2.1 \times 2.7 \mu\text{m}$. Dashed lines represent BOT and TOP values (Table S16) used to normalize data for panel C using methods of Section S1.4.5. (C) Normalized voxel intensity scatter plots representing estimated normalized signal (Pearson correlation coefficient, r).

Target mRNA	Channel	BACK	SIG+BACK	SIG	SIG/BACK	BOT	TOP
<i>Dmbx1</i>	Alexa546	23.9 ± 1.1	840 ± 80	810 ± 80	34 ± 4	24	3266
<i>Dmbx1</i>	Alexa647	35 ± 5	770 ± 70	730 ± 70	21 ± 3	35	3040
<i>EphA4</i>	Alexa546	28.7 ± 1.1	1720 ± 10	1693 ± 13	59 ± 2	29	3995
<i>EphA4</i>	Alexa647	29.7 ± 1.1	1490 ± 10	1455 ± 10	48.9 ± 1.8	30	3855

Table S16. Estimated signal-to-background for redundant 2-channel detection of *Dmbx1* and *EphA4*. Mean ± standard error, $N = 3$ replicate embryos. Analysis based on rectangular regions depicted in Figure S18A and S19A using methods of Section S1.4.2. BOT and TOP values used to calculate normalized voxel intensities for scatter plots of Figures 5C, S18C, and S19C using methods of Section S1.4.5.

S3.5 In situ validation of automatic background suppression with split-initiator probes for mRNA flow cytometry with cultured human and bacterial cells

The methods of Sections S1.5.2 and S1.5.3 are used to measure:

- signal, background, and signal-to-background (Figures S20A–S22A and Tables S17A–S19A).
- background components (AF, NSA, NSD; Figures S20A–S22A and Tables S17B–S19B).
- split-initiator HCR suppression (Figures S20B–S22B and Tables S17C–S19C).

Additional measurements of these quantities are provided in the 2-channel experiments of Figures S24–S28 and Tables S20–S24.

S3.5.1 Measurement of signal, background, signal-to-background, background components, and split-initiator HCR suppression for *d2eGFP* transgenic target in HEK cells




Figure S20. Measurement of signal and background, background components, and split-initiator HCR suppression for *d2eGFP* transgenic target in HEK cells (cf. Figure 6A). (A) Signal and background: use experiments of Types 1a and 1b in Table S7A (odd + even probes, hairpins) to measure SIG+BACK (GFP+ cells) and BACK (WT cells). Background components: use experiment of Type 2 in Table S7B (no probes, with hairpins) to measure NSA+AF (GFP+ cells); use experiment of Type 3 (no probes, no hairpins) to measure AF (GFP+ cells). (B) Split-initiator HCR suppression: use experiment of Types 4a in Table S7C (odd probes, hairpins) to measure SIG^{odd}+BACK^{odd} (GFP+ cells); use experiment of Type 5a in Table S7C (even probes, hairpins) to measure SIG^{even}+BACK^{even} (GFP+ cells). Distribution of single-cell fluorescence intensities. Protocol: in situ HCR v3.0 (Section S2.3). Probe set: 12 split-initiator probe pairs. Amplifier: B3-Alexa594. Sample: 55,000 HEK cells in suspension (GFP+ or WT).

Quantity	Channel	Reagents		
	B3-Alexa594	Probes	Hairpins	Cell type
A SIG+NSD+NSA+AF = SIG+BACK	13 220 ± 60	odd + even	✓	GFP+
	NSD+NSA+AF = BACK	128.5 ± 0.2	odd + even	✓
SIG	13 090 ± 60			
SIG/BACK	101.8 ± 0.5			
B NSA+AF	120.7 ± 0.5		✓	GFP+
	AF	104.7 ± 0.2	✓	GFP+
NSA	16.0 ± 0.5			
NSD	7.8 ± 0.5			
C SIG ^{odd} +BACK ^{odd}	149.1 ± 0.3	odd	✓	GFP+
	SIG ^{even} +BACK ^{even}	710 ± 3	even	✓
SIG ^{odd}	28.4 ± 0.6			
SIG ^{even}	589 ± 3			
SIG/SIG ^{odd}	461 ± 9			
SIG/SIG ^{even}	22.2 ± 0.1			

Table S17. Estimated signal-to-background, background components, and split-initiator HCR suppression for *d2eGFP* transgenic target in HEK cells (cf. Figure 6A). (A) Signal-to-background (SIG/BACK). (B) Background components (AF, NSA, NSD). (C) Split-initiator HCR suppression (SIG/SIG^{odd}, SIG/SIG^{even}). The signal estimates SIG^{odd} and SIG^{even} are calculated using the background approximation BACK^{odd} = BACK^{even} ≈ NSA+AF, which leads to lower bounds on SIG/SIG^{odd} and SIG/SIG^{even}. Mean ± standard error, $N = 55,000$ cells. Analysis based on single-cell intensities of Figure S20 using methods of Section S1.5.2.

S3.5.2 Measurement of signal, background, signal-to-background, background components, and split-initiator HCR suppression for *GAPDH* endogenous target in HEK cells




Figure S21. Measurement of signal and background, background components, and split-initiator HCR suppression for *GAPDH* endogenous target in HEK cells. (A) Signal and background: use experiments of Types 1a and 1b in Table S8A to measure SIG+BACK (even + odd probes, hairpins) and BACK (Tg(odd) + Tg(even) probes, hairpins). Background components: use experiment of Type 2 in Table S8B (no probes, with hairpins) to measure NSA+AF; use experiment of Type 3 (no probes, no hairpins) to measure AF. (B) Split-initiator HCR suppression: use experiment of Types 4a and 4b in Table S8C to measure SIG^{odd}+BACK^{odd} (odd probes, hairpins) and BACK^{odd} (Tg(odd) probes, hairpins); use experiments of Types 5a and 5b in Table S8C to measure SIG^{even}+BACK^{even} (even probes, hairpins) and BACK^{even} (Tg(even) probes, hairpins). Distribution of single-cell fluorescence intensities. Protocol: in situ HCR v3.0 (Section S2.3). Probe set: 10 split-initiator probe pairs. Amplifier: B4-Alexa594. Sample: 30,000 HEK cells in suspension (WT).

Quantity	Channel		Reagents		Cell type
	B4-Alexa594		Probes	Hairpins	
A SIG+NSD+NSA+AF = SIG+BACK NSD+NSA+AF = BACK	4775	± 15	odd + even	✓	WT
	414.0	± 0.9	Tg(odd) + Tg(even)	✓	WT
SIG	4362	± 15			
SIG/BACK	10.55	± 0.04			
B NSA+AF	353.7	± 0.8		✓	WT
	AF	304.0	± 0.7	✓	WT
NSA	50	± 1			
NSD	60	± 1			
C SIG ^{odd} +NSD ^{odd} +NSA+AF = SIG ^{odd} +BACK ^{odd} NSD ^{odd} +NSA+AF = BACK ^{odd} SIG ^{even} +NSD ^{even} +NSA+AF = SIG ^{even} +BACK ^{even} NSD ^{even} +NSA+AF = BACK ^{even}	450	± 7	odd	✓	WT
	371.2	± 0.8	Tg(odd)	✓	WT
	499	± 1	even	✓	WT
	397.3	± 0.8	Tg(even)	✓	WT
NSD ^{odd}	18	± 1			
NSD ^{even}	44	± 1			
SIG ^{odd}	79	± 7			
SIG ^{even}	102	± 1			
SIG/SIG ^{odd}	55	± 5			
SIG/SIG ^{even}	42.9	± 0.6			

Table S18. Estimated signal-to-background, background components, and split-initiator HCR suppression for *GAPDH* endogenous target in HEK cells. (A) Signal-to-background (SIG/BACK). (B) Background components (AF, NSA, NSD). (C) Split-initiator HCR suppression (SIG/SIG^{odd}, SIG/SIG^{even}). Mean ± standard error, $N = 30,000$ cells. Analysis based on single-cell intensities of Figure S21 using methods of Section S1.5.3.

S3.5.3 Measurement of signal, background, signal-to-background, background components, and split-initiator HCR suppression for *eGFP* transgenic target in *E. coli*




Figure S22. Measurement of signal and background, background components, and split-initiator HCR suppression for *eGFP* transgenic target in *E. coli* (cf. Figure 6A). (A) Signal and background: use experiments of Types 1a and 1b in Table S7A (odd + even probes, hairpins) to measure SIG+BACK (GFP+ cells) and BACK (WT cells). Background components: use experiment of Type 2 in Table S7B (no probes, with hairpins) to measure NSA+AF (GFP+ cells); use experiment of Type 3 (no probes, no hairpins) to measure AF (GFP+ cells). (B) Split-initiator HCR suppression: use experiment of Type 4a in Table S7C (odd probes, hairpins) to measure SIG^{odd}+BACK^{odd} (GFP+ cells); use experiment of Type 5a in Table S7C (even probes, hairpins) to measure SIG^{even}+BACK^{even} (GFP+ cells). Distribution of single-cell fluorescence intensities. Protocol: in situ HCR v3.0 (Section S2.4). Probe set: 12 split-initiator probe pairs. Amplifier: B3-Alexa594. Sample: 18,000 *E. coli* in suspension (WT: K12 MG1655; GFP+: K12 MG1655 pUA66-sdhC expressing *gfpmut2*).

Quantity	Channel	Reagents		
	B3-Alexa594	Probes	Hairpins	Cell type
A SIG+NSD+NSA+AF = SIG+BACK	3330 ± 10	odd + even	✓	GFP+
	NSD+NSA+AF = BACK	120 ± 20	odd + even	✓
SIG	3200 ± 30			
SIG/BACK	26 ± 5			
B NSA+AF	72.5 ± 0.7		✓	GFP+
	AF	55.7 ± 0.7	✓	GFP+
NSA	17 ± 1			
NSD	50 ± 20			
C SIG ^{odd} +BACK ^{odd}	71.3 ± 0.7	odd	✓	GFP+
	SIG ^{even} +BACK ^{even}	395 ± 15	even	✓
SIG ^{odd}	< 1			
SIG ^{even}	320 ± 10			
SIG/SIG ^{odd}	> 3000			
SIG/SIG ^{even}	9.9 ± 0.5			

Table S19. Estimated signal-to-background, background components, and split-initiator HCR suppression for *eGFP* transgenic target in *E. coli* (cf. Figure 6A). (A) Signal-to-background (SIG/BACK). (B) Background components (AF, NSA, NSD). (C) Split-initiator HCR suppression (SIG/SIG^{odd}, SIG/SIG^{even}). The signal estimates SIG^{odd} and SIG^{even} are calculated using the background approximation $BACK^{odd} = BACK^{even} \approx NSA+AF$, which leads to lower bounds on SIG/SIG^{odd} and SIG/SIG^{even}. Mean ± standard error, $N = 18,000$ cells. Analysis based on single-cell intensities of Figure S22 using methods of Section S1.5.2.

S3.6 qHCR flow cytometry: analog mRNA relative quantitation for high-throughput expression profiling of human and bacterial cells (cf. Figure 6)

S3.6.1 Testing for a crowding effect

In order to perform multiplexed quantitative flow cytometry using HCR, it is important that there is not a crowding effect in which amplification polymers tethered to one target molecule affect the signal intensity for a different target molecule. To test for a possible crowding effect, we analyzed two highly-expressed target mRNAs (*GAPDH* and *ACTB*) individually (1-target studies) and also simultaneously (2-target studies) within HEK cells. Figure S23 compares the signal intensity distributions for 1-target and 2-target studies, revealing similar intensity distributions whether targets were detected alone or together, suggesting that there is not a significant crowding effect (either antagonistic or synergistic).




Figure S23. Comparison of signal intensity distributions for individual and multiplexed floHCR of *GAPDH* and *ACTB*. (A) Raw single-cell fluorescence intensity scatter plots: *GAPDH* channel vs *ACTB* channel. (B) Single-cell fluorescence intensity histogram for *GAPDH* channel. (C) Single-cell fluorescence intensity histogram for *ACTB* channel. Orange data: signal plus background for *GAPDH* and *ACTB*. Cyan data: signal plus background for *GAPDH* and autofluorescence for *ACTB*. Blue data: background for *ACTB* and signal plus autofluorescence for *GAPDH*. Black data: autofluorescence for *GAPDH* and *ACTB*. Sample: 65,000 HEK cells in suspension (WT).


S3.6.2 Redundant 2-channel detection of *GAPDH* endogenous target in HEK cells

Figure S24. Measurement of signal and background, background components, and split-initiator HCR suppression for redundant 2-channel detection of *GAPDH* endogenous target (cf. Figure 6B). (A) Signal and background: use experiment of Type 1a in Table S8A to measure SIG+BACK (even + odd probes, hairpins). Background components: use experiment of Type 2 in Table S8B (no probes, with hairpins) to measure NSA+AF; use experiment of Type 3 (no probes, no hairpins) to measure AF. Split-initiator HCR suppression: use experiment of Type 4a in Table S8C (odd probes, hairpins) to measure SIG^{odd}+BACK^{odd} (odd probes, hairpins); use experiment of Type 5a in Table S8C to measure SIG^{even}+BACK^{even} (even probes, hairpins). Single-cell fluorescence intensity histograms for Ch1 and Ch2. (B) Raw single-cell fluorescence intensity scatter plots representing signal plus background. Dashed lines represent BOT and TOP values (Table S20) used to normalize data for panel C using methods of Section S1.5.4 (outliers excluded from normalized scatter plots marked with circles). (C) Normalized single-cell fluorescence intensity scatter plots representing estimated normalized signal (Pearson correlation coefficient, r). Protocol: in situ HCR v3.0 (Section S2.3). Probe sets: 10 split-initiator probe pairs per channel. Amplifiers: B5-Alexa488 (Ch1) and B4-Alexa594 (Ch2). Sample: 20,000 HEK cells in suspension (WT).

Quantity	Channel		Reagents		Cell type
	Ch1: B5-Alexa488	Ch2: B4-Alexa594	Probes	Hairpins	
A SIG+BACK	870 ± 5	3105 ± 14	odd + even	✓	WT
SIG	808 ± 5	2904 ± 14			
SIG/BACK	12.86 ± 0.08	14.43 ± 0.08			
B NSA+AF	62.8 ± 0.2	201.2 ± 0.6		✓	WT
AF	57.4 ± 0.2	166.5 ± 0.5		✓	WT
NSA	5.4 ± 0.3	34.7 ± 0.8			
C SIG ^{odd} +BACK ^{odd}	82.8 ± 0.3	244.5 ± 0.8	odd	✓	WT
SIG ^{even} +BACK ^{even}	63.0 ± 0.2	256.6 ± 0.8	even	✓	WT
SIG ^{odd}	20.0 ± 0.4	43 ± 1			
SIG ^{even}	<0.3	55 ± 1			
SIG/SIG ^{odd}	40.4 ± 0.8	67 ± 2			
SIG/SIG ^{even}	>3000	52 ± 1			
D BOT	62.8	201.2			
TOP	5265.2	19 049.4			

Table S20. Estimated signal-to-background, background components, and split-initiator HCR suppression for redundant 2-channel detection of *GAPDH* endogenous target (cf. Figure 6B). (A) Signal-to-background (SIG/BACK). (B) Background components (AF, NSA). (C) Split-initiator HCR suppression (SIG/SIG^{odd}, SIG/SIG^{even}). The signal estimates SIG, SIG^{odd} and SIG^{even} are calculated using the background approximation $BACK = BACK^{odd} = BACK^{even} \approx NSA+AF$. Mean ± standard error, $N = 20,000$ cells. Analysis based on single-cell intensities of Figure S24 using methods of Section S1.5.3. (D) BOT and TOP values used to calculate normalized single-cell intensities for scatter plots of Figures 6B and S24C using methods of Section S1.5.4.

S3.6.3 Redundant 2-channel detection of *PGK1* endogenous target in HEK cells




Figure S25. Measurement of signal and background, background components, and split-initiator HCR suppression for redundant 2-channel detection of *PGK1* endogenous target. (A) Signal and background: use experiment of Type 1a in Table S8A to measure SIG+BACK (even + odd probes, hairpins). Background components: use experiment of Type 2 in Table S8B (no probes, with hairpins) to measure NSA+AF; use experiment of Type 3 (no probes, no hairpins) to measure AF. Split-initiator HCR suppression: use experiment of Type 4a in Table S8C (odd probes, hairpins) to measure SIG^{odd}+BACK^{odd} (odd probes, hairpins); use experiment of Type 5a in Table S8C to measure SIG^{even}+BACK^{even} (even probes, hairpins). Single-cell fluorescence intensity histograms for Ch1 and Ch2. (B) Raw single-cell fluorescence intensity scatter plots representing signal plus background. Dashed lines represent BOT and TOP values (Table S21) used to normalize data for panel C using methods of Section S1.5.4 (outliers excluded from normalized scatter plots marked with circles). (C) Normalized single-cell fluorescence intensity scatter plots representing estimated normalized signal (Pearson correlation coefficient, r). Protocol: in situ HCR v3.0 (Section S2.3). Probe sets: 18 split-initiator probe pairs per channel. Amplifiers: B1-Alexa488 (Ch1) and B2-Alexa594 (Ch2). Sample: 54,000 HEK cells in suspension (WT).

Quantity	Channel		Reagents		Cell type
	Ch1: B1-Alexa488	Ch2: B2-Alexa594	Probes	Hairpins	
A SIG+BACK	4145 ± 11	1528 ± 4	odd + even	✓	WT
SIG	3843 ± 11	1248 ± 4			
SIG/BACK	12.72 ± 0.04	4.47 ± 0.02			
B NSA+AF	302.1 ± 0.5	279.4 ± 0.5		✓	WT
AF	289.3 ± 0.5	220.1 ± 0.4		✓	WT
NSA	12.9 ± 0.7	59.3 ± 0.6			
C SIG ^{odd} +BACK ^{odd}	301.3 ± 0.5	309.5 ± 0.6	odd	✓	WT
SIG ^{even} +BACK ^{even}	380.5 ± 0.6	374 ± 7	even	✓	WT
SIG ^{odd}	< 0.7	30.1 ± 0.8			
SIG ^{even}	78.4 ± 0.8	94 ± 7			
SIG/SIG ^{odd}	>5000	41 ± 1			
SIG/SIG ^{even}	49.0 ± 0.5	13 ± 1			
D BOT	302.1	279.4			
TOP	35 538.8	11 847.7			

Table S21. Estimated signal-to-background, background components, and split-initiator HCR suppression for redundant 2-channel detection of *PGKI* endogenous target. (A) Signal-to-background (SIG/BACK). (B) Background components (AF, NSA). (C) Split-initiator HCR suppression (SIG/SIG^{odd}, SIG/SIG^{even}). The signal estimates SIG, SIG^{odd} and SIG^{even} are calculated using the background approximation BACK = BACK^{odd} = BACK^{even} ≈ NSA+AF. Mean ± standard error, $N = 54,000$ cells. Analysis based on single-cell intensities of Figure S25 using methods of Section S1.5.3. (D) BOT and TOP values used to calculate normalized single-cell intensities for scatter plots of Figure S25C using methods of Section S1.5.4.


S3.6.4 Redundant 2-channel detection of *fusA* endogenous target in *E. coli*

Figure S26. Measurement of signal and background, background components, and split-initiator HCR suppression for redundant 2-channel detection of *fusA* endogenous target (cf. Figure 6B). (A) Signal and background: use experiment of Type 1a in Table S8A to measure SIG+BACK (even + odd probes, hairpins). Background components: use experiment of Type 2 in Table S8B (no probes, with hairpins) to measure NSA+AF; use experiment of Type 3 (no probes, no hairpins) to measure AF. Split-initiator HCR suppression: use experiment of Type 4a in Table S8C (odd probes, hairpins) to measure SIG^{odd}+BACK^{odd} (odd probes, hairpins); use experiment of Type 5a in Table S8C to measure SIG^{even}+BACK^{even} (even probes, hairpins). Single-cell fluorescence intensity histograms for Ch1 and Ch2. (B) Raw single-cell fluorescence intensity scatter plots representing signal plus background. Dashed lines represent BOT and TOP values (Table S22) used to normalize data for panel C using methods of Section S1.5.4 (outliers excluded from normalized scatter plots marked with ellipses). (C) Normalized single-cell fluorescence intensity scatter plots representing estimated normalized signal (Pearson correlation coefficient, r). Protocol: in situ HCR v3.0 (Section S2.4). Probe sets: 18 split-initiator probe pairs per channel. Amplifiers: B3-Alexa488 (Ch1) and B2-Alexa594 (Ch2). Sample: 3,400 *E. coli* K12 MG1655 in suspension (WT).

Quantity	Channel		Reagents		Cell type
	Ch1: B3-Alexa488	Ch2: B2-Alexa594	Probes	Hairpins	
A SIG+BACK	13 100 ± 300	15 800 ± 300	odd + even	✓	WT
SIG	13 000 ± 300	15 700 ± 300			
SIG/BACK	99 ± 3	135 ± 7			
B NSA+AF	130 ± 3	116 ± 5		✓	WT
AF	126 ± 7	120 ± 10		✓	WT
NSA	<7	<14			
C SIG ^{odd} +BACK ^{odd}	500 ± 100	400 ± 100	odd	✓	WT
SIG ^{even} +BACK ^{even}	1100 ± 200	2600 ± 200	even	✓	WT
SIG ^{odd}	300 ± 100	300 ± 100			
SIG ^{even}	900 ± 200	2500 ± 200			
SIG/SIG ^{odd}	40 ± 20	50 ± 20			
SIG/SIG ^{even}	14 ± 3	6.2 ± 0.6			
D BOT	130.3	116			
TOP	150 951.6	183 947.3			

Table S22. Estimated signal-to-background, background components, and split-initiator HCR suppression for redundant 2-channel detection of *fusA* endogenous target (cf. Figure 6B). (A) Signal-to-background (SIG/BACK). (B) Background components (AF, NSA). (C) Split-initiator HCR suppression (SIG/SIG^{odd}, SIG/SIG^{even}). The signal estimates SIG, SIG^{odd} and SIG^{even} are calculated using the background approximation $BACK = BACK^{odd} = BACK^{even} \approx NSA+AF$. Mean \pm standard error, $N = 3,400$ cells. Analysis based on single-cell intensities of Figure S26 using methods of Section S1.5.3. (D) BOT and TOP values used to calculate normalized single-cell intensities for scatter plots of Figures 6B and S26C using methods of Section S1.5.4.


S3.6.5 Multiplexed 2-channel detection of *GAPDH* and *PGK1* endogenous targets in HEK cells

Figure S27. Measurement of signal and background, background components, and split-initiator HCR suppression for multiplexed 2-channel detection of *GAPDH* and *PGK1* endogenous targets. (A) Single-cell fluorescence intensity histograms for Ch1 and Ch2. (B) Raw single-cell fluorescence intensity scatter plots for Ch1 vs Ch2. Signal and background: use experiment of Type 1a in Table S8A to measure SIG+BACK (even + odd probes, hairpins). Background components: use experiment of Type 2 in Table S8B (no probes, with hairpins) to measure NSA+AF; use experiment of Type 3 (no probes, no hairpins) to measure AF. (C) Split-initiator HCR suppression: use experiment of Type 4a in Table S8C (odd probes, hairpins) to measure SIG^{odd}+BACK^{odd} (odd probes, hairpins); use experiment of Type 5a in Table S8C to measure SIG^{even}+BACK^{even} (even probes, hairpins). Protocol: in situ HCR v3.0 (Section S2.3). Ch1: target mRNA *GAPDH*, probe set with 10 split-initiator probe pairs, amplifier B4-Alexa488. Ch2: target mRNA *PGK1*, probe set with 18 split-initiator probe pairs, amplifier B2-Alexa594. Sample: 18,000 HEK cells in suspension (WT).

Quantity	Channel		Reagents		Cell type
	Ch1: B4-Alexa488	Ch2: B2-Alexa594	Probes	Hairpins	
A SIG+BACK	6980 ± 30	2073 ± 8	odd + even	✓	WT
SIG	6760 ± 30	1806 ± 8			
SIG/BACK	29.6 ± 0.1	6.77 ± 0.05			
B NSA+AF	228.2 ± 0.7	266.6 ± 1.5		✓	WT
AF	219.7 ± 0.6	198.4 ± 0.6		✓	WT
NSA	8.4 ± 0.9	68 ± 2			
C SIG ^{odd} +BACK ^{odd}	301 ± 5	355 ± 9	odd	✓	WT
SIG ^{even} +BACK ^{even}	303 ± 14	362.8 ± 1.2	even	✓	WT
SIG ^{odd}	73 ± 5	88 ± 9			
SIG ^{even}	74 ± 14	96 ± 2			
SIG/SIG ^{odd}	93 ± 6	21 ± 2			
SIG/SIG ^{even}	91 ± 14	18.8 ± 0.4			

Table S23. Estimated signal-to-background, background components, and split-initiator HCR suppression for multiplexed 2-channel detection of *GAPDH* and *PGKI* endogenous targets. (A) Signal-to-background (SIG/BACK). (B) Background components (AF, NSA). (C) Split-initiator HCR suppression (SIG/SIG^{odd}, SIG/SIG^{even}). The signal estimates SIG, SIG^{odd} and SIG^{even} are calculated using the background approximation $BACK = BACK^{odd} = BACK^{even} \approx NSA+AF$. Mean ± standard error, $N = 18,000$ cells. Analysis based on single-cell intensities of Figure S27 using methods of Section S1.5.3.

S3.6.6 Multiplexed 2-channel detection of *fusA* and *icd* endogenous targets in *E. coli*




Figure S28. Measurement of signal and background, background components, and split-initiator HCR suppression for multiplexed 2-channel detection of *fusA* and *icd* endogenous targets. (A) Single-cell fluorescence intensity histograms for Ch1 and Ch2. (B) Raw single-cell fluorescence intensity scatter plots for Ch1 vs Ch2. Signal and background: use experiment of Type 1a in Table S8A to measure SIG+BACK (even + odd probes, hairpins). Background components: use experiment of Type 2 in Table S8B (no probes, with hairpins) to measure NSA+AF; use experiment of Type 3 (no probes, no hairpins) to measure AF. (C) Split-initiator HCR suppression: use experiment of Type 4a in Table S8C (odd probes, hairpins) to measure SIG^{odd}+BACK^{odd} (odd probes, hairpins); use experiment of Type 5a in Table S8C to measure SIG^{even}+BACK^{even} (even probes, hairpins). Protocol: in situ HCR v3.0 (Section S2.4). Ch1: target mRNA *fusA*, probe set with 18 split-initiator probe pairs, amplifier B3-Alexa488. Ch2: target mRNA *icd*, probe set with 20 split-initiator probe pairs, amplifier B1-Alexa594. Sample: 35,000 *E. coli* K12 MG1655 in suspension (WT).

Quantity	Channel		Reagents		Cell type
	Ch1: B3-Alexa488	Ch2: B1-Alexa594	Probes	Hairpins	
A SIG+BACK	1756 ± 9	2533 ± 12	odd + even	✓	WT
SIG	1673 ± 9	2470 ± 12			
SIG/BACK	20.1 ± 0.2	38.9 ± 0.5			
B NSA+AF	83.2 ± 0.7	63.5 ± 0.7		✓	WT
AF	82.3 ± 0.7	60.1 ± 0.7		✓	WT
NSA	1 ± 1	3.4 ± 0.9			
C SIG ^{odd} +BACK ^{odd}	86 ± 1	66.6 ± 0.8	odd	✓	WT
SIG ^{even} +BACK ^{even}	180 ± 8	93 ± 1	even	✓	WT
SIG ^{odd}	3 ± 1	3 ± 1			
SIG ^{even}	97 ± 8	29 ± 1			
SIG/SIG ^{odd}	600 ± 300	800 ± 300			
SIG/SIG ^{even}	17.2 ± 1.3	85 ± 4			

Table S24. Estimated signal-to-background, background components, and split-initiator HCR suppression for multiplexed 2-channel detection of *fusA* and *icd* endogenous targets. (A) Signal-to-background (SIG/BACK). (B) Background components (AF, NSA). (C) Split-initiator HCR suppression (SIG/SIG^{odd}, SIG/SIG^{even}). The signal estimates SIG, SIG^{odd} and SIG^{even} are calculated using the background approximation BACK = BACK^{odd} = BACK^{even} ≈ NSA+AF. Mean ± standard error, $N = 35,000$ cells. Analysis based on single-cell intensities of Figure S28 using methods of Section S1.5.3.

S3.7 dHCR imaging: digital mRNA absolute quantitation in an anatomical context (cf. Figure 7)

S3.7.1 Redundant 2-channel detection of single *BRAF* mRNAs in HEK cells using in situ HCR v3.0




Figure S29. Redundant 2-channel detection of single *BRAF* mRNAs in HEK cells using in situ HCR v3.0 (cf. Figure 7A). Confocal images: individual channels and merge (without and with DAPI nuclear stain). Maximum intensity projection in the axial direction over $7.14 \mu\text{m}$ (17 focal planes). Pixel size: $0.062 \times 0.062 \mu\text{m}$. Probe sets: 23 split-initiator probe pairs per channel. Amplifiers: B3-Alexa647 (Ch1) and B4-Alexa546 (Ch2). Red circles: dots detected in Ch1. Green circles: dots detected in Ch2. Yellow circles: dots detected in both channels.

	Dots		Colocalized dots	Colocalization fractions	
	N_1	N_2	N_{12}	C_1	C_2
Replicate 1	129	136	110	0.85	0.81
Replicate 2	63	65	53	0.84	0.82
Replicate 3	170	170	144	0.85	0.85
Mean				0.85 ± 0.003	0.82 ± 0.01

Table S25. Dot colocalization fractions for redundant 2-channel detection of single *BRAF* mRNAs in HEK cells using in situ HCR v3.0 (cf. Figure 7A). Mean \pm standard error, $N = 3$ replicate samples. Analysis based on the images of Figure S29 using the methods of Section S1.4.6 with the settings in Table S6.

S3.7.2 Redundant 2-channel detection of single *Dmbx1* mRNAs in whole-mount chicken embryos using in situ HCR v3.0




Figure S30. Redundant 2-channel detection of single *Dmbx1* mRNAs in whole-mount chicken embryos using in situ HCR v3.0 (cf. Figure 7B). Confocal images: individual channels and merge. Maximum intensity projection in the axial direction over 5.04-23.52 μm (12, 54, 56 focal planes for replicates 1, 2, 3 depending on sample thickness). Pixel size: $0.099 \times 0.099 \mu\text{m}$. Probe sets: 25 split-initiator probe pairs per channel. Amplifiers: B2-Alexa647 (Ch1) and B1-Alexa594 (Ch2). Embryos fixed stage HH 8. Red circles: dots detected in Ch1. Green circles: dots detected in Ch2. Yellow circles: dots detected in both channels.

	Dots		Colocalized dots	Colocalization fractions	
	N_1	N_2	N_{12}	C_1	C_2
Replicate 1	403	417	364	0.90	0.87
Replicate 2	992	990	794	0.80	0.80
Replicate 3	526	539	448	0.85	0.83
Mean				0.85 ± 0.03	0.84 ± 0.02

Table S26. Dot colocalization fractions for redundant 2-channel detection of single *Dmbx1* mRNAs in whole-mount chicken embryos using in situ HCR v3.0 (cf. Figure 7B). Mean \pm standard error, $N = 3$ replicate embryos. Analysis based on the images of Figure S30 using the methods of Section S1.4.6 with the settings in Table S6.

S3.7.3 Redundant 2-channel detection of single *kdrl* mRNAs in whole-mount zebrafish embryos using in situ HCR v2.0 (Shah *et al.*, 2016)




Figure S31. Redundant 2-channel detection of single *kdrl* mRNAs in whole-mount zebrafish embryos using in situ HCR v2.0. Spinning disk confocal images: individual channels and merge from Shah *et al.* (2016). Maximum intensity projection in the axial direction over 13 μm (39 focal planes). Pixel size: $0.217 \times 0.217 \mu\text{m}$. Probe sets: 39 standard probes per channel, each incorporating a 30-nt target-binding domain and a full HCR initiator. Amplifiers: B3-Alexa647 (Ch1) and B2-Alexa546 (Ch2). Embryos fixed 27 hpf. Red circles: dots detected in Ch1. Green circles: dots detected in Ch2. Yellow circles: dots detected in both channels.

	Dots		Colocalized dots	Colocalization fractions	
	N_1	N_2	N_{12}	C_1	C_2
Replicate 1	139	132	79	0.57	0.60
Replicate 2	220	215	113	0.51	0.53
Replicate 3	243	245	91	0.37	0.37
Mean				0.49 ± 0.06	0.50 ± 0.07

Table S27. Dot colocalization fractions for redundant 2-channel detection of single *kdrl* mRNAs in whole-mount zebrafish embryos using in situ HCR v3.0. Mean \pm standard error, $N = 3$ replicate embryos. Analysis based on the images of Figure S31 using the methods of Section S1.4.6 with the settings in Table S6.

S4 Probe sequences

Target mRNA sequences were obtained from the National Center for Biotechnology Information (NCBI) (McEntyre & Ostell, 2002) with the exception of *d2eGFP* (pd2EGFP-1, Clontech, Cat. #6008-1). Spatial and temporal expression information for whole-mount chicken embryos were obtained from the Gallus Expression in Situ Hybridization Analysis (GEISHA) (Bell *et al.*, 2004; Darnell *et al.*, 2007). Within a given probe set, each DNA standard probe or split-initiator probe pair initiates the same DNA HCR amplifier. Sequences are listed 5' to 3'. Probes are numbered consecutively moving along a target mRNA. For redundant detection experiments, two probe sets are used with each probe set taking alternating probe pairs from along the target (this leads to non-consecutive numbers within each probe set).

S4.1 Standard probes for Figures 3, S5, and S7

Organism: *G. gallus domesticus*

Target mRNA: **SRY (sex determining region Y)-box 10 (Sox10)**

Probe set: **5, 10, or 20 probes (each carrying 2 HCR initiators)**

HCR amplifier: **B3-Alexa647**

#	Initiator I1	Spacer	Probe Sequence (50 nt)	Spacer	Initiator I2
1	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	CATggACCCgTCACTCCATgTCTTgAgTCTTCCTCATCTAagAAggCCAAT	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
2	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	CCAgCagggATCAAagATTCAgTgTgAAATCTTAaggCaggACTgCTg	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
3	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	CgggCTATgAAATgAgAAAaggCTAAggCTgACAgTgCAgTTCCTgAATCC	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
4	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	TTCACgTTTTCAgCagACACAgTCAAATgCTgAggAgCAAaggACCTggT	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
5	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	ATTggAACACATCTgTggTgTggCAAgtgCATggTAGCTTCTTgTgTgC	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
6	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	AgAggCggggAgAAAagCTATAgCgTgCAgCTgTgAAAATCAgCAaggAA	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
7	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	ATAAAATCCATgCAggAAggggTgTgggATTAAACAgATgggACAggggg	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
8	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	gATggCgATAATgTgATgAACAAACgAgCAgTgATgTACACCCATCggC	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
9	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	ATCCACgAgAgTATCTTTCCATCCTgAgTgAAAgTAaggAgggAggTgCTg	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
10	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	ACCCgTAgAAggTCCCAACAACATCTCTCTgATCAgTTgTCAggAgTC	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
11	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	CCggCgAgAggCAgTggTggTCTTCAgAACCCACTgggCTCATCTCCACC	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
12	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	CCTCgCCCTgCgTggCCTTgCCATTTTTCgCCTCCgTggCTggTACTTg	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
13	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	TTCTTgTAgTgAgCCTggATAgAggCAgCCCCgCAgCCTCCCCCTCCAC	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
14	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	TgCCCATCggACATgggTgACCCTTCCCCAggATgCCTgTggTCCAggTg	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
15	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	TTAggggTggTgggAggAgTgggAgggCCgTggCTCTgACCTgAAgAgTg	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
16	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	CCCCAagggAACgCCCTTCTCgCTTggAgTCAgCTTTgCCTgCCTgCAgC	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
17	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	ggCCTgggTggCCggCgTgTCCgTTgggTggCAggTATTggTCAAATTCg	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
18	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	TCCATgCTgCTTggAgATCCAaggCTgAgTgTCCACTggCCgCAgCCAagg	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
19	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	AACCCTgTgAAggCTgCAgCTCCTgTCCCAgTgCCCCCTgTTCTCCCTC	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC
20	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	gggTCCAgTCATAgCCgCTCAGCACCTggCTgACCgCTTACggATgCAg	TAAAA	AAAgTCTAATCCgTCCCTgCCTCTATATCTCCACTC

S4.2 Split-initiator probes for Figures 3, S6, S8, S9, and S10

Organism: *G. gallus domesticus*

Target mRNA: **SRY (sex determining region Y)-box 10 (Sox10)**

Probe set: **5, 10, or 20 split-initiator probe pairs (each probe carries half an HCR initiator)**

HCR amplifier: **B3-Alexa647**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	gTCCCTgCCTCTATATCT	TT	gTCTTCCTCATCTAgAAggCCAATA	2	CCATggACCCgTCACTCCATgTCTT	TT	CCACTCAACTTTAACCCg
3	gTCCCTgCCTCTATATCT	TT	TgTgAATCTTAggCAggACTgCTgC	4	TCCAgCAgggATCAAgATTTCATgCA	TT	CCACTCAACTTTAACCCg
5	gTCCCTgCCTCTATATCT	TT	gCTgACAgTgCagTTCCTgAATCCT	6	ACgggCTATgAAATgAgAAAggCTA	TT	CCACTCAACTTTAACCCg
7	gTCCCTgCCTCTATATCT	TT	TgCTggAggAgCAaggACCTggTCT	8	TTCACgTTTTCAgCAGACACAgTCA	TT	CCACTCAACTTTAACCCg
9	gTCCCTgCCTCTATATCT	TT	CAAgTgCATggTAGCTTCTTggTg	10	AATATTggAACACATCTgggTgTT	TT	CCACTCAACTTTAACCCg
11	gTCCCTgCCTCTATATCT	TT	AgCTgTgAAAATCAgCAaggAAgCA	12	gAggCggggAgAAAAGCTATAgCgT	TT	CCACTCAACTTTAACCCg
13	gTCCCTgCCTCTATATCT	TT	gggATTAAACAgATgggACAggggg	14	TTATAAAATCCATgCAggAAggggT	TT	CCACTCAACTTTAACCCg
15	gTCCCTgCCTCTATATCT	TT	AgCAGTgATgTACACCCCATCggCC	16	AgATggCgATAATgTgATgAACAAA	TT	CCACTCAACTTTAACCCg
17	gTCCCTgCCTCTATATCT	TT	TgAgTgAAAgTAggAgggAggTgCT	18	TATATCCACgAgAgTATCTTTCCAT	TT	CCACTCAACTTTAACCCg
19	gTCCCTgCCTCTATATCT	TT	CTCTCTgATCAgTTgTCAggAgTCA	20	TACCCgTTAgAAggTCCCACAACAC	TT	CCACTCAACTTTAACCCg
21	gTCCCTgCCTCTATATCT	TT	gAACCCACTgggCTCATCTCCACCT	22	CCCggCgAgAggCAgTggTggTCTT	TT	CCACTCAACTTTAACCCg
23	gTCCCTgCCTCTATATCT	TT	TTCCgCCTCCgTggCTggTACTTgT	24	CCCTCgCCCTgCgTggCCTTgCCAT	TT	CCACTCAACTTTAACCCg
25	gTCCCTgCCTCTATATCT	TT	AgCCCCgCCAgCCTCCCCCTCCACC	26	ATTCTTgTAgTgAgCCTggATAgAg	TT	CCACTCAACTTTAACCCg
27	gTCCCTgCCTCTATATCT	TT	CCCAGgATgCCTgTggTCCAggTgg	28	gTgCCCATCggACATgggTgACCCT	TT	CCACTCAACTTTAACCCg
29	gTCCCTgCCTCTATATCT	TT	gCCgTggCTCTgACCTgAAgAgTgC	30	CTTAggggTggTgggAggAgTgggA	TT	CCACTCAACTTTAACCCg
31	gTCCCTgCCTCTATATCT	TT	gAgTCAgCTTTgCCTgCCTgCAgCT	32	TCCCCAAgggAACgCCCTTCTCgCT	TT	CCACTCAACTTTAACCCg
33	gTCCCTgCCTCTATATCT	TT	ggTggCAggTATTggTCAAATTCgT	34	TggCCTgggTggCCggCgTgTCCgT	TT	CCACTCAACTTTAACCCg
35	gTCCCTgCCTCTATATCT	TT	AgTgTCCACTggCCgCAgCCAgggC	36	CTCCATgCTgCTTggAgATCCAggC	TT	CCACTCAACTTTAACCCg
37	gTCCCTgCCTCTATATCT	TT	CCCAGTgCCCCCTgTTCTCCCTCC	38	AAACCCgTgAAggCTgCAgCTCCT	TT	CCACTCAACTTTAACCCg
39	gTCCCTgCCTCTATATCT	TT	TggCTgACCgCTCACggATgCAgA	40	AgggTCCAgTCATAgCCgCTCAgCA	TT	CCACTCAACTTTAACCCg

S4.3 Standard probes for Figures S9 and S10

Organism: *G. gallus domesticus*

Target mRNA: **SRY (sex determining region Y)-box 10 (Sox10)**

Probe set: **20 probe pairs (odd probes carry full HCR initiator, even probes carry no initiator)**

HCR amplifier: **B3-Alexa647**

Odd #	Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)
1	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	gTCTTCCTCATCTAgAAggCCAATA	2	CCATggACCCgTCACTCCATgTCTT
3	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	TgTgAATCTTAggCAggACTgCTgC	4	TCCAgCAgggATCAAgATTCATgCA
5	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	gCTgACAgTgCagTTCCTgAATCCT	6	ACgggCTATgAAATgAgAAAggCTA
7	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	TgCTggAggAgCAaggACCTggTCT	8	TTCACgTTTTCAgCagACACAgTCA
9	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	CAAgTgCATggTAGCTTTCTTggTg	10	AATATTggAACCACATCTgggTgTT
11	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	AgCTgTgAAAATCAgCAaggAAgCA	12	gAggCggggAgAAAAgCTATAgCgT
13	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	gggATTAAACAgATgggACAggggg	14	TTATAAAATCCATgCaggAAggggT
15	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	AgCagTgATgTACACCCATCggCC	16	AgATggCgATAATgTgATgAACAAA
17	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	TgAgTgAAAgTAGgAgggAggTgCT	18	TATATCCACgAgAgTATCTTTCCAT
19	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	CTCTCTgATCagTTgTCAggAgTCA	20	TACCCgTTAgAAggTCCACAAACAC
21	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	gAACCCACTgggCTCATCTCCACCT	22	CCCggCgAgAggCagTggTggTCTT
23	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	TTCCgCCTCCgTggCTggTACTTgT	24	CCCTCgCCCTgCgTggCCTTgCCAT
25	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	AgCCCCgCCAgCCTCCCCCTCCACC	26	ATTCTTgTAGTgAgCCTggATAgAg
27	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	CCCaggATgCCTgTggTCCAggTgg	28	gTgCCCATCggACATgggTgACCCT
29	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	gCCgTggCTCTgACCTgAAgAgTgC	30	CTAgggggTggTgggAggAgTgggA
31	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	gAgTCAgCTTTgCCTgCCTgCagCT	32	TCCCCAagggAACgCCCTTCTCgCT
33	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	ggTggCaggTATTggTCAAAATTCgT	34	TggCCTgggTggCCgCgTgTCCgT
35	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	AgTgTCCACTgCCgCagCCAaggC	36	CTCCATgCTgCTTggAgATCCAggC
37	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	CCCAGTgCCCCCTgTTCTCCCTCC	38	AAACCCTgTgAAggCTgCagCTCCT
39	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg	TACAA	TggCTgACCgCTTACggATgCagA	40	AgggTCCAgTCATAgCCgCTCagCA

Organism: *G. gallus domesticus*Target mRNA: **SRY (sex determining region Y)-box 10 (Sox10)**Probe set: **20 probe pairs (odd probes carry no initiator, even probes carry full HCR initiator)**HCR amplifier: **B3-Alexa647**

Odd #	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	Initiator I1
1	gTCTTCCTCATCTAgAAggCCAATA	2	CCATggACCCgTCACTCCATgTCTT	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
3	TgTgAATCTTAggCAggACTgCTgC	4	TCCAgCagggATCAAgATTCATgCA	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
5	gCTgACAgTgCAgTTCCTgAATCCT	6	ACgggCTATgAAATgAgAAAggCTA	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
7	TgCTggAggAgCAAggACCTggTCT	8	TTCACgTTTTCAgCAgACACAgTCA	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
9	CAAgtgCATggTAgCTTTCTTggTg	10	AATATTggAACCACATCTgggTgTT	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
11	AgCTgTgAAAATCAgCAAaggAAgCA	12	gAggCggggAgAAAAGCTATAgCgT	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
13	gggATTAAACAgATgggACAgggggg	14	TTATAAAATCCATgCAggAAggggT	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
15	AgCAgTgATgTACACCCCATCggCC	16	AgATggCgATAATgTgATgAACAAA	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
17	TgAgTgAAAgTAggAgggAggTgCT	18	TATATCCACgAgAgTATCTTTCCAT	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
19	CTCTCTgATCagTTgTCAggAgTCA	20	TACCCgTTAgAAggTCCCACAACAC	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
21	gAACCCACTgggCTCATCTCCACCT	22	CCCggCgAgAggCAgTggTggTCTT	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
23	TTCCgCCTCCgTggCTggTACTTgT	24	CCCTCgCCCTgCgTggCCCTgCCAT	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
25	AgCCCCgCCAgCCTCCCCCTCCACC	26	ATTCTTgTAgTgAgCCTggATAgAg	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
27	CCCAGgATgCCTgTggTCCAggTgg	28	gTgCCCATCggACATgggTgACCCT	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
29	gCCgTggCTCTgACCTgAAgAgTgC	30	CTTAGgggTggTgggAggAgTgggA	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
31	gAgTCAgCTTTgCCTgCCTgCAgCT	32	TCCCCAAGggAACgCCCTTCTCgCT	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
33	ggTggCAggTATTggTCAAATTCgT	34	TggCCTgggTggCCggCgTgTCCgT	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
35	AgTgTCCACTggCCgCAgCAaggC	36	CTCCATgCTgCTTggAgATCCAaggC	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
37	CCCAGTgCCCCCTgTTCTCCCTCC	38	AAACCCTgTgAAggCTgCAgCTCCT	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg
39	TggCTgACCgCTTACAggATgCAgA	40	AgggTCCAgTCATAgCCgCTCAgCA	TACTT	gTCCCTgCCTCTATATCTCCACTCAACTTTAACCCg

S4.4 Split-initiator probes for Figure S11

Organism: *G. gallus domesticus*

Target mRNA: **EPH receptor A4** (*EphA4*)

Probe set: **20 split-initiator probe pairs** (each probe carries half an HCR initiator)

HCR amplifier: **B2-Alexa647**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	CCTCgTAAATCCTCATCA	AA	gCgCCgACggggACCCcgCCATgC	2	CAGACgCCgACgAggAgCgggAggA	AA	ATCATCCAgTAAACCgCC
5	CCTCgTAAATCCTCATCA	AA	CCCAGCTCCCCtgcACcAgCggg	6	CCTCCTCCAgCgggCTcGcATCC	AA	ATCATCCAgTAAACCgCC
9	CCTCgTAAATCCTCATCA	AA	TgACTgggCTCCATCACATTgCAAA	10	ATCCAATCAgTTCgTAACCAATTAT	AA	ATCATCCAgTAAACCgCC
13	CCTCgTAAATCCTCATCA	AA	TCCTTgTCgTTgTTTgATTcATAgT	14	gCAAActggCTCTCTcGAATAAAAC	AA	ATCATCCAgTAAACCgCC
17	CCTCgTAAATCCTCATCA	AA	AgTgggCACTTCTTATAgAAgACAC	18	ggAAActgTgCCAggTTCgAACTg	AA	ATCATCCAgTAAACCgCC
21	CCTCgTAAATCCTCATCA	AA	ACgTCCTTCTCTCCgAgTTgTTgA	22	CCATCTgCCCCgAgTACATTTTTg	AA	ATCATCCAgTAAACCgCC
25	CCTCgTAAATCCTCATCA	AA	ATTTTgCAAgCTTggCATTCACCAT	26	TCTgTTgAgAgCgCCTTgTAgTATC	AA	ATCATCCAgTAAACCgCC
29	CCTCgTAAATCCTCATCA	AA	AAgAAgCCCCgATCACAggTgCAgg	30	ATggATgCAGCATCATTTTCTgCTC	AA	ATCATCCAgTAAACCgCC
33	CCTCgTAAATCCTCATCA	AA	gCgCTCCACTCCAAgTTCActgACg	34	TCgTCCcGTCCCTTgTTCTgTg	AA	ATCATCCAgTAAACCgCC
37	CCTCgTAAATCCTCATCA	AA	AAATgTACACCActgCCACAggACC	38	gTTTTCAgCCCgTTCTgCTgggggC	AA	ATCATCCAgTAAACCgCC
41	CCTCgTAAATCCTCATCA	AA	TgCTTggACACTCCATTCACTgCCC	42	gACACAgCTTggTCCTggCTggggT	AA	ATCATCCAgTAAACCgCC
45	CCTCgTAAATCCTCATCA	AA	gCAACgCTgTgCCTcGTATCTCTT	46	ggCCTgTCAggTTCcAgCCAggCCA	AA	ATCATCCAgTAAACCgCC
49	CCTCgTAAATCCTCATCA	AA	gCTgTCTTCACAATgCgATAgCTgC	50	AAACCTTTgATgTCAgTATTCCTgg	AA	ATCATCCAgTAAACCgCC
53	CCTCgTAAATCCTCATCA	AA	AActCAAACggCCCACTgAAgTCTC	54	ATgggggAAggAACTgTgTTAgTTg	AA	ATCATCCAgTAAACCgCC
57	CCTCgTAAATCCTCATCA	AA	gCTgCAATgAgAATgACCACgAgAA	58	TTgCTgCgCCTCTgCTgATgACAA	AA	ATCATCCAgTAAACCgCC
61	CCTCgTAAATCCTCATCA	AA	TATgTAAAggATCCACATATgTTC	62	TCCCTCACAgCTTgATTTggATCCT	AA	ATCATCCAgTAAACCgCC
65	CCTCgTAAATCCTCATCA	AA	CATTTAgTAACAACgCCTTCCAAgT	66	TACTCAgTTATgATCATTACTggTT	AA	ATCATCCAgTAAACCgCC
69	CCTCgTAAATCCTCATCA	AA	TCCcGATgCACATAgCTCATgTCAg	70	TTgACCgTATgTTTCgAgCAgCTA	AA	ATCATCCAgTAAACCgCC
73	CCTCgTAAATCCTCATCA	AA	CCAAATTTAggTCTgTCgCTgCgTT	74	AgTTTgTCCAgCATgTTgACAATCT	AA	ATCATCCAgTAAACCgCC
77	CCTCgTAAATCCTCATCA	AA	ggggAgCTgggATCCAgCAGggCTg	78	CTgACAgAAACAACCgCCgAgAACT	AA	ATCATCCAgTAAACCgCC

S4.5 Split-initiator probes for Figures 4 and S12

Organism: *G. gallus domesticus*

Target mRNA: forkhead box D3 (*FoxD3*)

Probe set: 12 split-initiator probe pairs (each probe carries half an HCR initiator)

HCR amplifier: B4-Alexa488

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	CCTCAACCTACCTCCAAC	AA	CgCCgCCCgATAgAgTCATCCCCgC	2	gCgCggTCTggCCggACATATCgCT	AT	TCTCACCATATTCgCTTC
3	CCTCAACCTACCTCCAAC	AA	CgTCgATATCCACgTCCTCggCCgC	4	CCggCgCgTCgTCTCCCTCgCCCAC	AT	TCTCACCATATTCgCTTC
5	CCTCAACCTACCTCCAAC	AA	CCCgCAggTgCTCCTgCTgTgCCg	6	AgCCCTgCATCATgAgCgCCgTCTg	AT	TCTCACCATATTCgCTTC
7	CCTCAACCTACCTCCAAC	AA	AgggCCCggCCAgCCCgTAggCgCC	8	CggggggCAgCCCgTAggggCggCC	AT	TCTCACCATATTCgCTTC
9	CCTCAACCTACCTCCAAC	AA	gCAgAgCggCggggTgCgggTAggC	10	gCCCgACgggCgggATgTAggggTA	AT	TCTCACCATATTCgCTTC
11	CCTCAACCTACCTCCAAC	AA	gCAgCgggCACgCgggCggCAgCAT	12	CTTTgCggCTCAgCTCgCCCgACgg	AT	TCTCACCATATTCgCTTC
13	CCTCAACCTACCTCCAAC	AA	ggCTgggCCCgAgCTgCgCgTTgAA	14	CCCCAAACTgCTgAgCTgCAgCTg	AT	TCTCACCATATTCgCTTC
15	CCTCAACCTACCTCCAAC	AA	gCTCggATTTCACgATggAgCCCgC	16	CgATgCTgAACgAggggCggCTgCT	AT	TCTCACCATATTCgCTTC
17	CCTCAACCTACCTCCAAC	AA	TggCggCggggCCgCCgATgATgTT	18	ggAAAgTCTgCgCgCTgggCgCCgA	AT	TCTCACCATATTCgCTTC
19	CCTCAACCTACCTCCAAC	AA	CCgACTgCACggTgACgggCggCCg	20	ACgCCAgCggCTggTgCgCCACAg	AT	TCTCACCATATTCgCTTC
21	CCTCAACCTACCTCCAAC	AA	gCgCgATggCCgCggTggTCCTggC	22	TgATgTTggTAggCACgCTgAggAT	AT	TCTCACCATATTCgCTTC
23	CCTCAACCTACCTCCAAC	AA	CCgTTTCCAgAgATACgTCCgggg	24	AAATAAAAACCCgAAAgCgACCTC	AT	TCTCACCATATTCgCTTC

Organism: *G. gallus domesticus*Target mRNA: **diencephalon/mesencephalon homeobox 1 (*Dmbx1*)**Probe set: **20 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B1-Alexa514**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	gAggAggggCagCAAACgg	AA	CgCTCAgTgAgTTCATggCATgCAG	2	CTgCCTgCTggTgCaggTTgTACAT	TA	gAAgAgTCTTCCTTTACg
5	gAggAggggCagCAAACgg	AA	AAATgATgTCAgCCAgTCTCTCCgC	6	ggTgCTgggATCCATAACgTgCCTC	TA	gAAgAgTCTTCCTTTACg
9	gAggAggggCagCAAACgg	AA	AAATgATgTCAgCCAgTCTCTCCgC	10	gCTTCTgCagCTgCTCCTTCTggAg	TA	gAAgAgTCTTCCTTTACg
13	gAggAggggCagCAAACgg	AA	CagTgTCAggTATTgTggACTgggT	14	CCTCACTgggATgCTCTgAgTCTT	TA	gAAgAgTCTTCCTTTACg
17	gAggAggggCagCAAACgg	AA	CCTCCTCTCTgTCAgTCTgTCCTC	18	TAgCCTCATCCAAggTgCTCTAAA	TA	gAAgAgTCTTCCTTTACg
21	gAggAggggCagCAAACgg	AA	CgCTgATgggAgACTCTgATTTTgg	22	CACTgCTAgATgAAggAgTCACggT	TA	gAAgAgTCTTCCTTTACg
25	gAggAggggCagCAAACgg	AA	CTgCCATgTgCTggCggAATTgCTC	26	AggAgTAAAggACCAAgTTgTTggT	TA	gAAgAgTCTTCCTTTACg
29	gAggAggggCagCAAACgg	AA	gACAgTgCagggAgCTCagAggCgC	30	CgTgggAgAgAgATTggTAgTACgA	TA	gAAgAgTCTTCCTTTACg
33	gAggAggggCagCAAACgg	AA	TCTCAACTTgTTgTTTTACTgTT	34	CATgCTgCTTTgCCCggAgCCTTAg	TA	gAAgAgTCTTCCTTTACg
37	gAggAggggCagCAAACgg	AA	gACCTCCgTgCATCTTCTTATggg	38	ggTTCCAggAgTgACATgTCTggTg	TA	gAAgAgTCTTCCTTTACg
41	gAggAggggCagCAAACgg	AA	CggTgCTAgTAAgACATTAgTAAAT	42	AgCCAgTAGCagTgTCTgATgCAAT	TA	gAAgAgTCTTCCTTTACg
45	gAggAggggCagCAAACgg	AA	TCACAgCagTCCAAAggAgCagTTC	46	gCTCTTCTgAATgTTACAggCTTA	TA	gAAgAgTCTTCCTTTACg
49	gAggAggggCagCAAACgg	AA	gTTTTCCAAAgaATgCATCgACAA	50	TATgTACAAGACAAAgCagACTCT	TA	gAAgAgTCTTCCTTTACg
53	gAggAggggCagCAAACgg	AA	ggggAATAAAAgCAAAgAggCCAC	54	gACTAgCTACCAAACTgAgAgAgA	TA	gAAgAgTCTTCCTTTACg
57	gAggAggggCagCAAACgg	AA	TTTgCTCTAAgCACCATTAgACTC	58	gAgCagTgAATTgCATAATggTTTT	TA	gAAgAgTCTTCCTTTACg
61	gAggAggggCagCAAACgg	AA	ggAAgTgCTTAAACAggAAATTCAC	62	AgTAAAggAAAAACACTTgCCTTT	TA	gAAgAgTCTTCCTTTACg
65	gAggAggggCagCAAACgg	AA	AATTTgCTTTATTTTCTCCCA	66	AACAATCAAgTCAAAgTAACCATg	TA	gAAgAgTCTTCCTTTACg
69	gAggAggggCagCAAACgg	AA	CTAgACCAAAATgCTCTCCAAAAG	70	AgTTTTTATTgTTCTCTATTgTCgA	TA	gAAgAgTCTTCCTTTACg
73	gAggAggggCagCAAACgg	AA	TAAgAACAgCTTgCATTAAATCgTgg	74	TAgAATTTggTgATCggAgCgTTTT	TA	gAAgAgTCTTCCTTTACg
77	gAggAggggCagCAAACgg	AA	CTTgCCCTCagCATTgCagCATT	78	AATAgAAAgCCCCgATTATCACCC	TA	gAAgAgTCTTCCTTTACg

Organism: *G. gallus domesticus*Target mRNA: **SRY (sex determining region Y)-box 10 (Sox10)**Probe set: **20 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B3-Alexa546**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	gTCCCTgCCTCTATATCT	TT	gTCTTCCTCATCTAAGgCCAATA	2	CCATggACCCgTCACTCCATgTCTT	TT	CCACTCAACTTTAACCCg
3	gTCCCTgCCTCTATATCT	TT	TgTgAATCTTAGgCAGgACTgCTgC	4	TCCAgCAGggATCAAATTTCATgCA	TT	CCACTCAACTTTAACCCg
5	gTCCCTgCCTCTATATCT	TT	gCTgACAgtgCAgTTCCCTgAATCCT	6	ACgggCTATgAAATgAgAAAggCTA	TT	CCACTCAACTTTAACCCg
7	gTCCCTgCCTCTATATCT	TT	TgCTggAggAgCAAaggACCTggTCT	8	TTCACgTTTTCAgCAGACACAgTCA	TT	CCACTCAACTTTAACCCg
9	gTCCCTgCCTCTATATCT	TT	CAAgTgCATggTAgCTTTCTTggTg	10	AATATTggAACACATCTgggTgTT	TT	CCACTCAACTTTAACCCg
11	gTCCCTgCCTCTATATCT	TT	AgCTgTgAAAATCAgCAAaggAAgCA	12	gAggCggggAgAAAACTATAgCgT	TT	CCACTCAACTTTAACCCg
13	gTCCCTgCCTCTATATCT	TT	gggATTAAACAgATgggACAggggg	14	TTATAAAATCCATgCAggAAggggT	TT	CCACTCAACTTTAACCCg
15	gTCCCTgCCTCTATATCT	TT	AgCAGTgATgTACACCCCATCggCC	16	AgATggCgATAATgTgATgAACAAA	TT	CCACTCAACTTTAACCCg
17	gTCCCTgCCTCTATATCT	TT	TgAgTgAAAgTAggAgggAggTgCT	18	TATATCCACgAgAgTAICTTTCCAT	TT	CCACTCAACTTTAACCCg
19	gTCCCTgCCTCTATATCT	TT	CICTCTgATCAgTTgTCAggAgTCA	20	TACCCgTTAgAAggTCCCACAACAC	TT	CCACTCAACTTTAACCCg
21	gTCCCTgCCTCTATATCT	TT	gAACCCACTgggCTCATCTCCACCT	22	CCCggCgAgAggCAGTggTggTCTT	TT	CCACTCAACTTTAACCCg
23	gTCCCTgCCTCTATATCT	TT	TTCCgCCTCCgTggCTggTACTTgT	24	CCCTCgCCCTgCgTggCCTTgCCAT	TT	CCACTCAACTTTAACCCg
25	gTCCCTgCCTCTATATCT	TT	AgCCCCgCAgCCTCCCCTCCACC	26	ATTCTTgTAGTgAgCCTggATAgAg	TT	CCACTCAACTTTAACCCg
27	gTCCCTgCCTCTATATCT	TT	CCCAGgATgCCTgTggTCCAgTgg	28	gTgCCCATCggACATgggTgACCCT	TT	CCACTCAACTTTAACCCg
29	gTCCCTgCCTCTATATCT	TT	gCCgTggCTCTgACCTgAAgAgTgC	30	CTTAggggTggTgggAggAgTgggA	TT	CCACTCAACTTTAACCCg
31	gTCCCTgCCTCTATATCT	TT	gAgTCAgCTTTgCCTgCCTgCAgCT	32	TCCCCAagggAACgCCCTTCTCgCT	TT	CCACTCAACTTTAACCCg
33	gTCCCTgCCTCTATATCT	TT	ggTggCAGgTATTggTCAAATTCgT	34	TggCCTgggTggCCggCgTgTCCgT	TT	CCACTCAACTTTAACCCg
35	gTCCCTgCCTCTATATCT	TT	AgTgTCCACTggCCgCAGCCAaggC	36	CTCCATgCTgCTTggAgATCCAggC	TT	CCACTCAACTTTAACCCg
37	gTCCCTgCCTCTATATCT	TT	CCCAGTgCCCCCTgTTCTCCCTCC	38	AAACCTTgTgAAggCTgCAGCTCCT	TT	CCACTCAACTTTAACCCg
39	gTCCCTgCCTCTATATCT	TT	TggCTgACCgCTTACgATgCAGa	40	AgggTCCAgTCATAgCCgCTCAGCA	TT	CCACTCAACTTTAACCCg

Organism: *G. gallus domesticus*Target mRNA: **EPH receptor A4 (EphA4)**Probe set: **20 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B2-Alexa647**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	CCTCgTAAATCCTCATCA	AA	gCgCCgACggggACCCcgCCATgC	2	CAGACgCCgACgAggAgCgggAggA	AA	ATCATCCAgTAAACCgCC
5	CCTCgTAAATCCTCATCA	AA	CCCAGCTCCCCCTgCACCGAgCggg	6	CCTCCTCCAAGCgggCTCgCgATCC	AA	ATCATCCAgTAAACCgCC
9	CCTCgTAAATCCTCATCA	AA	TgACTgggCTCCATCACATTgCAAA	10	ATCCAATCAgTTCgTAACCAATTAT	AA	ATCATCCAgTAAACCgCC
13	CCTCgTAAATCCTCATCA	AA	TCCTTgTCgTTgTTTgATTCATAgT	14	gCAAACCTggCTCTCTCgAATAAAAC	AA	ATCATCCAgTAAACCgCC
17	CCTCgTAAATCCTCATCA	AA	AgTgggCACTTCTTATAgAAGACAC	18	ggAAACTgTgCCAggTTCgAACTg	AA	ATCATCCAgTAAACCgCC
21	CCTCgTAAATCCTCATCA	AA	ACgTCCTTCTCTCCgAgTgTgTgA	22	CCATCTgCCCCgCgTACATTTTTg	AA	ATCATCCAgTAAACCgCC
25	CCTCgTAAATCCTCATCA	AA	ATTTTgCAAAGCTTggCATTCACCAT	26	TCTgTTgAgAgCgCCTTgTAgTATC	AA	ATCATCCAgTAAACCgCC
29	CCTCgTAAATCCTCATCA	AA	AAgAAgCCCCgATCACAggTgCAgg	30	ATggATgCAGCATATTTCTgCTC	AA	ATCATCCAgTAAACCgCC
33	CCTCgTAAATCCTCATCA	AA	gCgCTCCACTCCAAGTTCAGTgACg	34	TCgTCCCgTCCCTTgTTCgTg	AA	ATCATCCAgTAAACCgCC
37	CCTCgTAAATCCTCATCA	AA	AAATgTACACCACTgCCACAggACC	38	gTTTTCAgCCGTTCTgCTgggggC	AA	ATCATCCAgTAAACCgCC
41	CCTCgTAAATCCTCATCA	AA	TgCTTggACACTCCATTCACTgCCC	42	gACACAgCTTggTCCTggCTggggT	AA	ATCATCCAgTAAACCgCC
45	CCTCgTAAATCCTCATCA	AA	gCAACgCTgTgCCTCgTTATCTCTT	46	ggCCTgTCAGgTCCAgCCAaggCCA	AA	ATCATCCAgTAAACCgCC
49	CCTCgTAAATCCTCATCA	AA	gCTgTCTTCACAATgCgATAgCTgC	50	AAACCTTTgATgTCAgTATTCCTgg	AA	ATCATCCAgTAAACCgCC
53	CCTCgTAAATCCTCATCA	AA	AACTCAAACggCCCACTgAAGTCTC	54	ATgggggAAggAACTgTgTAgTTg	AA	ATCATCCAgTAAACCgCC
57	CCTCgTAAATCCTCATCA	AA	gCTgCAATgAgAAgACCACgAgAA	58	TTgCTgCgCCTCTgCTgATgACAA	AA	ATCATCCAgTAAACCgCC
61	CCTCgTAAATCCTCATCA	AA	TATgTAAAAGgATCCACATATgTTC	62	TCCCTCACAgCTTgATTggATCCT	AA	ATCATCCAgTAAACCgCC
65	CCTCgTAAATCCTCATCA	AA	CATTTAgTAACAACgCCTTCCAAgT	66	TACTCAGTTATgATCATTACTggTT	AA	ATCATCCAgTAAACCgCC
69	CCTCgTAAATCCTCATCA	AA	TCCCgATgCACATAgCTCATgTCAG	70	TTgACCAgTATgTTTCgAgCAgCTA	AA	ATCATCCAgTAAACCgCC
73	CCTCgTAAATCCTCATCA	AA	CCAAATTTAggTCTgTCgCTgCgTT	74	AgTTTgTCCAgCATgTTgACAATCT	AA	ATCATCCAgTAAACCgCC
77	CCTCgTAAATCCTCATCA	AA	ggggAgCTgggATCCAAGCaggCTg	78	CTgACAAGAAACAACCgCCgAgAACT	AA	ATCATCCAgTAAACCgCC

S4.6 Split-initiator probes for Figures S13–S17

Organism: *G. gallus domesticus*

Target mRNA: **EPH receptor A4** (*EphA4*)

Probe set: **20 split-initiator probe pairs** (each probe carries half an HCR initiator)

HCR amplifier: **B1-Alexa546**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
3	gAggAggggCagCAAACgg	AA	TACACgCgggAgCCggTgACggCCC	4	TCCAgCAgggTCACTTCgTTggCgg	TA	gAAgAgTCTTCCTTTACg
7	gAggAggggCagCAAACgg	AA	TCATCCATTATgCTCACTTCCTCCC	8	TggTAGgTgCggATCggAgTgTTCT	TA	gAAgAgTCTTCCTTTACg
11	gAggAggggCagCAAACgg	AA	TATACCCTCTgAgCCCCCTCgCggg	12	TCTCTCAgCgTgAACTTgATTCAA	TA	gAAgAgTCTTCCTTTACg
15	gAggAggggCagCAAACgg	AA	ACCTCTgTATTCAGCTTCATgATCC	16	TTCTTgCTgAgAggCCCCACgTCCC	TA	gAAgAgTCTTCCTTTACg
19	gAggAggggCagCAAACgg	AA	gATgTATCAgCCCCAgTAATggTgT	20	CAggAgCCACgAACCTCCACCAGAg	TA	gAAgAgTCTTCCTTTACg
23	gAggAggggCagCAAACgg	AA	CAgTTgCCAATgggTACCAGCCATT	24	CgTTCTTCATAgCCAgCATTgCACA	TA	gAAgAgTCTTCCTTTACg
27	gAggAggggCagCAAACgg	AA	TgAggCgggCATTgTgCACATgCAA	28	gTAGAgCCTTCCCAgATgAgTAGC	TA	gAAgAgTCTTCCTTTACg
31	gAggAggggCagCAAACgg	AA	ggTgCggATggAgggCgAgTgCagg	32	TCgTTgACgTTggAAATCAggTTCT	TA	gAAgAgTCTTCCTTTACg
35	gAggAggggCagCAAACgg	AA	CgCTTgCACACCACgTTgTAggAgA	36	CAgTggCTgggCTCCCTgCCCCgC	TA	gAAgAgTCTTCCTTTACg
39	gAggAggggCagCAAACgg	AA	AggAggTCAgTgATggAAACCTTCg	40	ACCTCAAaggTgTAGTTggTgTgTg	TA	gAAgAgTCTTCCTTTACg
43	gAggAggggCagCAAACgg	AA	gCTgCTTggTTAgTTgTCACAgTgA	44	gCCTggATCAATgCAATTggggATg	TA	gAAgAgTCTTCCTTTACg
47	gAggAggggCagCAAACgg	AA	ACTTCATACTCCAaggATgACTCCAT	48	TCgTTTTggTCCTTTTCATAgTACT	TA	gAAgAgTCTTCCTTTACg
51	gAggAggggCagCAAACgg	AA	TgAAATACATATgAAgTCAgggggT	52	TATCCTgCTgCTgTCCTggCCCgCA	TA	gAAgAgTCTTCCTTTACg
55	gAggAggggCagCAAACgg	AA	ACTgTgggATTggTACCATCgCCAA	56	ACACTgCCAgCCACTgAAACAAGCA	TA	gAAgAgTCTTCCTTTACg
59	gAggAggggCagCAAACgg	AA	TCTgCCTCTTgCTTAgCTTTACTgT	60	ACACCTTggTTCAAATgTTTCTCCT	TA	gAAgAgTCTTCCTTTACg
63	gAggAggggCagCAAACgg	AA	TTCAgAgTCTTgATAgCCACACAgA	64	CTCCgTTgTTTgTCAgTgTAACCAg	TA	gAAgAgTCTTCCTTTACg
67	gAggAggggCagCAAACgg	AA	CgAAgCATCCCCACCAACTggATTA	68	AgATACTTCATTCCTgAgCCgATgC	TA	gAAgAgTCTTCCTTTACg
71	gAggAggggCagCAAACgg	AA	TggAgAgCAATggggCagTCCATTg	72	TTCTgCCAgCAgTCTAACATCAgCT	TA	gAAgAgTCTTCCTTTACg
75	gAggAggggCagCAAACgg	AA	CTCTTCaggCTgTTAgggTTgCggA	76	CTgggTCTggAgCTCTCgCTgCCTg	TA	gAAgAgTCTTCCTTTACg
79	gAggAggggCagCAAACgg	AA	TCCATTTAATggCTTggAgCCAgt	80	gCAgCTgTgAAgTTATCCTTgTATC	TA	gAAgAgTCTTCCTTTACg

Organism: *G. gallus domesticus*Target mRNA: early growth response 2 (*Egr2*)

Probe set: 20 split-initiator probe pairs (each probe carries half an HCR initiator)

HCR amplifier: B3-Alexa647

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	gTCCCTgCCTCTATATCT	TT	gCAGTgCTgggCTgCTTgCAGCTgT	2	CgTgCCAaggCTgTgCCAaggCTCA	TT	CCACTCAACTTTAACCg
3	gTCCCTgCCTCTATATCT	TT	gTCTggCgGTAACCTATTATggggg	4	CCCgCCgCAGCTCgCgCTggAggAC	TT	CCACTCAACTTTAACCg
5	gTCCCTgCCTCTATATCT	TT	CACgCCgCTCATCTgTgTgAACggC	6	gTCCACgCCgAgCATgCCgTCTgCg	TT	CCACTCAACTTTAACCg
7	gTCCCTgCCTCTATATCT	TT	AACCAACCCCAACCAgTgCgTACAA	8	CTCTTCggTCACCgTAAAgACAAAA	TT	CCACTCAACTTTAACCg
9	gTCCCTgCCTCTATATCT	TT	CCACCCCCCCCCCgCAGACgCAA	10	gTgCgCTCgTCCAACCgggCCCTT	TT	CCACTCAACTTTAACCg
11	gTCCCTgCCTCTATATCT	TT	ggTgTCAGCgTggAATAATTAAGg	12	CCCCggCAGgCACCTACggAAATA	TT	CCACTCAACTTTAACCg
13	gTCCCTgCCTCTATATCT	TT	TgCACTTgAgTAgCTgAgAgCCTgA	14	ggAggCCgCgAgCAGAgCCTTggCT	TT	CCACTCAACTTTAACCg
15	gTCCCTgCCTCTATATCT	TT	TTTTgCCTggCAGCCAAATggTgC	16	TgTgACAAgTgTggTAACgCgACTT	TT	CCACTCAACTTTAACCg
17	gTCCCTgCCTCTATATCT	TT	CgACCAgTCATTACTTTCCTCCgCA	18	CAGAAATAACgggATATCTCACC	TT	CCACTCAACTTTAACCg
19	gTCCCTgCCTCTATATCT	TT	ATATAAggACTgAggAACgggCCC	20	CCCTgAATgCCCGgACgTCACTgC	TT	CCACTCAACTTTAACCg
21	gTCCCTgCCTCTATATCT	TT	CgCgCCCTTgCgCTCCTTCTggCgC	22	TCCgggCTgCggggCggCggCggTC	TT	CCACTCAACTTTAACCg
23	gTCCCTgCCTCTATATCT	TT	ACCCCCACATgCAGCCgggTACgg	24	gCTgTCgTCgAgCgATCggTAAg	TT	CCACTCAACTTTAACCg
25	gTCCCTgCCTCTATATCT	TT	ggCAAACTTTCgTCTCgCgTCg	26	ACAACCATACTACAACCAGggAggg	TT	CCACTCAACTTTAACCg
27	gTCCCTgCCTCTATATCT	TT	CgCTAAgATgAggggAggCgAAAgC	28	TgTgggACCAggTggCAAAGCTgCC	TT	CCACTCAACTTTAACCg
29	gTCCCTgCCTCTATATCT	TT	CgCTgTCCCTTCgggAgCCTgggAA	30	CCATgTgCCACTCTCCggCAGAgCg	TT	CCACTCAACTTTAACCg
31	gTCCCTgCCTCTATATCT	TT	gAggTTgTgCTCCgCggCCgAgACA	32	TgCCgACTgAggACTCCACgACTCT	TT	CCACTCAACTTTAACCg
33	gTCCCTgCCTCTATATCT	TT	gTCgCgGCCAgCCgATgTgCgggg	34	AAAAAAAAAAAggAgAAAAAgCA	TT	CCACTCAACTTTAACCg
35	gTCCCTgCCTCTATATCT	TT	AgCCATggTTATCCAAggCTgTggC	36	TgATgCACgACgCTCCggCTgTgAC	TT	CCACTCAACTTTAACCg
37	gTCCCTgCCTCTATATCT	TT	CTACCACgACggCACCgCATgCAT	38	TCACACCACAAggCACCAGgAC	TT	CCACTCAACTTTAACCg
39	gTCCCTgCCTCTATATCT	TT	gAgCgCCgACCCCTgCCgTCgAAggg	40	TTATTCgTCgTCgCTTATAAACgCC	TT	CCACTCAACTTTAACCg

S4.7 Split-initiator probes Figures 5, S18, and S19

Organism: *G. gallus domesticus*Target mRNA: **diencephalon/mesencephalon homeobox 1 (*Dmbx1*)**Probe set: **20 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B1-Alexa546**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	gAggAggggCagCAAACgg	AA	CgCTCAgTgAgTTCATggCATgCAG	2	CTgCCTgCTggTgCAggTTgTACAT	TA	gAAgAgTCTTCCTTTACg
5	gAggAggggCagCAAACgg	AA	AAATgATgTCAgCCAgTCTCTCCgC	6	ggTgCTgggATCCATAACgTgCCTC	TA	gAAgAgTCTTCCTTTACg
9	gAggAggggCagCAAACgg	AA	AAATgATgTCAgCCAgTCTCTCCgC	10	gCTTCTgCAgCTgCTCCTTCTggAg	TA	gAAgAgTCTTCCTTTACg
13	gAggAggggCagCAAACgg	AA	CAGTgTCAggTATTgTggACTgggT	14	CCTCACTTgggATgCTCTgAgTCTT	TA	gAAgAgTCTTCCTTTACg
17	gAggAggggCagCAAACgg	AA	CCTCCTCTCTgTCAgTCTggTCCTC	18	TAgCCTCATCCAAggTgCTCTTAAA	TA	gAAgAgTCTTCCTTTACg
21	gAggAggggCagCAAACgg	AA	CgCTgATgggAgACTCTgATTTTgg	22	CACTgCTAgATgAAggAgTCACggT	TA	gAAgAgTCTTCCTTTACg
25	gAggAggggCagCAAACgg	AA	CTgCCATgTgCTggCggAATTgCTC	26	AggAgTAAAggACCAAgTTgTTggT	TA	gAAgAgTCTTCCTTTACg
29	gAggAggggCagCAAACgg	AA	gACAgTgCAgggAgCTCAgAggCgC	30	CgTgggAgAgAgATTggTAgTACgA	TA	gAAgAgTCTTCCTTTACg
33	gAggAggggCagCAAACgg	AA	TCTCAACTTgTTgTTTTACTgTT	34	CATgCTgCTTTgCCCggAgCCTTAg	TA	gAAgAgTCTTCCTTTACg
37	gAggAggggCagCAAACgg	AA	gACCTCCggTgCATCTTCTTATggg	38	ggTTCCAggAgTgACATgTCTggTg	TA	gAAgAgTCTTCCTTTACg
41	gAggAggggCagCAAACgg	AA	CggTgCTAgTAAgACATTAgTAAAT	42	AgCCAgtAgCagTgTCTgATgCAAT	TA	gAAgAgTCTTCCTTTACg
45	gAggAggggCagCAAACgg	AA	TCACAgCagTCCAAAggAgACAgtTC	46	gCTCTTCTgAATgTTACAggCTTA	TA	gAAgAgTCTTCCTTTACg
49	gAggAggggCagCAAACgg	AA	gTTTTCCCAAAGAAAgCATCgACAA	50	TATgTACAAGACAAAgCAGgACTCT	TA	gAAgAgTCTTCCTTTACg
53	gAggAggggCagCAAACgg	AA	ggggAATAAAAgCAAAAgAggCCAC	54	gACTAgCTACCAAAACTgAgAgAgA	TA	gAAgAgTCTTCCTTTACg
57	gAggAggggCagCAAACgg	AA	TTTgCTCTAAgCACATTAAgACTC	58	gAgCagTgAATgCATAATgTTTT	TA	gAAgAgTCTTCCTTTACg
61	gAggAggggCagCAAACgg	AA	ggAAgTgCTTAAACAggAAATTCAC	62	AgTAAAggAAAAACACTTgCCTTT	TA	gAAgAgTCTTCCTTTACg
65	gAggAggggCagCAAACgg	AA	AATTTggCTTTCATTTTCTCCCCA	66	AACAATCAAgTCAAAAgTAACCATg	TA	gAAgAgTCTTCCTTTACg
69	gAggAggggCagCAAACgg	AA	CTAgACCAAAATgCTCTCCAAAAAg	70	AgTTTTTATTgTTCTCTATTgTCgA	TA	gAAgAgTCTTCCTTTACg
73	gAggAggggCagCAAACgg	AA	TAAgAACAgCTTgCATTAATCgTgg	74	TAgAATTTggTgATCggAgCgTTTT	TA	gAAgAgTCTTCCTTTACg
77	gAggAggggCagCAAACgg	AA	CTTggCCTCCAgCATTgCagCATT	78	AATAgAAAgCCCCgATTATCACCC	TA	gAAgAgTCTTCCTTTACg

Organism: *G. gallus domesticus*Target mRNA: **diencephalon/mesencephalon homeobox 1 (*Dmbx1*)**Probe set: **20 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B2-Alexa647**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
3	CCTCgTAAATCCTCATCA	AA	AgTCgggCgCgTgCTgTgCTTgCTg	4	gggTgAgggCgTgCACTgAgggCCg	AA	ATCATCCAgTAAACCgCC
7	CCTCgTAAATCCTCATCA	AA	AggCCgTgCgACTTCgCCTTgTTT	8	CCAgTgCCTCCAgCTgTTgggCAgT	AA	ATCATCCAgTAAACCgCC
11	CCTCgTAAATCCTCATCA	AA	CACTgTggCTgCTTTCACAgTCCTT	12	CCAgCACAggTggTTCcgTCTTCCC	AA	ATCATCCAgTAAACCgCC
15	CCTCgTAAATCCTCATCA	AA	ACAgggTTAggTTgAggTCTgTgCT	16	gCgCTgATTCACTggCTgACTgCTC	AA	ATCATCCAgTAAACCgCC
19	CCTCgTAAATCCTCATCA	AA	CgTCCACACCAggACTCTTgTCCAC	20	TCgCTCTCTTgCAgTTCAAAGCCTT	AA	ATCATCCAgTAAACCgCC
23	CCTCgTAAATCCTCATCA	AA	AggAgTAggAgTgAgTTTgAgCCA	24	gCAggCggAAgAggCTCAggggTgA	AA	ATCATCCAgTAAACCgCC
27	CCTCgTAAATCCTCATCA	AA	TggCAGACggTgTCCCCATTTcGAA	28	TgTTgACATTCATgCCCAAgTAggg	AA	ATCATCCAgTAAACCgCC
31	CCTCgTAAATCCTCATCA	AA	ggATgggTgAggACCAgACCTgCTg	32	ggCTTggAAgAgAgCTggAggCCTg	AA	ATCATCCAgTAAACCgCC
35	CCTCgTAAATCCTCATCA	AA	gTAgggTgTCAAgCCCAAgggATgC	36	ACggTTgCCCATCTggCAgTCAgTT	AA	ATCATCCAgTAAACCgCC
39	CCTCgTAAATCCTCATCA	AA	ACTCCAggAAgAgATgAgggTggAA	40	AAAgtTTTCCCTgATAgggAgCACC	AA	ATCATCCAgTAAACCgCC
43	CCTCgTAAATCCTCATCA	AA	TCCTCCTCAAATATTTAAAgAAgAC	44	CTgTCTAAACACACATCCTCTCCCT	AA	ATCATCCAgTAAACCgCC
47	CCTCgTAAATCCTCATCA	AA	ACATTATCgCAgggATgAggTgAgg	48	AAAAAgggTgTATATAACACggTTg	AA	ATCATCCAgTAAACCgCC
51	CCTCgTAAATCCTCATCA	AA	gCggTggATgCTTCAACATTgTAA	52	TTCTgTAACACTgACAgTAACACAC	AA	ATCATCCAgTAAACCgCC
55	CCTCgTAAATCCTCATCA	AA	gggAgCgTggCTgATTTgTgACTTT	56	AACCCAAgAAgAgCAACTAgCTgTg	AA	ATCATCCAgTAAACCgCC
59	CCTCgTAAATCCTCATCA	AA	TCAgCTTTAgCAgAAgAgAgAAg	60	TTgCATCATTTCTgCCgTTATAA	AA	ATCATCCAgTAAACCgCC
63	CCTCgTAAATCCTCATCA	AA	TCTgTCTgTgAACAAgTgCTATTAg	64	CAgCAgCATTTggCCAgCATTTTgT	AA	ATCATCCAgTAAACCgCC
67	CCTCgTAAATCCTCATCA	AA	TTACACTTCACTgAAgACCAAAGg	68	AACCCATAATTTgTAAATgggggA	AA	ATCATCCAgTAAACCgCC
71	CCTCgTAAATCCTCATCA	AA	gTATgAACACAgTgggAgTTCATAC	72	TTgTCAAaggAACCATATAATTCATg	AA	ATCATCCAgTAAACCgCC
75	CCTCgTAAATCCTCATCA	AA	AgggTCTgAAgCTgCACAgCTTgAg	76	CACTTgTTACATTCTCACTTgCTAA	AA	ATCATCCAgTAAACCgCC
79	CCTCgTAAATCCTCATCA	AA	CAAgCCAATCTACTCCTCgCTgCAg	80	ggTTgCTTggggACATggtACTTTT	AA	ATCATCCAgTAAACCgCC

Organism: *G. gallus domesticus*Target mRNA: **EPH receptor A4 (*EphA4*)**Probe set: **20 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B1-Alexa546**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
3	gAggAggggCAgCAAACgg	AA	TACACgCgggAgCCggTgACggCCC	4	TCCAgCAgggTCACTTCgTTggCgg	TA	gAAgAgTCTTCCTTTACg
7	gAggAggggCAgCAAACgg	AA	TCATCCATTATgCTCACTTCCTCCC	8	TggTAggTgCggATCggAgTgTTCT	TA	gAAgAgTCTTCCTTTACg
11	gAggAggggCAgCAAACgg	AA	TATACCTCTgAgCCCCCTCgCggg	12	TCTCTCAgCgTgAACTTgATTTCAA	TA	gAAgAgTCTTCCTTTACg
15	gAggAggggCAgCAAACgg	AA	ACCTCTgTATTCAgCTTCATgATCC	16	TTCTTgCTgAgAggCCCCACgTCCC	TA	gAAgAgTCTTCCTTTACg
19	gAggAggggCAgCAAACgg	AA	gATgTATCAgCCCCAgTAATgTgT	20	CAggAgCCACgAACCTCCACCAgAg	TA	gAAgAgTCTTCCTTTACg
23	gAggAggggCAgCAAACgg	AA	CAGTTgCCAATgggTACCAGCATT	24	CgTTCTTCATAgCCAATgCACA	TA	gAAgAgTCTTCCTTTACg
27	gAggAggggCAgCAAACgg	AA	TgAggCgggCATTgTgCACATgCAA	28	gTAGAgCCTTCCCAgATggAgTAGC	TA	gAAgAgTCTTCCTTTACg
31	gAggAggggCAgCAAACgg	AA	ggTgCggATgAgggCgAgTgCAgg	32	TCgTTgACgTTggAAATCAggTTCT	TA	gAAgAgTCTTCCTTTACg
35	gAggAggggCAgCAAACgg	AA	CgCTTgCACACCACgTTgTAggAgA	36	CAgTggCTgggCTCCCCTgCCCCgC	TA	gAAgAgTCTTCCTTTACg
39	gAggAggggCAgCAAACgg	AA	AggAggTCAGTgATggAAACCTTCg	40	ACCTCAAaggTgTAGTTggTgTgTg	TA	gAAgAgTCTTCCTTTACg
43	gAggAggggCAgCAAACgg	AA	gCTgCTTggTTAgTTgTCACAgTgA	44	gCCTggATCAATgCAATTggggATg	TA	gAAgAgTCTTCCTTTACg
47	gAggAggggCAgCAAACgg	AA	ACTTCATACTCCAgATgACTCCAT	48	TCgTTTTggTCCTTTTCATAgTACT	TA	gAAgAgTCTTCCTTTACg
51	gAggAggggCAgCAAACgg	AA	TgAAATACATATgAAgTCAgggggT	52	TATCCTgCTgCTgTCCTggCCgCA	TA	gAAgAgTCTTCCTTTACg
55	gAggAggggCAgCAAACgg	AA	ACTgTgggATTggTACCATCgCCAA	56	ACACTgCCAgCCACTgAAACAAgCA	TA	gAAgAgTCTTCCTTTACg
59	gAggAggggCAgCAAACgg	AA	TCTgCCTCTTgCTTAgCTTTACTgT	60	ACACCTTggTTCAAATgTTTCTCCT	TA	gAAgAgTCTTCCTTTACg
63	gAggAggggCAgCAAACgg	AA	TTCAgAgTCTTgATAgCCACACAgA	64	CTCCgTTgTTTgTCAgTgTAACCAg	TA	gAAgAgTCTTCCTTTACg
67	gAggAggggCAgCAAACgg	AA	CgAAgCATCCCCACCAACTggATTA	68	AgATACTTCATTCCTgAgCCgATgC	TA	gAAgAgTCTTCCTTTACg
71	gAggAggggCAgCAAACgg	AA	TggAgAgCAATggggCAgTCCATTg	72	TTCTgCCAgCAGTCTAACATCAgCT	TA	gAAgAgTCTTCCTTTACg
75	gAggAggggCAgCAAACgg	AA	CTCTTCAggCTgTTAgggTTgCggA	76	CTgggTCTggAgCTCTCgCTgCCTg	TA	gAAgAgTCTTCCTTTACg
79	gAggAggggCAgCAAACgg	AA	TCCATTTAATggCTTggAgCCAgt	80	gCAGCTgTgAAgTTATCCTTgTATC	TA	gAAgAgTCTTCCTTTACg

Organism: *G. gallus domesticus*Target mRNA: **EPH receptor A4 (*EphA4*)**Probe set: **20 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B2-Alexa647**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	CCTCgTAAATCCTCATCA	AA	gCgCCgACggggACCCcgCCATgC	2	CAGACgCCgACgAggAgCgggAggA	AA	ATCATCCAgTAAACCgCC
5	CCTCgTAAATCCTCATCA	AA	CCCAGCTCCCCCTgCACGgAgCggg	6	CCTCCTCCAAGCgggCTCgCgATCC	AA	ATCATCCAgTAAACCgCC
9	CCTCgTAAATCCTCATCA	AA	TgACTgggCTCCATCACATTgCAAA	10	ATCCAATCAgTTCgTAACCAATTAT	AA	ATCATCCAgTAAACCgCC
13	CCTCgTAAATCCTCATCA	AA	TCCTTgTCgTTgTTTgATTCATAgT	14	gCAAACtggCTCTCTCgAATAAAAC	AA	ATCATCCAgTAAACCgCC
17	CCTCgTAAATCCTCATCA	AA	AgTgggCACTTCTTATAgAAGACAC	18	ggAAACTgTgCCAggTTCgAACTg	AA	ATCATCCAgTAAACCgCC
21	CCTCgTAAATCCTCATCA	AA	ACgTCCTTCTCTCCgAgTgTgTgA	22	CCATCTgCCCCgCgTACATTTTTg	AA	ATCATCCAgTAAACCgCC
25	CCTCgTAAATCCTCATCA	AA	ATTTTgCAAAGCTTggCATTCACCAT	26	TCTgTTgAgAgCgCCTTgTAgTATC	AA	ATCATCCAgTAAACCgCC
29	CCTCgTAAATCCTCATCA	AA	AAgAAgCCCCgATCACAggTgCAgg	30	ATggATgCAGCATATTTCTgCTC	AA	ATCATCCAgTAAACCgCC
33	CCTCgTAAATCCTCATCA	AA	gCgCTCCACTCCAAGTTCAGTgACg	34	TCgTCCCgTCCCTTgTTCgTg	AA	ATCATCCAgTAAACCgCC
37	CCTCgTAAATCCTCATCA	AA	AAATgTACACCACTgCCACAggACC	38	gTTTTCAgCCGTTCTgCTgggggC	AA	ATCATCCAgTAAACCgCC
41	CCTCgTAAATCCTCATCA	AA	TgCTTggACACTCCATTCACTgCCC	42	gACACAgCTTggTCCTggCTggggT	AA	ATCATCCAgTAAACCgCC
45	CCTCgTAAATCCTCATCA	AA	gCAACgCTgTgCCTCgTTATCTCTT	46	ggCCTgTCAGgTCCAgCCAaggCCA	AA	ATCATCCAgTAAACCgCC
49	CCTCgTAAATCCTCATCA	AA	gCTgTCTTCACAATgCgATAgCTgC	50	AAACCTTTgATgTCAgTATTCCTgg	AA	ATCATCCAgTAAACCgCC
53	CCTCgTAAATCCTCATCA	AA	AACTCAAACggCCCACTgAAGTCTC	54	ATgggggAAggAACTgTgTTAgTTg	AA	ATCATCCAgTAAACCgCC
57	CCTCgTAAATCCTCATCA	AA	gCTgCAATgAgAATgACCACgAgAA	58	TTgCTgCgCCTCTgCTgATgACAA	AA	ATCATCCAgTAAACCgCC
61	CCTCgTAAATCCTCATCA	AA	TATgTAAAAGgATCCACATATgTTC	62	TCCCTCACAgCTTgATTggATCCT	AA	ATCATCCAgTAAACCgCC
65	CCTCgTAAATCCTCATCA	AA	CATTTAgTAACAACgCCTTCCAAgT	66	TACTCAGTTATgATCATTACTggTT	AA	ATCATCCAgTAAACCgCC
69	CCTCgTAAATCCTCATCA	AA	TCCCgATgCACATAgCTCATgTCAG	70	TTgACCAgTATgTTTCgAgCAgCTA	AA	ATCATCCAgTAAACCgCC
73	CCTCgTAAATCCTCATCA	AA	CCAAATTTAggTCTgTCgCTgCgTT	74	AgTTTgTCCAgCATgTTgACAATCT	AA	ATCATCCAgTAAACCgCC
77	CCTCgTAAATCCTCATCA	AA	ggggAgCTgggATCCAAGCAGggCTg	78	CTgACAAGAAACAACCgCCgAgAACT	AA	ATCATCCAgTAAACCgCC

S4.8 Split-initiator probes for Figure 6, S20–S28

Organism: *H. sapiens sapiens*

Target mRNA: destabilized enhanced green fluorescent protein (*Tg(d2eGFP)*)

Probe set: 12 split-initiator probe pairs (each probe carries half an HCR initiator)

HCR amplifier: B3-Alexa594 (Figures 6A and S20)

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	gTCCCTgCCTCTATATCT	TT	TTgTggCCgTTTACgTCgCCgTCCA	2	CCCTCgCCCTCgCCggACACgCTgA	TT	CCACTCAACTTTAACCCg
3	gTCCCTgCCTCTATATCT	TT	AgggTCAGCTTgCCgTAggTggCAT	4	AgCTTgCCggTggTgCAGATgAACT	TT	CCACTCAACTTTAACCCg
5	gTCCCTgCCTCTATATCT	TT	gTCACgAgggTgggCCAgggCACgg	6	AAgCACTgCACgCCgTAggTCAggg	TT	CCACTCAACTTTAACCCg
7	gTCCCTgCCTCTATATCT	TT	TgCTTCATgTggTCggggTAgCggC	8	ggCATggCggACTTgAAgAAgTCgT	TT	CCACTCAACTTTAACCCg
9	gTCCCTgCCTCTATATCT	TT	ATggTgCgCTCCTggACgTAgCCTT	10	TTgTAGTTgCCgTCgTCCTTgAAgA	TT	CCACTCAACTTTAACCCg
11	gTCCCTgCCTCTATATCT	TT	CCCTCgAACTTCACCTCggCgCggg	12	AgCTCgATgCggTTCACCAgggTgT	TT	CCACTCAACTTTAACCCg
13	gTCCCTgCCTCTATATCT	TT	CCgTCCTCCTTgAAgTCgATgCCCT	14	TACTCCAgCTTgTgCCCCAggATgT	TT	CCACTCAACTTTAACCCg
15	gTCCCTgCCTCTATATCT	TT	ATATAgACgTTgTggCTgTTgTAgT	16	ATgCCgTTCTTCTgCTTgTCggCCA	TT	CCACTCAACTTTAACCCg
17	gTCCCTgCCTCTATATCT	TT	TTgTggCggATCTTgAAgTTCACCT	18	gCgAgCTgCACgCTgCCgTCCTCgA	TT	CCACTCAACTTTAACCCg
19	gTCCCTgCCTCTATATCT	TT	ATgggggTgTTCTgCTggTAgTggT	20	TCgggCAgCAgCACggggCCgTCgC	TT	CCACTCAACTTTAACCCg
21	gTCCCTgCCTCTATATCT	TT	gCggACTgggTgCTCAggTAgTggT	22	CgCTTCTCgTTggggTCTTTgCTCA	TT	CCACTCAACTTTAACCCg
23	gTCCCTgCCTCTATATCT	TT	ACgAACTCCAgCAggACCATgTgAT	24	ATgCCgAgAgTgATCCcgCggCgg	TT	CCACTCAACTTTAACCCg

Organism: *H. sapiens sapiens*Target mRNA: **glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*)**Probe set: **10 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B5-Alexa488 (Figures 6B, S24, and S23)**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
3	CTCACTCCCAATCTCTAT	AA	gggTCATTgATggCAACAATATCCA	4	TAAACCATgTAgTTgAggTCAATgA	AA	CTACCCTACAAATCCAAT
7	CTCACTCCCAATCTCTAT	AA	TTTCATTgATgACAAgCTTCCCgT	8	TCTCgCTCCTggAAgATggTgATgg	AA	CTACCCTACAAATCCAAT
11	CTCACTCCCAATCTCTAT	AA	gCCTTCTCCATggTggTgAAgACgC	12	TTggCTCCCCCTgCAAATgAgCCC	AA	CTACCCTACAAATCCAAT
15	CTCACTCCCAATCTCTAT	AA	AggCTgTTgTCATACTTCTCATggT	16	gTgCaggAggCATTgCTgATgATCT	AA	CTACCCTACAAATCCAAT
19	CTCACTCCCAATCTCTAT	AA	gCATggACTgTggTCATgAgTCCCT	20	TCCACAgTCTTCTgggTggCagTgA	AA	CTACCCTACAAATCCAAT
23	CTCACTCCCAATCTCTAT	AA	gCCTTggCagCgCCAgTAgAggCag	24	TTCagCTCaggATgACCTTgCCCA	AA	CTACCCTACAAATCCAAT
27	CTCACTCCCAATCTCTAT	AA	ggTTTTTCTAgACggCAggTCAggT	28	ACCTTCTTgATgTCATCATATTTgg	AA	CTACCCTACAAATCCAAT
31	CTCACTCCCAATCTCTAT	AA	CTgTTgAAgTCagAggAgACCACCT	32	gCgTCAAAGgTggAggAgTgggTgT	AA	CTACCCTACAAATCCAAT
35	CTCACTCCCAATCTCTAT	AA	CTggTggTCCAggggTCTTACTCCT	36	TCTCTTCTCTTgTgCTCTTgCTgg	AA	CTACCCTACAAATCCAAT
39	CTCACTCCCAATCTCTAT	AA	CTACATggCAACTgTgAggAggggA	40	CCTAggCCCCCTCCCTCTTCAAagg	AA	CTACCCTACAAATCCAAT

Organism: *H. sapiens sapiens*Target mRNA: **glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*)**Probe set: **10 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B4-Alexa594 (Figures 6B and S24), B4-Alexa488 (Figures 6C and S27)**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	CCTCAACCTACCTCCAAC	AA	ACCagCgCCCAATACgACCAAATC	2	TTACCagAgTTAAAAGCagCCCTgg	AT	TCTCACCATATTCgCTTC
5	CCTCAACCTACCTCCAAC	AA	CCATgggTggAATCATATTTggAACA	6	TCAgCCTTgACggTgCCATggAATT	AT	TCTCACCATATTCgCTTC
9	CCTCAACCTACCTCCAAC	AA	gCATCgCCCCACTTgATTTTggAgg	10	gTggACTCCACgACgTACTCagCgC	AT	TCTCACCATATTCgCTTC
13	CCTCAACCTACCTCCAAC	AA	gCagAgggggCagAgATgATgACCC	14	ACACCCATgACgAACATgggggCAT	AT	TCTCACCATATTCgCTTC
17	CCTCAACCTACCTCCAAC	AA	TTggCCAggggTgCTAAgCagTTgg	18	ACgATACCAAAGTTgTCATggATgA	AT	TCTCACCATATTCgCTTC
21	CCTCAACCTACCTCCAAC	AA	TCACgCCACAgTTCCCGgAggggC	22	ATgATgTTCTgAgAgCCCCgCggC	AT	TCTCACCATATTCgCTTC
25	CCTCAACCTACCTCCAAC	AA	CggAAggCCATgCCAgTgAgCTTCC	26	ACCACtGACAGtTgTgCagTggggA	AT	TCTCACCATATTCgCTTC
29	CCTCAACCTACCTCCAAC	AA	AgggggCCCTCCgACgCTTgCTTCA	30	TgCTCagTgTAgCCCAggATgCCCT	AT	TCTCACCATATTCgCTTC
33	CCTCAACCTACCTCCAAC	AA	TggTCgTTgAgggCAATgCCAgCCC	34	TCATACCagggAAATgAgCTTgACAA	AT	TCTCACCATATTCgCTTC
37	CCTCAACCTACCTCCAAC	AA	CAgggACTCCCCAgCagTgAgggTC	38	TTCagTgTggTgggggACTgAgTgT	AT	TCTCACCATATTCgCTTC

Organism: *H. sapiens sapiens*Target mRNA: **actin beta (ACTB)**Probe set: **10 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B2-Alexa594 (Figure S23)**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	CCTCgTAAATCCTCATCA	AA	gCggggCggACgCggTCTCggCggT	2	gATCggCAAAGgCgAggCTCTgTgC	AA	ATCATCCAgTAAACCgCC
5	CCTCgTAAATCCTCATCA	AA	gCTggCggCgggTgTggACgggCgg	6	gCgCggCgATATCATCATCCATggT	AA	ATCATCCAgTAAACCgCC
9	CCTCgTAAATCCTCATCA	AA	TgCgCAAgTTAggTTTTgTCAAgAA	10	AAgCCATgCCAATCTCATCTTgTTT	AA	ATCATCCAgTAAACCgCC
13	CCTCgTAAATCCTCATCA	AA	ACCAAAACAAAACAAAAAACAAA	14	CTgAgTCAAgCCAAAAAAAAAAAAA	AA	ATCATCCAgTAAACCgCC
17	CCTCgTAAATCCTCATCA	AA	CACCTTCACCgTCCAgtTTTTAAA	18	gggATgCTCgTCCAACCgACTgCT	AA	ATCATCCAgTAAACCgCC
21	CCTCgTAAATCCTCATCA	AA	AgTCCTCggCCACATTgTgAACTTT	22	ATTAAAAAACAACAATgTgCAATC	AA	ATCATCCAgTAAACCgCC
25	CCTCgTAAATCCTCATCA	AA	ACAACgCATCTCATATTTggAATgA	26	TTTTAggATggCAAgggACTTCCTg	AA	ATCATCCAgTAAACCgCC
29	CCTCgTAAATCCTCATCA	AA	ATTCTCCTTAgAgAgAAgTggggTg	30	TgTgTggACTTgggAgAggACTggg	AA	ATCATCCAgTAAACCgCC
33	CCTCgTAAATCCTCATCA	AA	ACACgAAAgCAATgCTATCACCTCC	34	TTAAAAAATTTTgCATTACATAAT	AA	ATCATCCAgTAAACCgCC
37	CCTCgTAAATCCTCATCA	AA	CAAATAAAAAAgTATTAaggCgAA	38	CACgAAggCTCATCATTCAAAATAA	AA	ATCATCCAgTAAACCgCC

Organism: *H. sapiens sapiens*Target mRNA: phosphoglycerate kinase 1 (*PGK1*)

Probe set: 18 split-initiator probe pairs (each probe carries half an HCR initiator)

HCR amplifier: B1-Alexa488 (Figures 6B and S25)

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	gAggAggggCAgCAAACgg	AA	CCgCCCCCTCCCgGCCgCTgCTCTC	2	CTACCgCCCCACACCCgCCTCCCg	TA	gAAgAgTCTTCCTTTACg
5	gAggAggggCAgCAAACgg	AA	CAACgAgggAgCCgACTgCCgACgT	6	gCTggggAgAgAggTCggTgATTCg	TA	gAAgAgTCTTCCTTTACg
9	gAggAggggCAgCAAACgg	AA	gACTCTCATAACgACCCgCTTCCCT	10	gTTgTTCTTCATAggAACATTgAAg	TA	gAAgAgTCTTCCTTTACg
13	gAggAggggCAgCAAACgg	AA	CTTCCCTTCTTCTCCACATgAAAg	14	AACCTTgTTCCCAgAAgCATCTTTT	TA	gAAgAgTCTTCCTTTACg
17	gAggAggggCAgCAAACgg	AA	gCCAAAAGCATCATTgACATgACA	18	CATggAgCTgTgggCTCTgTgAgCA	TA	gAAgAgTCTTCCTTTACg
21	gAggAggggCAgCAAACgg	AA	gCTCTCCAAGgCCTTTgCAAAGTAG	22	CAggATggCCAggAAgggTCgCTCT	TA	gAAgAgTCTTCCTTTACg
25	gAggAggggCAgCAAACgg	AA	CTTCTCAgCTTTgACATTAaggTCT	26	AACAggCAAggTAATCTTCACCCA	TA	gAAgAgTCTTCCTTTACg
29	gAggAggggCAgCAAACgg	AA	CCAgCCAgCAggTATgCCAgAAgCC	30	gCTTTCAggACCACAgTCCAAGCCC	TA	gAAgAgTCTTCCTTTACg
33	gAggAggggCAgCAAACgg	AA	AgCTTCCATTCAAATACCCCCACA	34	CATgAgAgCTTTggTTCCCCgggCA	TA	gAAgAgTCTTCCTTTACg
37	gAggAggggCAgCAAACgg	AA	gTACTAAATATTgCTgAgAgCATCC	38	TgTgCACAggAACTAAAAGgCAggA	TA	gAAgAgTCTTCCTTTACg
41	gAggAggggCAgCAAACgg	AA	ggCCACTAgCTgAATCTTgACATgg	42	TTAAGggTTCTTggCACTgCATCTC	TA	gAAgAgTCTTCCTTTACg
45	gAggAggggCAgCAAACgg	AA	CTAAAAAATTCAAATgggATCTTgA	46	ACTCTAgAATgCACAAATggTTTAgT	TA	gAAgAgTCTTCCTTTACg
49	gAggAggggCAgCAAACgg	AA	TAATCATAATAACCTACATCAAAAg	50	TgCTgAgTAGTgAAACAgTgACAAA	TA	gAAgAgTCTTCCTTTACg
53	gAggAggggCAgCAAACgg	AA	TCAATggACACTTTTATTgTTTACT	54	gACAggAAAAAAAAAAAAATCACgg	TA	gAAgAgTCTTCCTTTACg
57	gAggAggggCAgCAAACgg	AA	CTgCCCCACTTCTTgCATTCAgCAA	58	TCTAATTgTCCCATCTCTCCACTgC	TA	gAAgAgTCTTCCTTTACg
61	gAggAggggCAgCAAACgg	AA	CTgATAAAAAATAAAgTTAgAATAA	62	gACTTTTTAAATTATgATCATgTgT	TA	gAAgAgTCTTCCTTTACg
65	gAggAggggCAgCAAACgg	AA	CAAgAgTTgAAAgTggTCACCTCTg	66	AACATggAggTATATACCTgAAAAA	TA	gAAgAgTCTTCCTTTACg
69	gAggAggggCAgCAAACgg	AA	gAgCCTTCCTCCATggTATgAAATA	70	TgAAgAAgTggAAATATATgTggAA	TA	gAAgAgTCTTCCTTTACg

Organism: *H. sapiens sapiens*Target mRNA: phosphoglycerate kinase 1 (*PGK1*)

Probe set: 18 split-initiator probe pairs (each probe carries half an HCR initiator)

HCR amplifier: B2-Alexa594 (Figures 6B, 6C, S25, and S27)

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
3	CCTCgTAAATCCTCATCA	AA	ACCgCgCgggCAgAACAgggCCCA	4	gCTCCgAggCTTgCAgAATgCggA	AA	ATCATCCAgTAAACCgCC
7	CCTCgTAAATCCTCATCA	AA	gTTAgAAAgCgACATTTTggAAATA	8	AACgTCCAgCTTgTCCAgCgTCAGC	AA	ATCATCCAgTAAACCgCC
11	CCTCgTAAATCCTCATCA	AA	CTTAATCCTCTggTTgTTgTTATC	12	gCagAATTTgATgCTTgggACAgCA	AA	ATCATCCAgTAAACCgCC
15	CCTCgTAAATCCTCATCA	AA	ggAgTACTTgTCAggCATgggCACA	16	TTTgAgTTCTACAgCAACTggCTCT	AA	ATCATCCAgTAAACCgCC
19	CCTCgTAAATCCTCATCA	AA	AgCCTTCTgTggCAgATTgACTCCT	20	CAgCTCCTTCTTCATCAAAAACCCA	AA	ATCATCCAgTAAACCgCC
23	CCTCgTAAATCCTCATCA	AA	ATCAAACCTgTCAGCAgTgACAAAg	24	AgTggCTTggCCAgtCTTggCATTC	AA	ATCATCCAgTAAACCgCC
27	CCTCgTAAATCCTCATCA	AA	AgTgACAgCCTCAGCATACTTCTTg	28	ACCATTCCACACAATCTgCTTAGCC	AA	ATCATCCAgTAAACCgCC
31	CCTCgTAAATCCTCATCA	AA	CCTAgAAgTggCTTTCACCACCTCA	32	TCCACCACCTATgATggTgATgCAg	AA	ATCATCCAgTAAACCgCC
35	CCTCgTAAATCCTCATCA	AA	CTCCAAACTggCACCACCCCAgTg	36	CCCAGgAAgACTTTACCTTCCAgg	AA	ATCATCCAgTAAACCgCC
39	CCTCgTAAATCCTCATCA	AA	CAGAAAAgCTAAgTTgACTTAGgg	40	ggTTTTAgCTAATgCCAAGTggAgA	AA	ATCATCCAgTAAACCgCC
43	CCTCgTAAATCCTCATCA	AA	AgATgAgCTgAgATgCTgTgCAACT	44	AATgTATgCAAAATCCAaggTgCAgT	AA	ATCATCCAgTAAACCgCC
47	CCTCgTAAATCCTCATCA	AA	TTTAACAaggCAAAATATAAATATAT	48	AACTAAgCTAACACTgCTCACTTTC	AA	ATCATCCAgTAAACCgCC
51	CCTCgTAAATCCTCATCA	AA	TACAAATggAATTTTCATCTTgTTC	52	ATggATCATCAATTTTgTCTCACTA	AA	ATCATCCAgTAAACCgCC
55	CCTCgTAAATCCTCATCA	AA	CTATTCTCACCCCTCCTAACAAAgt	56	TAgACATCTgATCCgTTCCTCAAgA	AA	ATCATCCAgTAAACCgCC
59	CCTCgTAAATCCTCATCA	AA	AggCCCTTgATAAAgAATggACATT	60	gCACTAgCACAAgTCTgCCATAAA	AA	ATCATCCAgTAAACCgCC
63	CCTCgTAAATCCTCATCA	AA	ggCTggggCTTTTTTgTTATAAgCC	64	gAgTgggAATCTTgAATgggAggAA	AA	ATCATCCAgTAAACCgCC
67	CCTCgTAAATCCTCATCA	AA	gTgAACAAATATAAgCATATTACTTA	68	TTCTTTAAAAAATAAAAAAAAg	AA	ATCATCCAgTAAACCgCC
71	CCTCgTAAATCCTCATCA	AA	AATTgTgACAAAACCTATACCgAgAg	72	gTTATgTAGACTTTgATCTAATCT	AA	ATCATCCAgTAAACCgCC

Organism: *E. coli*Target mRNA: enhanced green fluorescent protein (*Tg(EGFP)*)

Probe set: 12 split-initiator probe pairs (each probe carries half an HCR initiator)

HCR amplifier: B3-Alexa594 (Figures 6A and S22)

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	gTCCCTgCCTCTATATCT	TT	TgAAAAGTTCTTCTCCTTACgCAT	2	ATTCAACAAGAATTgggACAACCTCC	TT	CCACTCAACTTTAACCCg
3	gTCCCTgCCTCTATATCT	TT	ATTTgTgCCCATTAACATCACCATC	4	CACCTTCACCCCTCTCCACTgACAgA	TT	CCACTCAACTTTAACCCg
5	gTCCCTgCCTCTATATCT	TT	TAAgggTAAgTTTTCCgTATgTTgC	6	gTAgTTTTCCAgTAgTgCAAATAAA	TT	CCACTCAACTTTAACCCg
7	gTCCCTgCCTCTATATCT	TT	TAgTgACAAgTgTTggCCATggAAC	8	CAAAGCATTgAACACCATAACCgAA	TT	CCACTCAACTTTAACCCg
9	gTCCCTgCCTCTATATCT	TT	gCTgTTTCATATgATCTgggTATCT	10	CgggCATggCACTCTTgAAAAgTC	TT	CCACTCAACTTTAACCCg
11	gTCCCTgCCTCTATATCT	TT	ATATAgTTCTTTCTgTACATAACC	12	TCTTgTAgTTCCCGTCATCTTTgAA	TT	CCACTCAACTTTAACCCg
13	gTCCCTgCCTCTATATCT	TT	CACCTTCAAACCTgACTTCAgCACg	14	TTAACTCgATTCTATTAACAAgggT	TT	CCACTCAACTTTAACCCg
15	gTCCCTgCCTCTATATCT	TT	TTCCATCTTCTTTAAAATCAATACC	16	TgTATTCCAATTTgTgTCCAAGAAT	TT	CCACTCAACTTTAACCCg
17	gTCCCTgCCTCTATATCT	TT	TgATgTATACATTgTgTgAgTTATA	18	TgATTCCATTCCTTTgTTTgTCTgC	TT	CCACTCAACTTTAACCCg
19	gTCCCTgCCTCTATATCT	TT	TgTTgTgTCTAATTTTgAAgTTAAC	20	CTgCTAgTTgAACgCTTCCAICTTC	TT	CCACTCAACTTTAACCCg
21	gTCCCTgCCTCTATATCT	TT	CAATTggAgTATTTTgTTgATAATg	22	TgTCTggTAAAAGgACAgggCCATC	TT	CCACTCAACTTTAACCCg
23	gTCCCTgCCTCTATATCT	TT	gggCAgATTgTgTggACAggTAATg	24	CTCTCTTTTCgTTgggATCTTTCgA	TT	CCACTCAACTTTAACCCg

Organism: *E. coli*Target mRNA: **GTP-binding protein chain elongation factor EF-G (*fusA*)**Probe set: **18 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B2-Alexa594 (Figures 6B and S26)**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	CCTCgTAAATCCTCATCA	AA	gTgCgATgggTgTTgTACgAgCCAT	2	gCgCACTgATACCgATgTTACggTA	AA	ATCATCCAgTAAACCgCC
5	CCTCgTAAATCCTCATCA	AA	CATgAACTTACCgATTTTATggTT	6	CCATCCAgTCCATggTTgCAGCgCC	AA	ATCATCCAgTAAACCgCC
9	CCTCgTAAATCCTCATCA	AA	gCTCATACTgCTTAgCCATACCgA	10	gggTgTCgATgATgTTgATgCgATg	AA	ATCATCCAgTAAACCgCC
13	CCTCgTAAATCCTCATCA	AA	CAACTgCgCAGTAAACCATTACCgC	14	CggTTTCAGACTgCggCTgAACACC	AA	ATCATCCAgTAAACCgCC
17	CCTCgTAAATCCTCATCA	AA	TCAggAAgTTCgCACCCATgCggTC	18	gACgggTTTTgATCTgTAAACAAC	AA	ATCATCCAgTAAACCgCC
21	CCTCgTAAATCCTCATCA	AA	TCATTTTCACCggtCAACAACACC	22	ggTCAGCgTCgTTCAGTTgATAgC	AA	ATCATCCAgTAAACCgCC
25	CCTCgTAAATCCTCATCA	AA	ATTCgATCAGgTTCTggTgCCATTC	26	TCAGCTCTTCAGAAgCTTCAGCTgC	AA	ATCATCCAgTAAACCgCC
29	CCTCgTAAATCCTCATCA	AA	TTTCgTTgTTCAGAACgCgCTgACg	30	ACgCAGAACCCAGgTTACCAGgAT	AA	ATCATCCAgTAAACCgCC
33	CCTCgTAAATCCTCATCA	AA	CgTTgATCgCAGgTACgTCAACCgg	34	gAgTgTCTTTACCgTCgTCCAggAT	AA	ATCATCCAgTAAACCgCC
37	CCTCgTAAATCCTCATCA	AA	ggTTACCAACAACgggTCggTAgC	38	CACCggAgTAAACACggAAgAAggT	AA	ATCATCCAgTAAACCgCC
41	CCTCgTAAATCCTCATCA	AA	CgATAgCAGCAGCgATgTCgCCCgC	42	TgTCACCAGTggTTACgTCTTTCAg	AA	ATCATCCAgTAAACCgCC
45	CCTCgTAAATCCTCATCA	AA	TCggTTCAACTgCgATggAgATTAC	46	CCATTTTTTCCTggTCAGCTTTggT	AA	ATCATCCAgTAAACCgCC
49	CCTCgTAAATCCTCATCA	AA	ATTCACgCTTCATACggTCAACgAT	50	gTTTACCTACgTTCgCTTCAACgTT	AA	ATCATCCAgTAAACCgCC
53	CCTCgTAAATCCTCATCA	AA	CACgACCACCAgACTgTTTCgCgTg	54	TgTCgATAACAACATgACCATACTg	AA	ATCATCCAgTAAACCgCC
57	CCTCgTAAATCCTCATCA	AA	ATTCgCCAaggATTACACCACCTTT	58	ggATACCTTTATCAACggCCgggAT	AA	ATCATCCAgTAAACCgCC
61	CCTCgTAAATCCTCATCA	AA	CATggTAAgAACCGAAgTgCAGACg	62	TAAACgCAgTTCAGAggAgTCAAC	AA	ATCATCCAgTAAACCgCC
65	CCTCgTAAATCCTCATCA	AA	CTTCAACCTTCATgATCggCTCAAg	66	CACCggTgTTCTCTTCcggAgTTTC	AA	ATCATCCAgTAAACCgCC
69	CCTCgTAAATCCTCATCA	AA	CAGCgTggATCTTAAACgCCAgTAAC	70	ATCCgAACATTTTCAGACAgCggTAC	AA	ATCATCCAgTAAACCgCC

Organism: *E.coli*Target mRNA: **GTP-binding protein chain elongation factor EF-G *fusA***Probe set: **18 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B3-Alexa488 (Figures 6B, S26, and S28)**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
3	gTCCCTgCCTCTATATCT	TT	TAgTAgTggTTTTACCggCgTCgAT	4	CACCggTgTAgAACAgAATACgTTC	TT	CCACTCAACTTTAACCCg
7	gTCCCTgCCTCTATATCT	TT	TggTAATACCACgTTCCTgCTCCTg	8	AgAATgCAGTAGTCgCAGCggAAgT	TT	CCACTCAACTTTAACCCg
11	gTCCCTgCCTCTATATCT	TT	CTTCgATTgTgAAgTCAACgTgCCC	12	CATCgAgAACACgCATggAACgTTC	TT	CCACTCAACTTTAACCCg
15	gTCCCTgCCTCTATATCT	TT	CTTTATATTTgTTTgCCTgACgCCA	16	TTTTgTTAACgAACgCAATgCgCgg	TT	CCACTCAACTTTAACCCg
19	gTCCCTgCCTCTATATCT	TT	gCTgCAGCggAACCCgggTTCgCgCC	20	TgAAATgTTCTTCAGCACCAATCgC	TT	CCACTCAACTTTAACCCg
23	gTCCCTgCCTCTATATCT	TT	TATCTTCgTATTCgAAggTTACgCC	24	TAgCCAgTTCAACCATgTCTgCCgg	TT	CCACTCAACTTTAACCCg
27	gTCCCTgCCTCTATATCT	TT	gTTCTTCACCACCCAggTATTTTTC	28	gAgCACCTTTgATTTCTgCTTCAgT	TT	CCACTCAACTTTAACCCg
31	gTCCCTgCCTCTATATCT	TT	gCATCgCCTgAACACCTTTgTTCTT	32	ATggCAGgTAATCAATTACCgCATC	TT	CCACTCAACTTTAACCCg
35	gTCCCTgCCTCTATATCT	TT	CgTCATCACTTgCgTgACgTTCAgC	36	TTTTgAACgCCAgTgCAGAgAACgg	TT	CCACTCAACTTTAACCCg
39	gTCCCTgCCTCTATATCT	TT	TCAgTACggTATCACCAGgTTAAC	40	AACgCTCACgTgCAGCTTTCACggA	TT	CCACTCAACTTTAACCCg
43	gTCCCTgCCTCTATATCT	TT	TgATCggCgCATCCgggTCACACAg	44	gCTCAGggAATTCATACgTTCCA	TT	CCACTCAACTTTAACCCg
47	gTCCCTgCCTCTATATCT	TT	CTTTAgCCAgACggCCCAgAgCCA	48	CAGTCCATACACggAAAAGCgggTC	TT	CCACTCAACTTTAACCCg
51	gTCCCTgCCTCTATATCT	TT	ggATAgTTTCACggTAAgCAACCTg	52	TACCTTCAACATCggTAACCTTCTg	TT	CCACTCAACTTTAACCCg
55	gTCCCTgCCTCTATATCT	TT	ggTTTgAACCCggCTCCAgCgggTA	56	TgTCgTTgATgAACTCgTAGCCTTT	TT	CCACTCAACTTTAACCCg
59	gTCCCTgCCTCTATATCT	TT	CCAgCggACCTgCTTTCAGCTgTTC	60	TACCCATgTCTACTACCgggTAgCC	TT	CCACTCAACTTTAACCCg
63	gTCCCTgCCTCTATATCT	TT	CTTTAAAgCgATAgAAgCAgCCA	64	gAACTggTTTCgCTTCTTAAAgCC	TT	CCACTCAACTTTAACCCg
67	gTCCCTgCCTCTATATCT	TT	gACgACggCTCAAgTCACCgATAAC	68	CAGATTCTgACCTTTgAgCATAACC	TT	CCACTCAACTTTAACCCg
71	gTCCCTgCCTCTATATCT	TT	TggTCAGAgAACgCAGCTgAgTTgC	72	ATTCATAgTgTATgATgCACgACC	TT	CCACTCAACTTTAACCCg

Organism: *E. coli*Target mRNA: **isocitrate dehydrogenase (*icd*)**Probe set: **20 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B1-Alexa594 (Figure S28)**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	gAggAggggCagCAAACgg	AA	CCggAACAACTACTTTACTTTCCAT	2	TTTgCaggggTgATCTTCTTgCCTTg	TA	gAagAgTCTTCCTTTACg
3	gAggAggggCagCAAACgg	AA	gATTTTCagAACgTTgAgTTTgCC	4	CATCACCTTCAATgTAAgggATAAT	TA	gAagAgTCTTCCTTTACg
5	gAggAggggCagCAAACgg	AA	TggCTggggTTACATCTACACCgAT	6	CgACTgCagCgTCgACCACTTTCag	TA	gAagAgTCTTCCTTTACg
7	gAggAggggCagCAAACgg	AA	TTTTACgCTCgCCTTTATAggCTTT	8	CACCggTgTAAATTTCCATCCAggA	TA	gAagAgTCTTCCTTTACg
9	gAggAggggCagCAAACgg	AA	ACCTgACCATAAACCTgTgTgATTT	10	CAAgAgTTTCagCagCagCCAgAC	TA	gAagAgTCTTCCTTTACg
11	gAggAggggCagCAAACgg	AA	TggCAACgCgATATTCACgAATCag	12	CAACCggAgTggTCagCggACCTTT	TA	gAagAgTCTTCCTTTACg
13	gAggAggggCagCAAACgg	AA	CAACgTTCagAgAgCgAATACCgCC	14	TgTAgAgATCCagTTCCTggCgCag	TA	gAagAgTCTTCCTTTACg
15	gAggAggggCagCAAACgg	AA	gATAgTAACgTACCggACgCagCA	16	ggTgTTTAAACCgggCTTggAgTgCC	TA	gAagAgTCTTCCTTTACg
17	gAggAggggCagCAAACgg	AA	ggAAgATAACCATATCggTCagTTC	18	CCgCATAAATgTCTTCCgAgTTTTTC	TA	gAagAgTCTTCCTTTACg
19	gAggAggggCagCAAACgg	AA	CggCagAgTCTgCTTTCCATTCgAT	20	gCaggAATTTAATCACTTTCTCggC	TA	gAagAgTCTTCCTTTACg
21	gAggAggggCagCAAACgg	AA	gAATTTTCTTCAACCCCATCTCTTC	22	TACCgATACCACAATgTTCCgggAA	TA	gAagAgTCTTCCTTTACg
23	gAggAggggCagCAAACgg	AA	TggTgCCTTCTTCCgAACACggCTT	24	ATTcGATCgCgCgCgAACCgACg	TA	gAagAgTCTTCCTTTACg
25	gAggAggggCagCAAACgg	AA	CagAgTCACgATCgTTAgCAATTgC	26	TgATgTTgCCTTTgTgCACCgAgT	TA	gAagAgTCTTCCTTTACg
27	gAggAggggCagCAAACgg	AA	CTTTAAACgCTCCTTCggTgAACTT	28	CTTCACgCgCCAgCTggTAgCCCCA	TA	gAagAgTCTTCCTTTACg
29	gAggAggggCagCAAACgg	AA	CACCgTCgATCagTTCACCgCCAAA	30	TCgggTTTTTAACTTTCAgCCACgg	TA	gAagAgTCTTCCTTTACg
31	gAggAggggCagCAAACgg	AA	CTTTAATgACgATCTCTTTgCCAgt	32	gTTgCagAAATgCATCagCAATCAC	TA	gAagAgTCTTCCTTTACg
33	gAggAggggCagCAAACgg	AA	CATATTCagCCggACgCagCagAT	34	CgTTCagggTTCATACagCgATAAC	TA	gAagAgTCTTCCTTTACg
35	gAggAggggCagCAAACgg	AA	CTgCCAaggCgTCagAAATgTAgTC	36	gggCgATACCgATACCgCCAACCTg	TA	gAagAgTCTTCCTTTACg
37	gAggAggggCagCAAACgg	AA	CgCATTCgTCACCgATgTTTgCACC	38	CagTACCgTgggTggCTTCAAACAg	TA	gAagAgTCTTCCTTTACg
39	gAggAggggCagCAAACgg	AA	CTTgTCTTgACCggCATATTTTCgg	40	CggAgAgAATAATAgAgCCAaggATT	TA	gAagAgTCTTCCTTTACg

S4.9 Split-initiator probes for Figures 7, S29, and S30

Organism: *H. sapiens sapiens*

Target mRNA: **B-Raf proto-oncogene, serine/threonine kinase (BRAF)**

Probe set: **23 split-initiator probe pairs (each probe carries half an HCR initiator)**

HCR amplifier: **B3-Alexa647**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	gTCCCTgCCTCTATATCT	TT	ACCgCTCAGCgCCgCCATCTTATAA	2	gCCCggCTCCgCgCCgCCACCACCg	TT	CCACTCAACTTTAACCCg
5	gTCCCTgCCTCTATATCT	TT	ggCAGggTCCgCAGCCgAAgAggCC	6	TTTgATATTCCACACCTCCTCCggA	TT	CCACTCAACTTTAACCCg
9	gTCCCTgCCTCTATATCT	TT	ATATATTgATgTgATTATgCTCC	10	gCTggTgTATTCTTCATAggCCTCC	TT	CCACTCAACTTTAACCCg
13	gTCCCTgCCTCTATATCT	TT	AgAgCTAgAAACAgAAAAATCAGTT	14	AgAAgATgTAACggTATCCATTgAT	TT	CCACTCAACTTTAACCCg
17	gTCCCTgCCTCTATATCT	TT	CTTggggTTgCTCCgTgCCACATCT	18	gACTCTAACgATAggTTTTgTggT	TT	CCACTCAACTTTAACCCg
21	gTCCCTgCCTCTATATCT	TT	CTCTCCATCCTgAATTCTgTAAACA	22	ATCAGTgTCCCAACCAATTggTTTC	TT	CCACTCAACTTTAACCCg
25	gTCCCTgCCTCTATATCT	TT	TTTTcGTACAAAgTTgTgTTgTA	26	gTCACAAAATgCTAAggTgAAAAAC	TT	CCACTCAACTTTAACCCg
29	gTCCCTgCCTCTATATCT	TT	AACTTCTgTACTACAACgCTggTgA	30	TTggTCATAATTAACACACATCAGT	TT	CCACTCAACTTTAACCCg
33	gTCCCTgCCTCTATATCT	TT	TAgggCAGTCTCTgCTAAggACgCC	34	gggTgCggAAggggATgATCCAgAT	TT	CCACTCAACTTTAACCCg
37	gTCCCTgCCTCTATATCT	TT	TggTCggAAgggCTgTggAATTggA	38	AAATTgATTTCgATgATCTTCATCT	TT	CCACTCAACTTTAACCCg
41	gTCCCTgCCTCTATATCT	TT	TCTAATCAAgtCATCAATATTgACA	42	TCTCCATCACACgAAATCCTTgg	TT	CCACTCAACTTTAACCCg
45	gTCCCTgCCTCTATATCT	TT	TggAgATTCTgTAAggCTTTCACg	46	AgATgACTTCCTTTCCTgCTgAggT	TT	CCACTCAACTTTAACCCg
49	gTCCCTgCCTCTATATCT	TT	CTgCCCATCAGgAATCTCCCAATCA	50	AgATCCAATTCTTTgTCCCACTgTA	TT	CCACTCAACTTTAACCCg
53	gTCCCTgCCTCTATATCT	TT	AggTgTAggTgCTgTCACATTCAAC	54	TTCATTTTTgAAggCTTgTAACTgC	TT	CCACTCAACTTTAACCCg
57	gTCCCTgCCTCTATATCT	TT	ggTCTCAATgATATggAgATggTgA	58	ATCTATAAgTTTgATCATCTCAAAT	TT	CCACTCAACTTTAACCCg
61	gTCCCTgCCTCTATATCT	TT	CACTgTAGCTAgACCAAAATCACCT	62	CTgATgggACCACTCCATCgAgAT	TT	CCACTCAACTTTAACCCg
65	gTCCCTgCCTCTATATCT	TT	CTgAAAgCTgTATggATTTTTATCT	66	AACAATTCAAATgCATATACATCT	TT	CCACTCAACTTTAACCCg
69	gTCCCTgCCTCTATATCT	TT	ACTAAAATCCTCTgTTTTggAAACCA	70	TgTTTTTggAgAAgCACAAGCATAT	TT	CCACTCAACTTTAACCCg
73	gTCCCTgCCTCTATATCT	TT	TTTTgTTgCTACTCTCCTgAACTCT	74	AAgCAAACATATgTTCATTTATTTT	TT	CCACTCAACTTTAACCCg
77	gTCCCTgCCTCTATATCT	TT	ATTATATCTAgTCTTTAACCACACA	78	TAAgTATAAATTTTAgTTTggggAA	TT	CCACTCAACTTTAACCCg
81	gTCCCTgCCTCTATATCT	TT	AAgTAAAgCCTCTAgAAgAggCTCT	82	AAgTgAATgATACAAACCCggAACA	TT	CCACTCAACTTTAACCCg
85	gTCCCTgCCTCTATATCT	TT	TCTTCTggAgTCCCTAgTggACATg	86	CTgCAAACACAggCATAggTAgggT	TT	CCACTCAACTTTAACCCg
89	gTCCCTgCCTCTATATCT	TT	ATTAAATCTACTgACTTCCTAAAT	90	AAAATTATTAAgAATAATAATAgAA	TT	CCACTCAACTTTAACCCg

Organism: *H. sapiens sapiens*Target mRNA: **B-Raf proto-oncogene, serine/threonine kinase (*BRAF*)**Probe set: **23 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B4-Alexa546**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
3	CCTCAACCTACCTCCAAC	AA	CTCCATgTCCCCgTTgAACAgAgCC	4	ggCgCCgCgCCgCgCCgCCTCg	AT	TCTCACCATATTCgCTTC
7	CCTCAACCTACCTCCAAC	AA	ATgTTCCTgTgTCAACTAATCATT	8	ACCAAATTTgTCCAATAgggCCTCT	AT	TCTCACCATATTCgCTTC
11	CCTCAACCTACCTCCAAC	AA	TTCTCTTgTTggAgTgCATCTAgC	12	gTCCCCAgAgATTCCAATAACTgT	AT	TCTCACCATATTCgCTTC
15	CCTCAACCTACCTCCAAC	AA	AggTAgCACTgAAAaggCTAgAAgAg	16	gggATTTTgAAAACTgAAAgAgAT	AT	TCTCACCATATTCgCTTC
19	CCTCAACCTACCTCCAAC	AA	CACTgTCCCTgTgTTTgTgggCagg	20	gACTgTAACTCCACACCTTgCaggT	AT	TCTCACCATATTCgCTTC
23	CCTCAACCTACCTCCAAC	AA	CAATTCTTCTCCAgTAAgCCAaggAA	24	TggAACATTCTCCAACACTTCCACA	AT	TCTCACCATATTCgCTTC
27	CCTCAACCTACCTCCAAC	AA	ACCCTggAAAAGCAgCTTTCgACAA	28	TTTATAACCACATgTTTgACAgCgg	AT	TCTCACCATATTCgCTTC
31	CCTCAACCTACCTCCAAC	AA	gAACTTggAgACAAACAgCAAATCA	32	TTCCCTgTggTATTgggTggTgTTCA	AT	TCTCACCATATTCgCTTC
35	CCTCAACCTACCTCCAAC	AA	AAATTTggggCCCAATAgAgTCCgAg	36	ggATTTTgAAggAgACgACTggTg	AT	TCTCACCATATTCgCTTC
39	CCTCAACCTACCTCCAAC	AA	AgCTgATgAggATCggTCTCgTTgC	40	TTCTATTgTgTTATATgCACATTg	AT	TCTCACCATATTCgCTTC
43	CCTCAACCTACCTCCAAC	AA	gggggTAgCAGACAAACCTgTggTT	44	AgTTAgTgAgCCAaggTAATgAggCA	AT	TCTCACCATATTCgCTTC
47	CCTCAACCTACCTCCAAC	AA	CATTCgATTCCCTgTCTTCTgAggAT	48	ACTCgAgTCCCgTCTACCAAgTgTT	AT	TCTCACCATATTCgCTTC
51	CCTCAACCTACCTCCAAC	AA	TCCCTTgTAgACTgTTCCAATgAT	52	TTTCACTgCCACATCACCATgCCAC	AT	TCTCACCATATTCgCTTC
55	CCTCAACCTACCTCCAAC	AA	ATgTCgTgTTTTCTgAgTACTCCT	56	ATAgCCATgAAgAgTAggATATTC	AT	TCTCACCATATTCgCTTC
59	CCTCAACCTACCTCCAAC	AA	TATATTACTCTTgAggTCTCTg	60	TTTTACTgTgAggTCTTCATgAAgA	AT	TCTCACCATATTCgCTTC
63	CCTCAACCTACCTCCAAC	AA	CAAAATggATCCAgACAACCTgTTCA	64	CATTCTgATgACTTCTgTgCCATC	AT	TCTCACCATATTCgCTTC
67	CCTCAACCTACCTCCAAC	AA	CTTCATggCTTTTggACAgTTACTC	68	CTTTTTgAggCACTCTgCCATTAAT	AT	TCTCACCATATTCgCTTC
71	CCTCAACCTACCTCCAAC	AA	CgCACCATATCCCCCTgCCTggATg	72	CACTCATTTgTTTCAGTggACAggA	AT	TCTCACCATATTCgCTTC
75	CCTCAACCTACCTCCAAC	AA	gAgAgTATTTTTATCAATTTAACAT	76	TgTTCTTTggTTCACCTAAAAAAA	AT	TCTCACCATATTCgCTTC
79	CCTCAACCTACCTCCAAC	AA	AACCCTTggATgTTAAAAATCCAAT	80	CAATTTTAgCAATgTCTATgTATT	AT	TCTCACCATATTCgCTTC
83	CCTCAACCTACCTCCAAC	AA	AACTgAAgTTTACTACTTAAAATAA	84	ATAgCTggCAACAAAAGTTgCATgA	AT	TCTCACCATATTCgCTTC
87	CCTCAACCTACCTCCAAC	AA	CAggCTAACCGACTgCCAACCTCTC	88	gATCTgTTCAGTTTgCCTTATCTAA	AT	TCTCACCATATTCgCTTC
91	CCTCAACCTACCTCCAAC	AA	ATTgTTATAAAAAGAAATAgTTATA	92	gAAATAAAAAGACATCCACATTTTCC	AT	TCTCACCATATTCgCTTC

Organism: *G. gallus domesticus*Target mRNA: **diencephalon/mesencephalon homeobox 1 (*Dmbx1*)**Probe set: **25 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B1-Alexa594**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
1	gAggAggggCagCAAACgg	AA	CgCTCagTgAgTTCATggCATgCag	2	CTgCCTgCTggTgCaggTTgTACAT	TA	gAAgAgTCTTCCTTTACg
5	gAggAggggCagCAAACgg	AA	AAATgATgTCAgCCAgTCTCTCCgC	6	ggTgCTgggATCCATAACgTgCCTC	TA	gAAgAgTCTTCCTTTACg
9	gAggAggggCagCAAACgg	AA	AAATgATgTCAgCCAgTCTCTCCgC	10	gCTTCTgCagCTgCTCCTTCTggAg	TA	gAAgAgTCTTCCTTTACg
13	gAggAggggCagCAAACgg	AA	CAgTgTCAggTATTgTggACTgggT	14	CCTCACTTgggATgCTCTgAgTCTT	TA	gAAgAgTCTTCCTTTACg
17	gAggAggggCagCAAACgg	AA	CCTCCTCTCTgTCAgTCTgTCCCTC	18	TAgCCTCATCCAAggTgCTCTTAAA	TA	gAAgAgTCTTCCTTTACg
21	gAggAggggCagCAAACgg	AA	CgCTgATgggAgACTCTgATTTTgg	22	CACTgCTAgATgAAggAgTCACggT	TA	gAAgAgTCTTCCTTTACg
25	gAggAggggCagCAAACgg	AA	CTgCCATgTgCTggCggAATTgCTC	26	AggAgTAATggACCAAgTTgTTggT	TA	gAAgAgTCTTCCTTTACg
29	gAggAggggCagCAAACgg	AA	gACAgTgCagggAgCTCagAggCgC	30	CgTgggAgAgAgATTggTAgTACgA	TA	gAAgAgTCTTCCTTTACg
33	gAggAggggCagCAAACgg	AA	TCTCAATACTTgTTgTTTTACTgTT	34	CATgCTgCTTTgCCCggAgCCTTAg	TA	gAAgAgTCTTCCTTTACg
37	gAggAggggCagCAAACgg	AA	gACCTCCggTgCATCTTCTTATggg	38	ggTTCCAggAgTgACATgTCTggTg	TA	gAAgAgTCTTCCTTTACg
41	gAggAggggCagCAAACgg	AA	CggTgCTAgTAAgACATTAgtAAAT	42	AgCCAgTAGCAgTgTCTgATgCAAT	TA	gAAgAgTCTTCCTTTACg
45	gAggAggggCagCAAACgg	AA	TCACAgCagTCCAAAgggACAgTTC	46	gCTCTTCTgAATgTTACAggCTTA	TA	gAAgAgTCTTCCTTTACg
49	gAggAggggCagCAAACgg	AA	gTTTTCCCAAAGAAATgCATCgACAA	50	TATgTACAAGACAAAAGCaggACTCT	TA	gAAgAgTCTTCCTTTACg
53	gAggAggggCagCAAACgg	AA	ggggAATAAAAAGCAAAAgAggCCAC	54	gACTAgCTACCAAAACTgAgAgAgA	TA	gAAgAgTCTTCCTTTACg
57	gAggAggggCagCAAACgg	AA	TTTgCTCTAAgCACCATTAAGACTC	58	gAgCagTgAATTgCATAATggTTTT	TA	gAAgAgTCTTCCTTTACg
61	gAggAggggCagCAAACgg	AA	ggAAgTgCTTAAACAggAAATTCAC	62	AgTAAAggAAAAACACTTgCCTTT	TA	gAAgAgTCTTCCTTTACg
65	gAggAggggCagCAAACgg	AA	AATTTgCTTTCATTTTTCTCCCA	66	AACAATCAAgTCAAAAgTAACCATg	TA	gAAgAgTCTTCCTTTACg
69	gAggAggggCagCAAACgg	AA	CTAgACCAAAATgCTCTCCAAAAG	70	AgTTTTTATTgTTCTCTATTgTCgA	TA	gAAgAgTCTTCCTTTACg
73	gAggAggggCagCAAACgg	AA	TAAgAACAgCTTgCATTAAATCgTgg	74	TAgAATTTggTgATCggAgCgTTTT	TA	gAAgAgTCTTCCTTTACg
77	gAggAggggCagCAAACgg	AA	CTTggCCTCCAgCATTgCagCATT	78	AATAgAAAAGCCCCgATTATCACCC	TA	gAAgAgTCTTCCTTTACg
81	gAggAggggCagCAAACgg	AA	gAAATCACTTTgCagTTggTgAgTT	82	AgAgAAATgAggCCAAAACTgTggA	TA	gAAgAgTCTTCCTTTACg
85	gAggAggggCagCAAACgg	AA	AAgTCCTTTgggTTggTAgAAAAT	86	TATTTggTTTggAgAAgATAAATAA	TA	gAAgAgTCTTCCTTTACg
89	gAggAggggCagCAAACgg	AA	gCATTTTTggTCTTAggCAATACTA	90	ACCTAgCACCTgCCACAgAgCCAgT	TA	gAAgAgTCTTCCTTTACg
93	gAggAggggCagCAAACgg	AA	gACCAgATgATggCCTgCagTgAAT	94	CTCCATTTCTTCTTAAATCgAgCA	TA	gAAgAgTCTTCCTTTACg
97	gAggAggggCagCAAACgg	AA	AggTAACTCAACCCAggCTTCTgC	98	AACACCCTTCCCCCTCgTgTTAA	TA	gAAgAgTCTTCCTTTACg

Organism: *G. gallus domesticus*Target mRNA: **diencephalon/mesencephalon homeobox 1 (*Dmbx1*)**Probe set: **25 split-initiator probe pairs (each probe carries half an HCR initiator)**HCR amplifier: **B2-Alexa647**

Odd #	1st Half of Initiator I1	Spacer	Probe Sequence (25 nt)	Even #	Probe Sequence (25 nt)	Spacer	2nd Half of Initiator I1
3	CCTCgTAAATCCTCATCA	AA	AgTCgggCgCgTgCTgTgCTTgCTg	4	gggTgAgggCgTgCACTgAgggCCg	AA	ATCATCCAgTAAACCgCC
7	CCTCgTAAATCCTCATCA	AA	AggCCgTgCgACTTCgCCTTgTTT	8	CCAgTgCCTCCAgCTgTTgggCAgT	AA	ATCATCCAgTAAACCgCC
11	CCTCgTAAATCCTCATCA	AA	CACTgTggCTgCTTTCACAgTCCTT	12	CCAgCACAggTggTTCcCgTCTCCC	AA	ATCATCCAgTAAACCgCC
15	CCTCgTAAATCCTCATCA	AA	ACAgggTTAggTTgAggTCTgTgCT	16	gCgCTgATTCACTggCTgACTgCTC	AA	ATCATCCAgTAAACCgCC
19	CCTCgTAAATCCTCATCA	AA	CgTCCACACCAggACTCTTgTCCAC	20	TCgCTCTCTTgCAgTTCAAAGCCTT	AA	ATCATCCAgTAAACCgCC
23	CCTCgTAAATCCTCATCA	AA	AggAgTAggAgTgAgTTTgAgCCA	24	gCAggCggAAgAggCTCAggggTgA	AA	ATCATCCAgTAAACCgCC
27	CCTCgTAAATCCTCATCA	AA	TggCAGACggTgTCCCCATTTcGAA	28	TgTTgACATTCATgCCCAAgTAggg	AA	ATCATCCAgTAAACCgCC
31	CCTCgTAAATCCTCATCA	AA	ggATgggTgAggACCAgACCTgCTg	32	ggCTTggAAgAgAgCTggAggCCTg	AA	ATCATCCAgTAAACCgCC
35	CCTCgTAAATCCTCATCA	AA	gTAgggTgTCAAgCCCAAgggATgC	36	ACggTTgCCCCTCTgCAgTCAgTT	AA	ATCATCCAgTAAACCgCC
39	CCTCgTAAATCCTCATCA	AA	ACTCCAaggAAgAgATgAgggTggAA	40	AAAgTTTCCCTgATAgggAgCACC	AA	ATCATCCAgTAAACCgCC
43	CCTCgTAAATCCTCATCA	AA	TCCTCCTCAAATATTTAAAgAAgAC	44	CTgTCTAAACACACATCCTCTCCCT	AA	ATCATCCAgTAAACCgCC
47	CCTCgTAAATCCTCATCA	AA	ACATTATCgCAgggATgAggTgAgg	48	AAAAAgggTgTATATAACACggTTg	AA	ATCATCCAgTAAACCgCC
51	CCTCgTAAATCCTCATCA	AA	gCggTggATgCTTCAACATTgTAA	52	TTCTgTAACACTgACAgTAACACAC	AA	ATCATCCAgTAAACCgCC
55	CCTCgTAAATCCTCATCA	AA	gggAgCgTggCTgATTTgTgACTTT	56	AACCCAAgAAgCAACTAgCTgTg	AA	ATCATCCAgTAAACCgCC
59	CCTCgTAAATCCTCATCA	AA	TCAgCTTTAgCAgAAgAgAgAAg	60	TTgCATCATTTCTgCCgTTATAA	AA	ATCATCCAgTAAACCgCC
63	CCTCgTAAATCCTCATCA	AA	TCTgTCTgTgAAACAAGTgCTATTAg	64	CAgCAgCATTTggCCAgCATTTgT	AA	ATCATCCAgTAAACCgCC
67	CCTCgTAAATCCTCATCA	AA	TTACACTTCACTgAAgACCAAAGg	68	AACCCATAATTTgTAAATgggggA	AA	ATCATCCAgTAAACCgCC
71	CCTCgTAAATCCTCATCA	AA	gTATgAACACAgTgggAgTTCATAC	72	TTgTCAAaggAACCATATAATTCATg	AA	ATCATCCAgTAAACCgCC
75	CCTCgTAAATCCTCATCA	AA	AgggTCTgAAgCTgCACAgCTTgAg	76	CACTTgTTACATTTCTCACTTgCTAA	AA	ATCATCCAgTAAACCgCC
79	CCTCgTAAATCCTCATCA	AA	CAAgCCAATCTACTCCTCgCTgCAg	80	ggTTgCTTggggACATggTACTTTT	AA	ATCATCCAgTAAACCgCC
83	CCTCgTAAATCCTCATCA	AA	TggATTACTAAATgAAgggTCATT	84	CTTCTCAAgAAggAAAAACACTCTg	AA	ATCATCCAgTAAACCgCC
87	CCTCgTAAATCCTCATCA	AA	TTCTTggTACggTgAgTTCAAAggA	88	TTAgATCTgggTTTCTCCTCCCT	AA	ATCATCCAgTAAACCgCC
91	CCTCgTAAATCCTCATCA	AA	gACCCgCCTgACACCCTTgAgATTC	92	gCCCAGCTCTgCTgCgTgTAgTgg	AA	ATCATCCAgTAAACCgCC
95	CCTCgTAAATCCTCATCA	AA	gTTATgTAggCTATgCACACgTTgC	96	TggTATgAAgTAAgATgggAgCAA	AA	ATCATCCAgTAAACCgCC
99	CCTCgTAAATCCTCATCA	AA	TTTTTAAgATgCATTATgCAgTTg	100	TgCCTCAgTTTAAgggATTTAgATg	AA	ATCATCCAgTAAACCgCC

References

- Bell, G.W., Yatskievych, T.A., & Antin, P.B.** (2004). GEISHA, a whole-mount in situ hybridization gene expression screen in chicken embryos. *Dev. Dyn.*, **229**(3), 677–687.
- Choi, H. M. T., Chang, J. Y., Trinh, L. A., Padilla, J. E., Fraser, S. E., & Pierce, N. A.** (2010). Programmable in situ amplification for multiplexed imaging of mRNA expression. *Nat. Biotechnol.*, **28**(11), 1208–12.
- Choi, H. M. T., Beck, V. A., & Pierce, N. A.** (2014). Next-generation in situ hybridization chain reaction: higher gain, lower cost, greater durability. *ACS Nano*, **8**(5), 4284–4294.
- Choi, H. M. T., Calvert, C. R., Husain, N., Huss, D., Barsi, J. C., Deverman, B. E., Hunter, R. C., Kato, M., Lee, S. M., Abelin, A. C. T., Rosenthal, A. Z., Akbari, O. S., Li, Y., Hay, B. A., Sternberg, P. W., Patterson, P. H., Davidson, E. H., Mazmanian, S. K., Prober, D. A., van de Rijn, M., Leadbetter, J. R., Newman, D. K., Readhead, C., Bronner, M. E., Wold, B., Lansford, R., Sauka-Spengler, T., Fraser, S. E., & Pierce, N. A.** (2016). Mapping a multiplexed zoo of mRNA expression. *Development*, **143**, 3632–3637.
- Coupric, M., & Bertrand, G.** (1997). Topological gray-scale watershed transform. *Pages 136–146 of: Vision geometry v. Proceedings of SPIE.*
- Darnell, D. K., Kaur, S., Stanislaw, S., Davey, S., Konieczka, J. H., Yatskievych, T. A., & Antin, P. B.** (2007). GEISHA: an in situ hybridization gene expression resource for the chicken embryo. *Cytogenet. Genome Res.*, **117**(1-4), 30–35.
- McEntyre, J., & Ostell, J.** (2002). *The NCBI handbook [Internet]*. Bethesda, MD: National Center for Biotechnology Information (US).
- Shah, S., Lubeck, E., Schwarzkopf, M., He, T.-F., Greenbaum, A., Sohn, C. H., Lignell, A., Choi, H. M. T., Gradinaru, V., Pierce, N. A., & Cai, L.** (2016). Single-molecule RNA detection at depth via hybridization chain reaction and tissue hydrogel embedding and clearing. *Development*, **143**, 2862–2867.
- Trivedi, V., Choi, H. M. T., Fraser, S. E., & Pierce, N. A.** (2018). Multidimensional quantitative analysis of mRNA expression within intact vertebrate embryos. *Development*, **145**, DOI:10.1242/dev.156869.
- Yoo, T.S., Ackerman, M.J., Lorenzen, W.E., Schroeder, W., Chalana, V., Aylward, S., Metaxas, D., & Whitaker, R.** (2002). Engineering and algorithm design for an image processing API: a technical report on ITK - the Insight Toolkit. *Studies in Health Technology and Informatics*, **85**, 586–592.