

S3 Table.

	PT vs. S	DT vs. S
Genera enriched in plaque* (Genera with a significantly higher abundance in PT or DT groups than in the S group)	<i>Fusobacterium,</i> <i>Capnocytophaga,</i> <i>Actinomyces,</i> <i>Leptotrichia,</i> <i>Corynebacterium,</i> <i>Aggregatibacter,</i> <i>Campylobacter,</i> <i>Tannerella,</i> <i>Gemella,</i> <i>Selenomonas</i>	<i>Fusobacterium,</i> <i>Capnocytophaga,</i> <i>Lautropia,</i> <i>Corynebacterium,</i> <i>Aggregatibacter,</i> <i>Campylobacter,</i> <i>Tannerella,</i> <i>Eikenella,</i> <i>Paludibacter,</i> <i>Kingella,</i> <i>Gemella,</i> <i>Selenomonas,</i> <i>Treponema</i>
Genera enriched in saliva* (Genera with a significantly higher abundance in the S group than in either the PT or DT groups)	<i>Streptococcus,</i> <i>Haemophilus,</i> <i>Granulicatella,</i> <i>Rothia,</i> <i>[Prevotella],</i> <i>Oribacterium,</i> <i>Bulleidia</i>	<i>Streptococcus,</i> <i>Granulicatella,</i> <i>Rothia,</i> <i>[Prevotella],</i> <i>Veillonella,</i> <i>Oribacterium,</i> <i>Bulleidia,</i> <i>Moryella</i>