

Chemotaxonomy of Mycotoxigenic Small-Spored *Alternaria* Fungi – Do Multitoxin Mixtures Act as an Indicator for Species Differentiation?

Theresa Zwickel^{1,2*}, Sandra M. Kahl^{3,4}, Michael Rychlik² and Marina E. H. Müller^{3*}

¹ Federal Institute for Risk Assessment (BfR), Berlin, Germany, ² Chair of Analytical Food Chemistry, Technical University of Munich, Munich, Germany, ³ Leibniz Centre for Agricultural Landscape Research (ZALF), Müncheberg, Germany, ⁴ University of Potsdam, Institute of Biochemistry und Biology, Potsdam, Germany

*Correspondence: Marina E. H. Müller; mmueller@zalf.de

Table-S3: Limits of detection (LODs) and limits of quantification (LOQs) determined according to DIN EN ISO 32645 (10 calibration points, Mandel fitting test at a significance level of 0.01 showed linear regression for all analytes, $R^2 > 0.96$ for all analytes, confidence interval and t-test at a significance level of 0.05, maximal permissible uncertainty was set to 33.33%)

Analyte	range [$\mu\text{g/L}$]	LOD [$\mu\text{g/kg}$]	LOQ [$\mu\text{g/kg}$]
ALT	0.25 – 4.5	3.0	9.5
AME	0.25 – 4.5	0.5	1.8
AOH	0.50 – 4.5	0.6	2.1
TEA	0.25 – 4.5	1.4	4.6
TEN	0.25 – 4.5	0.9	3.1
AAL TB	0.25 – 4.5	1.7	5.5
ATX-I	0.50 – 4.5	1.1	3.7
ATX-II	0.50 – 4.5	1.0	3.4
STTX-III	0.25 – 4.5	0.9	3.2
ALP	0.25 – 4.5	1.2	4.0
AA-III	0.25 – 4.5	0.9	3.1
ATL	0.50 – 4.5	1.2	3.7