

638

Table S1. Clinical characteristics for subjects included in third trimester placental analyses ($n = 36$).

	Preeclampsia ($n = 23$)	Healthy pregnancy ($n = 13$)
Baseline characteristics		
Maternal age, years	30 (6) ^a	32 (5) ^a
Primiparae, n (%)	12 (52) ^{b,*}	1 (8) ^b
Characteristics at delivery		
Body mass index, kg/m^2	29 (8) ^b	29 (6) ^b
Systolic blood pressure, mmHg	151 (12) ^{a,*}	119 (9) ^a
Diastolic blood pressure, mmHg	100 (8) ^{a,*}	71 (9) ^a
Gestational age delivery, weeks	32 (4) ^{b,*}	39 (1) ^b
Placental weight, g	350 (193) ^{b,*}	580 (241) ^b
Fetal birth weight, g	1780 (995) ^{b,*}	3590 (630) ^b
FGR, n (%)	12 (52) ^b	n.a.
Severe preeclampsia [†] , n (%)	16 (70) ^b	n.a.

Continuous variables are presented as mean with standard deviation (a) or median with interquartile range (b). FGR, fetal growth restriction; n.a.; not applicable.

Comparisons between outcome groups (two-sided T-tests or Mann-Whitney U-test for continuous variables, Chi-square test for categorical variables): * $P < 0.05$ vs healthy pregnancies. †Preeclampsia was sub-phenotyped as severe if diagnosed with one or more features (40, 64).

639

640