

1 **Scientific Reports**

2
3 **Thiamine tetrahydrofurfuryl disulfide promotes voluntary**
4 **activity through dopaminergic activation in the medial**
5 **prefrontal cortex**

6
7 **Masato Saiki^{1,#}, Takashi Matsui^{1,2,#}, Mariko Soya^{1,#}, Tomomi**
8 **Kashibe¹, Takeru Shima¹, Takeshi Shimizu³, Takehiro Naruto⁴,**
9 **Takahito Kitayoshi⁴, Kouji Akimoto⁴, Shinji Ninomiya⁴, and Hideaki**
10 **Soya^{1,2,*}**

11
12 *¹Laboratory of Exercise Biochemistry and Neuroendocrinology, Faculty of Health and Sport*
13 *Sciences, University of Tsukuba, Tsukuba, Ibaraki, Japan, 305-8574; ² Sports Neuroscience*
14 *Division, Advanced Research Initiative for Human High Performance (ARIHHP), University*
15 *of Tsukuba; Tsukuba, Ibaraki, Japan, 305-8574; ³Sports Research and Development Core,*
16 *University of Tsukuba, Tsukuba, Ibaraki, Japan, 305-8574; ⁴Takeda Consumer Healthcare*
17 *Company Limited, Tokyo, Japan, 100-0005. #These authors contributed equally to this work.*


18
19 ***Correspondence**

20 Hideaki Soya, Ph.D.

21 E-mail: soya.hideaki.gt@u.tsukuba.ac.jp

22 Tel/Fax: +81-29-853-2620

Supplementary Information



1 **Figure S1 Dopamine D1 and D2 receptor antagonists increase dopamine and serotonin**
 2 **release after TTFD injection. (A) Extracellular dopamine levels. (B) Extracellular serotonin**
 3 **levels. ** $P < 0.01$ versus saline + aCSF group (two-way ANOVA with Bonferroni's post hoc**
 4 **tests).**