Supplementary Table 1. Clinicopathologic and surgical *characteristics of cRCC patients in our surgical cohort*

| No. gender (%) | |
|--|----------------------|
| Male | 78 (57) |
| Female | 60 (43) |
| | |
| No. race (%) | |
| White | 132 (96) |
| Black | 2 (1) |
| Asian | 4 (3) |
| 14 III (10D) | 54.4 (45.5.00.0) |
| Median age at surgery, years (IQR) | 54.1 (45.5-63.9) |
| No survival task nimus (0/) | |
| No. surgical technique (%) Partial | 400 (00 4) |
| - 40 - 50 - 50 - 50 - 50 - 50 - 50 - 50 | 122 (88.4) |
| Radical | 16 (11.6) |
| Open | 113 (81.9) |
| Laparoscopic | 10 (7.2) |
| Robotic | 15 (10.9) |
| No precenting symptom (0/) | |
| No. presenting symptom (%) | 400 (00.4) |
| Incidental, asymptomatic | 122 (88.4) |
| Local | 16 (11.6) |
| No. histologic subtype (%) | |
| Clear cell renal cell carcinoma | 91 (65.9) |
| Multilocular cystic renal cell neoplasm of low malignant potential | 18 (13) |
| Benign multilocular cyst | 16 (11.6) |
| Papillary renal cell carcinoma | 5 (3.6) |
| Clear cell capillary renal cell carcinoma | 4 (2.9) |
| Unclassified renal cell carcinoma | 2 (1.4) |
| Mixed epithelial and stromal tumor family | 2 (1.4) |
| mixed opinional and stromar tumor runniy | Z (1. 4) |
| No. Fuhrman grade (%) | |
| 1 | 22 (15.9) |
| 2 | 85 (61.6) |
| 3 | 9 (6.5) |
| NA | 22 (15.9) |
| | 22 (10.0) |
| No. pathologic stage (%) | |
| pT1a | 97 (70.3) |
| pT1b | 15 (10.9) |
| pT2a | 3 (2.2) |

| рТ3а | 4 (2.9) |
|------|---------|
| pT3b | 1 (0.7) |