

## Habitat and landscape factors influence pollinators in a tropical megacity, Bangkok, Thailand

**Supplemental Table S3.** Detailed information about each of the pollinator taxa observed in this study. Taxa are sorted numerically by “Order” (highest to lowest percentage), then alphabetically by “Family”, then numerically by “Frequency”. The frequencies of some taxa are unknown, as they were only identified to genus or family in the field, and were later identified to species or genus in the lab based on collected or photographed specimens. The nine most common taxa are highlighted in yellow. Distribution information was obtained from the Catalogue of Life website (<http://www.catalogueoflife.org>; accessed May 2018). All pollinator taxa that could be sufficiently identified are native to Thailand.

Order	Family	Genus	Frequency (# of individuals)	Percent	Distribution
Hymenoptera (98.6%)	Apidae	<i>Tetragonula</i> spp.	12,087	64.317	native
		<i>Apis florea</i> Fabricius, 1787	2,441	12.989	native
		<i>Apis cerana</i> Fabricius, 1793	2,345	12.478	native
		<i>Apis dorsata</i> Fabricius, 1793	1,148	6.109	native
		<i>Xylocopa aestuans</i> Linnaeus, 1758	273	1.453	native
		<i>Xylocopa nasalis</i> Westwood, 1842	12	0.064	native
		<i>Ceratina</i> spp.	9	0.048	-
		<i>Ceratina cognata</i> Smith, 1879	(specimen)	-	native
		<i>Ceratina dentipes</i> Friese, 1914	(specimen)	-	native
	<i>Amegilla</i> spp.	7	0.037	-	
	Crabronidae	spp.	2	0.011	-
	Eumenidae	<i>Delta conoideum</i> Gmelin, 1790	32	0.170	native
		<i>Delta pyriforme</i> Fabricius, 1775	2	0.011	native
		<i>Polistes</i> spp.	2	0.011	-
	Halictidae	<i>Lasioglossum</i> spp.	84	0.447	-
		<i>Nomia</i> spp.	3	0.016	-
	Megachilidae	<i>Megachile</i> spp.	30	0.160	-
	Scoliidae	<i>Campsomeris</i> spp.	11	0.059	-
		spp.	6	0.032	-

	Unknown	spp.	27	0.144	-	
	Vespidae	spp.	15	0.080	-	
		<i>Ropalidia</i> spp.	(specimen)	-	-	
Lepidoptera (1.16%)	Hesperiidae	spp.	1	0.005	-	
	Lycaenidae	spp.	57	0.303	-	
		<i>Chilades pandava</i> Horsfield, 1829		14	0.074	native
	Nymphalidae	<i>Danaus chrysippus</i> Linnaeus, 1758		54	0.287	native
		<i>Junonia lemonias</i> Linnaeus, 1758		14	0.074	native
		<i>Ideopsis similis</i> Linnaeus, 1758		2	0.011	native
	Papilionidae	<i>Graphium doson</i> (Felder & Felder, 1864)		5	0.027	native
	Pieridae	<i>Delias eucharis</i> Drury, 1773		43	0.229	native
<i>Catopsilia Pomona</i> Fabricius, 1775			17	0.090	native	
	Unknown	spp.	11	0.059	-	
Diptera (0.15%)	Sacrophagidae	spp.	2	0.011	-	
	Syrphidae	spp.	20	0.106	-	
		<i>Eristalinus arvorum</i> Fabricius, 1787	(specimen)	-	-	native
	Unknown	spp.	6	0.032	-	
Coleoptera (0.032%)	Chrysomelidae	sp.	1	0.005	-	
	Curculionoidea	spp.	5	0.027	-	
Hemiptera (0.021%)	Lygaeidae	spp.	4	0.021	-	
Passeriformes (0.005%)	Nectariniidae	<i>Cinnyris jugularis</i> Linnaeus, 1766	1	0.005	native	