

Appendix S1
Ecological Applications

Estimating sensitivity of seabed habitats to disturbance by bottom trawling based on the longevity of benthic fauna.

Adriaan D. Rijnsdorp^{1,2}, Stefan G. Bolam³, Clement Garcia³, Jan Geert Hiddink⁴, Niels T. Hintzen¹, P. Daniel van Denderen⁵, Tobias van Kooten^{1,6}

¹ Wageningen Marine Research, P.O. Box 68, 1970 AB IJmuiden, The Netherlands;

² Aquaculture and Fisheries Group, Wageningen University, P.O. Box 338, 6700 AH Wageningen, The Netherlands;

³ The Centre for Environment, Fisheries and Aquaculture Science, Pakefield Road, Suffolk NR33 0HT, UK;

⁴ School of Ocean Sciences, Bangor University, Menai Bridge, Anglesey, LL59 5AB, UK ;

⁵ Centre for Ocean Life, National Institute of Aquatic Resources, Technical University of Denmark, Kemitorvet B-202, 2800 Kongens Lyngby, Denmark;

⁶ Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, The Netherlands.

Corresponding author: Adriaan D. Rijnsdorp (adriaan.rijnsdorp@wur.nl)




Figure S1. Frequency distribution of environmental variables of the benthos sampling stations (left) and the frequency distribution of environmental variables of all North Sea grid cells (right).