

Supplemental Material

Table S1. Clinical Characteristics of the Included Subjects and Excluded Subjects without baPWV Measurement.

Variables	Included (n=462)	Excluded (n=104)	P value
Age, yr	63.8±8.7	63.3±8.0	0.63
Men, n (%)	396 (85.7)	85 (81.7)	0.31
Body mass index, kg/m ²	24.8±3.6	25.2±3.8	0.34
Systolic blood pressure, mm Hg	129.0±16.2	125.9±15.6	0.08
Diastolic blood pressure, mm Hg	74.8±10.6	72.6±10.6	0.06
Heart rate, bpm	66.5±12.0	66.6±11.6	0.95
Total cholesterol, mg/dL	169.9±31.7	163.7±27.8	0.07
Triglycerides, mg/dL	138.0±94.0	133.1±62.1	0.61
HDL cholesterol, mg/dL	50.4±13.1	47.6±12.9	0.04
LDL cholesterol, mg/dL	92.8±27.4	89.5±25.3	0.26
Glucose, mg/dL	119.5±37.2	108.6±27.7	0.005
HbA1c, % (n=347)	6.4±1.0	6.4±1.0	0.96
Creatinine, mg/dL	0.85±0.24	0.88±0.25	0.37
eGFR, ml/min/1.73 m ²	71.9±17.6	68.9±15.4	0.11
Smoker, n (%)	313 (70.2)	78 (76.5)	0.20
Complications, n (%)			
Hypertension	433 (93.7)	100 (96.2)	0.32
Dyslipidemia	436 (94.4)	98 (94.2)	0.96
Diabetes mellitus	178 (38.5)	36 (34.6)	0.46
Prior myocardial infarction	244 (52.8)	43 (41.4)	0.03
Prior coronary intervention	404 (87.4)	97 (93.3)	0.07
Medication use, n (%)			
Antiplatelet drugs	454 (98.3)	100 (96.2)	0.21
ARBs/ACEIs	320 (69.3)	69 (66.4)	0.56
β-blockers	219 (47.4)	47 (45.2)	0.68
Antidiabetic drugs	145 (31.4)	31 (29.8)	0.75
Statins	394 (85.3)	89 (85.6)	0.94
Left ventricular ejection fraction, %	60.8±9.9	60.6±11.4	0.83
BNP, pg/mL	35.1±61.6	57.0±112.3	0.007
FMD, %	4.8±2.6	4.9±2.9	0.53

All results are presented as mean±SD.

HDL, high-density lipoprotein; LDL, low-density lipoprotein; eGFR, estimated glomerular filtration rate; ARB, angiotensin receptor blocker; ACEI, angiotensin-converting enzyme inhibitor; BNP, brain natriuretic peptide; FMD, flow-mediated vasodilation; baPWV, brachial ankle pulse wave velocity.