

## Web Appendix

**Table 1. Response Rate to IMCI Global Implementation Survey**

WHO Region	Total number of WHO MS	Member States (N) to which a survey questionnaire was sent	WHO MS that responded*	WHO MS that reported no IMCI (did not participate in the survey)	WHO MS that responded IMCI not in national plan/no district implementing nor component implemented	Response rate
African	47	47	45	4		95.7
Americas	35	33	19	4		57.6
Eastern Mediterranean	21	16	14			87.5
Europe	53	7	6			85.7
South-east Asia	11	11	9			81.8
Western Pacific	27	16	11		1	68.8
<b>Total</b>	<b>194</b>	<b>130</b>	<b>104</b>	<b>8</b>	<b>1</b>	<b>80.0</b>

MS = Member State; \* Includes those MSs that responded not having adopted IMCI and therefore not answering the questions of the survey

**Table 2. Respondents to IMCI Global Implementation Survey by WHO Region**

WHO Region	Questionnaires included in final analysis	Three-team response (MoH/WHO/UNICEF)	Response by MoH/WHO team	Response by MoH alone	Response by WHO alone	Response by WHO/UNICEF team	Unknown
African	41	1	1	22	14	-	3
Americas	15	-	5	9	1	-	-
Eastern Mediterranean	14	3	7	4	-	-	-
Europe	6	1	-	3	1	1	-
South-east Asia	9	-	-	6	3	-	-
Western Pacific	10	-	1	9	-	-	-
<b>Total</b>	<b>95</b>	<b>5</b>	<b>15</b>	<b>52</b>	<b>19</b>	<b>1</b>	<b>3</b>