

Supplemental Tables for
Validation and Refinement of the Age, Comorbidities, and Albumin index in Elderly Patients with Diffuse Large B-Cell Lymphoma: An Effective Tool for Comprehensive Geriatric Assessment
Hui Liu et al.

Table S1. Toxicities after therapy

Grade ≥ 3 toxicities	ACA "excellent" group (score 0, n=39)	ACA "good" group (score 1, n = 31)	ACA "moderate to poor" group (score ≥ 2 , n =29)	<i>P</i>
Hematologic	22 (56.4)	14 (45.2)	18 (64.3)	0.330
Gastrointestinal	1 (2.6)	2 (6.5)	0 (0.0)	0.490
Pulmonary	1 (2.6)	1 (3.2)	0 (0.0)	1.000
Infectious	1 (2.6)	4 (12.9)	9 (32.1)	0.002
Cardiovascular	0 (0.0)	1 (3.2)	2 (7.1)	0.188
Nervous	0 (0.0)	0 (0.0)	1 (3.6)	0.286

ACA index, Age, Comorbidities, and Albumin index
Statistical significance was set at $P < 0.05$.

Table S2. IACA index

Prognostic variables	IACA value		
	0	1	2
IADL scale	8	6 to 7	≤ 5
ACA index	Excellent	Good	Moderate to poor

ACA index, Age, Comorbidities, and Albumin index; IADL, instrumental activities of daily living

* low-risk, score 0; intermediate-risk, score 1 to 2; high-risk, score 3 to 4.