

The role of lipids in the inception, maintenance and complications of dengue virus infection

Carlos Fernando Odir Rodrigues Melo¹, Jeany Delafiori¹, Mohamad Ziad Dabaja¹, Diogo Noin de Oliveira¹, Tatiane Melina Guerreiro¹, Tatiana Elias Colombo², Maurício Lacerda Nogueira², Jose Luiz Proenca-Modena³, Rodrigo Ramos Catharino^{1*}

¹INNOVARE Biomarkers Laboratory, School of Pharmaceutical Sciences, University of Campinas, Brazil.

²School of Medicine from São José do Rio Preto (FAMERP), Brazil.

³Laboratory of Study of Emerging Viruses (LEVE), Department of Genetic, Evolution and Bioagents, Institute of Biology, University of Campinas, Brazil.


* Corresponding author: Rodrigo Ramos Catharino

Supplementary Information #1
MS/MS Reactions of the Species Elected by PCA


MS/MS spectrum of 743 m/z









MS/MS spectrum of 769 m/z



MS/MS spectrum of 795 m/z



MS/MS spectrum of 859 m/z




MS/MS spectrum of 762 m/z




MS/MS spectrum of 784 m/z




MS/MS spectrum of 838 m/z




MS/MS spectrum of 768 m/z



MS/MS spectrum of 770 m/z



MS/MS spectrum of 792 m/z



MS/MS spectrum of 798 m/z

