

S2 Table: Distribution of diet-related characteristics in two arms at baseline

Characteristic	Intervention n = 239 (%)	Control n = 239 (%)	p-value*
<i>1. Access to nearest local market</i>			
Accessible by walk	183 (77%)	189 (79%)	0.618
Not accessible by walk	56 (23%)	50 (21%)	
<i>2. Habit of consuming main meals †</i>			
With family	186 (78%)	202 (85%)	0.276
Alone or with others	53 (22%)	37 (15%)	
<i>3. Type of diet consumed †</i>			
Complete vegetarian	8 (3%)	7 (3%)	0.782
Partial- or non-vegetarian	231 (97%)	232 (97%)	
<i>4. Type of oil used for cooking</i>			
Only coconut oil	216 (90%)	222 (93%)	0.206
More than one type of oil	23 (10%)	17 (7%)	
<i>5. Daily fruit intake of at least 2 servings</i>			
No	236 (99%)	238 (99.6%)	0.327
Yes	3 (1%)	1(0.4%)	
<i>6. Daily vegetable intake of at least 3 servings</i>			
No	201 (84%)	206 (86%)	0.798
Yes	38 (16%)	33 (14%)	
<i>7. Use of locally available fruits</i>			
No	180 (75%)	195 (82%)	0.243
Yes	59 (25%)	44 (18%)	
<i>8. Use of locally available vegetables</i>			
No	204 (85%)	214 (90%)	0.229
Yes	35 (15%)	25 (10%)	

* All p-values are based on weighted samples using chi-square test taking into account cluster design; † Variables related to selected individual in the household.